

## ON THE SEMI-CANONICAL PROPERTY IN THE PRODUCT SPACE $X \times I$

A. OKUYAMA AND Y. YASUI

**ABSTRACT.** As one of the several properties in generalized metric spaces, the semi-canonical property has been discussed from the viewpoint of the extension of mappings. In this paper, that property will be discussed in product space  $X \times I$  and reduced to a property of  $X$ .

**1. Introduction.** By a pair  $(X, A)$  we mean a topological space  $X$  with a closed subset  $A$  of  $X$ . Let  $(X, A)$  be a pair. As in [6], a collection  $\mathcal{V} = \{V_\lambda : \lambda \in \Lambda\}$  of open subsets of  $X$  is called a *semi-canonical cover* for  $(X, A)$  if

(1)  $\bigcup_{\lambda \in \Lambda} V_\lambda = X - A$ , and

(2) for each  $x \in A$  and each neighborhood  $U$  of  $x$  in  $X$  there exists a neighborhood  $W$  of  $x$  in  $X$  such that  $\text{St}(W, \mathcal{V}) \subset U$ , where

$$\text{St}(W, \mathcal{V}) = \bigcup \{V \in \mathcal{V} : V \cap W \neq \emptyset\}$$

denotes the star of  $W$  with respect to  $\mathcal{V}$ .

If a semi-canonical cover exists for  $(X, A)$ ,  $(X, A)$  is called a *semi-canonical pair*.

It was proved by D. Hyman ([6], [7]) that  $(X, A)$  is a semi-canonical pair if  $X$  is the image of a metric space by a closed continuous map. It is also mentioned by M. Cauty [3] that, if  $X$  is a stratifiable space (cf. [2]), then any pair  $(X, A)$  is semi-canonical. However, quite recently S. San-ou [11] pointed out that Cauty's statement was false by constructing an  $M_1$ -space  $X$  (cf. [4]) such that  $(X, A)$  was not semi-canonical for some closed subset  $A$  of  $X$ .

The purpose of this paper is to discuss the semi-canonical property in the product space  $X \times I$  of a  $T_1$  space  $X$  with the unit closed interval  $I$  and to reduce it to a property in  $X$ .

**THEOREM 1.** *Let  $X$  be a  $T_1$  space. Then  $(X \times I, X \times \{0\})$  is a semi-canonical pair if and only if  $X$  is metrizable.*

By Theorem 1 it can be easily seen that, if  $X$  is any nonmetrizable  $M_1$ -space, then  $X \times I$  is an  $M_1$ -space such that  $(X \times I, X \times \{0\})$  is never semi-canonical.

---

Received by the editors November 8, 1976 and, in revised form, April 3, 1977.

AMS (MOS) subject classifications (1970). Primary 54E35; Secondary 54D30.

Key words and phrases. Semi-canonical cover, semi-canonical pair,  $M_1$ -space, metric space, compact-covering map, open map,  $X$ -base, countable character, first countable space.

© American Mathematical Society 1978

**THEOREM 2.** *Let  $X$  be a  $T_1$  space. Then  $(X \times I, K \times \{0\})$  is a semi-canonical pair for each compact subset  $K$  if and only if  $X$  is a regular space which is a compact-covering,<sup>1</sup> open image of a metric space.*

**THEOREM 3.** *Let  $X$  be a  $T_1$  space. Then  $(X \times I, \{(x, 0)\})$  is a semi-canonical pair for each point  $x \in X$  if and only if  $X$  is a regular, first countable space.*

Throughout this paper, the following notations will be used:  $X_0$  and  $X_n$  denote the subspaces  $X \times \{0\}$  and  $X \times \{1/n\}$  of  $X \times I$  for  $n = 1, 2, \dots$ ;  $\pi$  denotes the projection from  $X \times I$  onto  $X$ ; and  $I_n$  denotes the subspace  $[0, 1/n]$  of  $I$  for  $n = 1, 2, \dots$ .

All spaces in this paper are  $T_1$ , and all maps are continuous.

**2. Proof of Theorem 1.** The sufficiency of the condition is clear, since every pair  $(X, A)$  in a metric space  $X$  is semi-canonical (cf. [6]). To prove necessity, suppose that there exists a semi-canonical cover  $\mathcal{V}$  for  $(X \times I, X_0)$ . Put

$$\mathcal{U}_n = \pi(\mathcal{V}|X_n) \quad (= \{\pi(V \cap X_n) : V \in \mathcal{V}\})$$

for  $n = 1, 2, \dots$ . Then  $\{\mathcal{U}_n : n = 1, 2, \dots\}$  is clearly a sequence of open covers of  $X$ .

Let us show that, for each point  $x \in X$ , the system  $\{\text{St}^2(x, \mathcal{U}_n) : n = 1, 2, \dots\}$  forms a neighborhood base at  $x$ , where  $\text{St}^2(x, \mathcal{U})$  denotes the set  $\text{St}(\text{St}(x, \mathcal{U}), \mathcal{U})$ . Then  $X$  is metrizable by a theorem of K. Morita [10]. To complete the proof, let  $x$  be any point of  $X$  and  $G$  an arbitrary neighborhood of  $x$  in  $X$ . Since  $\mathcal{V}$  is a semi-canonical cover for  $(X \times I, X_0)$ , there exist a neighborhood  $H_1$  of  $x$  in  $X$  and a positive integer  $m$  such that  $\text{St}(H_1 \times I_m, \mathcal{V}) \subset G \times I$  holds. Again, for the neighborhood  $H_1$  of  $x$  there exist a neighborhood  $H_2$  of  $x$  in  $X$  and a positive integer  $n$  such that  $n \geq m$  and  $\text{St}(H_2 \times I_n, \mathcal{V}) \subset H_1 \times I$ . Now, let us show  $\text{St}^2(x, \mathcal{U}_n) \subset G$ . Pick an arbitrary point  $y$  in  $\text{St}^2(x, \mathcal{U}_n)$ . Then there are two members  $U, U'$  of  $\mathcal{U}_n$  with  $x \in U, y \in U'$  and  $U \cap U' \neq \emptyset$ . Let  $z$  be a point of  $U \cap U'$ . By the definition of  $\mathcal{U}_n$  there exist  $V, V'$  in  $\mathcal{V}$  such that

$$U = \pi(V \cap X_n) \quad \text{and} \quad U' = \pi(V' \cap X_n).$$

Hence,  $(x, 1/n) \in V, (z, 1/n) \in V \cap V'$  and  $(y, 1/n) \in V'$  hold. The first inclusion  $(x, 1/n) \in V$  implies  $V \subset H_1 \times I$ , because  $(x, 1/n)$  belongs to  $H_2 \times I_n$ ; the second one implies  $V' \subset G \times I$ , because  $(z, 1/n) \in V$  shows  $(z, 1/n) \in H_1 \times I_n$  and  $(z, 1/n) \in V'$  yields  $V' \cap (H_1 \times I_n) \neq \emptyset$ ; and, as a consequence, the last inclusion  $(y, 1/n) \in V'$  implies  $y \in G$ , which completes the proof.

### 3. Some lemmas.

**LEMMA 1.** *Let  $X$  be a space. If the pair  $(X \times I, \{(x, 0)\})$  is semi-canonical for every point  $x \in X$ , then  $X$  is a regular space.*

<sup>1</sup> A continuous map  $f: X \rightarrow Y$  is called *compact-covering* if every compact subset of  $Y$  is the image of some compact subset of  $X$ .

PROOF. Using the same notations as in the proof of Theorem 1, it has been shown that, for a given point  $x$  of  $X$  and an arbitrary neighborhood  $G$  of  $x$ , there exists an open cover  $\mathcal{U}_n$  of  $X$  such that  $\text{St}^2(X, \mathcal{U}_n) \subset G$  holds. Clearly,  $\text{St}(x, \mathcal{U}_n)$  is a neighborhood of  $x$ , whose closure is contained in  $\text{St}^2(x, \mathcal{U}_n)$  and hence in  $G$ . This proves that  $X$  is a regular space.

If  $A \subset X$ , then an  $X$ -base for  $A$  is a collection  $\mathcal{U}$  of open subsets of  $X$  such that, if  $x \in A$  and  $V$  is a neighborhood of  $x$  in  $X$ , then  $x \in U \subset V$  for some  $U \in \mathcal{U}$ .

LEMMA 2. Let  $X$  be a regular ( $T_1$ ) space and  $K$  a compact subset of  $X$ . If there exists a countable  $X$ -base for  $K$ , then there exists an  $X$ -base  $\bigcup_{n=1}^{\infty} \mathcal{P}_n$  for  $K$  such that

- (1)  $\mathcal{P}_n$  is a finite collection whose union covers  $K$  for  $n = 1, 2, \dots$ ,
- (2)  $\{P: P \in \mathcal{P}_{n+1}\}$  refines  $\mathcal{P}_n$  for  $n = 1, 2, \dots$ , and
- (3) for each point  $x$  of  $K$  and each neighborhood  $G$  of  $x$  in  $X$ , there exist a positive integer  $n$  and a neighborhood  $H$  of  $x$  in  $X$  such that  $\text{St}(H, \mathcal{P}_n) \subset G$ .

PROOF. Let  $\mathfrak{B}$  be the given countable  $X$ -base for  $K$ . Since  $\mathfrak{B}|K$  is a countable base for  $K$  itself,  $K$  is metrizable. Hence, for any subset  $E$  of  $K$ , the diameter  $\delta(E)$  of  $E$  is well defined and also, for any cover  $\mathfrak{C}$  of  $K$ , the mesh  $\bar{\mathfrak{C}} = \sup\{\delta(E); E \in \mathfrak{C}\}$  is well defined.

For each  $n$ , let  $\mathcal{U}_n$  be a finite subcollection of  $\mathfrak{B}$  such that

- (1) $_n$   $\mathcal{U}_n$  covers  $K$ , and
- (2) $_n$   $\text{mesh } \mathcal{U}_n|K < 1/2^n$ .

Let  $\{\mathcal{V}_n: n = 1, 2, \dots\}$  be the set of all finite subcollections of  $\mathfrak{B}$ , each of which forms a minimal cover with respect to  $K$ ; that is, any proper subcollection of  $\mathcal{V}_n$  does not cover  $K$  for  $n = 1, 2, \dots$ . Put  $\mathcal{W}_1 = \mathcal{U}_1 \wedge \mathcal{V}_1$  ( $= \{U \cap V: U \in \mathcal{U}_1, V \in \mathcal{V}_1\}$ ) and  $\mathcal{W}_{n+1} = \mathcal{W}_n \wedge \mathcal{U}_{n+1} \wedge \mathcal{V}_{n+1}$  for  $n = 1, 2, \dots$ . Then each  $\mathcal{W}_n$  is a finite collection of open subsets of  $X$  whose union covers  $K$ .

Next, by induction on  $n$ , let us construct a finite collection  $\mathcal{F}_n$  of closed subsets of  $K$ , a finite collection  $\mathcal{P}_n$  of open subsets of  $X$  and a function  $\varphi_n$  from  $\mathcal{F}_n$  onto  $\mathcal{P}_n$  such that the following conditions are satisfied:

- (3) $_n$   $\mathcal{F}_n$  is a closed cover of  $K$  which refines  $\mathcal{W}_n \wedge \mathcal{P}_{n-1}$ , where  $\mathcal{P}_0 = \{X\}$ ,
- (4) $_n$   $\mathcal{P}_n$  refines  $\mathcal{W}_n \wedge \mathcal{P}_{n-1}$ ,
- (5) $_n$  if  $F \in \mathcal{F}_n$ , then  $F \subset \varphi_n(F)$ ,
- (6) $_n$  if  $F \in \mathcal{F}_n$  and  $F \subset O \in \bigcup_{i=1}^{n-1} \mathcal{P}_i \cup \bigcup_{i=1}^n (\mathcal{U}_i \cup \mathcal{V}_i)$ , then  $\overline{\varphi_n(F)} \subset O$ , and

- (7) $_n$  if  $F \cap F' = \emptyset$ , then  $\varphi_n(F) \cap \varphi_n(F') = \emptyset$  for  $F, F' \in \mathcal{F}_n$ .

Let  $\mathcal{W}_1 = \{W_1, \dots, W_k\}$ . Since  $\mathcal{W}_1$  covers  $K$  and  $K$  is normal, there exists a closed cover  $\mathcal{F}_1 = \{F_1, \dots, F_k\}$  of  $K$  such that  $F_i \subset W_i$  for  $i = 1, \dots, k$ . Hence  $\mathcal{F}_1$  satisfies condition (3) $_1$ . Since  $X$  is regular and  $\mathcal{F}_1$  is a finite collection, each member of which is compact, and since  $\mathcal{U}_1$  and  $\mathcal{V}_1$  are also finite collections, it is easy to see that the function  $\varphi_1$  and  $\mathcal{P}_1 = \varphi_1(\mathcal{F}_1)$  are well defined to satisfy conditions (4) $_1$ –(7) $_1$ , as well. The situation in each step

is the same as above, and thus  $\mathcal{F}_n$ ,  $\varphi_n$  and  $\mathcal{P}_n$  are all constructed quite similarly.

Now, it remains to show that the sequence  $\{\mathcal{P}_n: n = 1, 2, \dots\}$  is the required one in Lemma 2. Since  $\mathcal{P}_n$  is finite and satisfies  $(3)_n$  and  $(5)_n$ ,  $\mathcal{P}_n$  satisfies the condition (1). By  $(3)_n$  and  $(6)_n$ ,  $\mathcal{P}_n$  satisfies the condition (2). To prove that  $\{\mathcal{P}_n: n = 1, 2, \dots\}$  satisfies the condition (3), let  $x$  be any point of  $K$  and  $G$  an arbitrary neighborhood of  $x$  in  $X$ . Since  $\mathcal{B}$  is an  $X$ -base for  $K$ , there exists a  $B_0 \in \mathcal{B}$  such that  $x \in B_0 \subset G$ .<sup>2</sup> Let  $\mathcal{V}$  be a finite subcollection of  $\mathcal{B}$  which is a minimal cover with respect to  $K$  and which keeps  $B_0$  as the only member of  $\mathcal{V}$  containing  $x$ . Since  $K$  is a compact  $T_2$  space and since  $\mathcal{B}$  is an  $X$ -base for  $K$ , such  $\mathcal{V}$  certainly exists; further, for some  $n$ ,  $\mathcal{V} = \mathcal{V}_n$ .

Let  $F_0 \in \mathcal{F}_n$  be a member with  $x \in F_0$ . Then  $F_0 \subset B_0$  holds, because  $\mathcal{F}_n$  refines  $\mathcal{W}_n$  which refines  $\mathcal{V}_n$  and  $B_0$  is the only member of  $\mathcal{V}$  containing  $x$ ; and also, by  $(5)_n$  and  $(6)_n$ , the inclusions  $F_0 \subset \varphi_n(F_0) \subset B_0$  hold. Since  $\varphi_n(F_0)$  is an open set containing  $x$ , there exists a positive integer  $m$  such that  $m \geq n$  and  $d(x, K - \varphi_n(F_0)) > 1/2^m$ , where  $d$  denotes the metric function on  $K$ . Since  $\mathcal{F}_{m+1}$  is a cover of  $K$  by  $(3)_{m+1}$ , there exists an  $F_1 \in \mathcal{F}_{m+1}$  containing  $x$ . To complete the proof, it suffices to show that

$$\text{St}(\varphi_{m+1}(F_1), \mathcal{P}_{m+1}) \subset \varphi_n(F_0),$$

because  $\varphi_{m+1}(F_1)$  is an open set in  $X$  containing  $x$  and  $\varphi_n(F_0)$  is contained in  $B_0$ , which is contained in  $G$ . Let  $P$  be an arbitrary member of  $\mathcal{P}_{m+1}$  and  $F$  the corresponding member of  $\mathcal{F}_{m+1}$  by  $P = \varphi_{m+1}(F)$ . If  $P \cap \varphi_{m+1}(F_1) \neq \emptyset$ , then by  $(7)_{m+1}$ ,  $F \cap F_1 \neq \emptyset$ . Since  $\mathcal{F}_{m+1}$  refines  $\mathcal{P}_{m+1}$  by  $(5)_{m+1}$  and  $\mathcal{P}_{m+1}$  refines  $\mathcal{W}_{m+1}$  by  $(4)_{m+1}$ , and since  $\mathcal{W}_{m+1}$  refines  $\mathcal{U}_{m+1}$  whose mesh restricting to  $K$  is less than  $1/2^{m+1}$ , the diameter  $\delta(F \cup F_1)$  is less than  $1/2^m$ . Since  $x$  belongs to  $F_1$ , by the choice of  $m$ ,  $F \cup F_1 \subset \varphi_n(F_0)$  holds. Again by  $(6)_{m+1}$ ,  $\varphi_{m+1}(F) \subset \varphi_n(F_0)$  and thus  $P \subset \varphi_n(F_0)$  holds, which completes the proof.

LEMMA 3. *Let  $X$  be a regular  $(T_1)$  space and  $K$  a compact subset of  $X$ . If there exists a countable  $X$ -base for  $K$ , then  $(X, K)$  is a semi-canonical pair.*

PROOF. Let  $\bigcup_{n=1}^{\infty} \mathcal{P}_n$  be an  $X$ -base for  $K$  obtained by Lemma 2. For each  $n$ , put  $G_n = \bigcup \{P: P \in \mathcal{P}_n\}$ . Then, by conditions (1) and (2) in Lemma 2,  $\overline{G}_{n+1} \subset G_n$  for  $n = 1, 2, \dots$  and  $K \subset \bigcap_{n=1}^{\infty} G_n$ , and by condition (3) and by the fact that  $K$  is compact, it is easily seen that  $K = \bigcap_{n=1}^{\infty} G_n$ .

Now, put  $\mathcal{V}_0 = \{X - \overline{G}_2\}$  and  $\mathcal{V}_n = \mathcal{P}_n | (G_n - \overline{G}_{n+2})$  for  $n = 1, 2, \dots$ , and put  $\mathcal{V} = \bigcup_{n=0}^{\infty} \mathcal{V}_n$ . Then it will be shown that  $\mathcal{V}$  is a semi-canonical cover for  $(X, K)$ . Clearly,  $\mathcal{V}$  is an open cover of  $X - K$ . To complete the proof, let  $x$  be any point of  $K$  and  $U$  an arbitrary neighborhood of  $x$  in  $X$ . By condition (3) in Lemma 2, there exist a positive integer  $n$  and a neighborhood  $H$  of  $x$  in  $X$  such that  $\text{St}(H, \mathcal{P}_n) \subset U$ . Put  $W = H \cap G_{n+1}$ . Then  $W$  is a neighborhood of  $x$  in  $X$  such that  $W \cap V = \emptyset$  for each  $V \in \bigcup_{i=1}^n \mathcal{V}_i$ .

<sup>2</sup> If  $K$  is singleton, then  $\bigcup_{n=1}^{\infty} \mathcal{P}_n$  is easily chosen from the given countable  $X$ -base for  $K$ , because  $X$  is regular. So, assuming that  $K$  is not a singleton,  $B_0$  is picked out from  $\mathcal{B}$  such that  $K - B_0 \neq \emptyset$ .

Therefore

$$\text{St}(W, \mathcal{V}) = \text{St}\left(W, \bigcup_{i>n} \mathcal{V}_i\right) \subset \text{St}\left(W, \bigcup_{i>n} \mathcal{P}_i\right) \subset \text{St}(H, \mathcal{P}_n) \subset U$$

by condition (2) in Lemma 2, and that completes the proof.

**4. Proofs of Theorems 2 and 3.** The following characterization of the compact-covering open images of metric spaces, due to E. Michael and K. Nagami [9] will be used in the proof of Theorem 2.

**THEOREM M-N (E. MICHAEL AND K. NAGAMI).**<sup>3</sup> For a  $T_2$  space  $X$ , the following conditions are equivalent:

- (1)  $X$  is the compact-covering open image of a metric space.
- (2) Every compact subset of  $X$  is metrizable and of countable character in  $X$ .<sup>4</sup>
- (3) Every compact subset of  $X$  has a countable  $X$ -base.

**PROOF OF THEOREM 2. Necessity.** Let  $(X \times I, K \times \{0\})$  be a semi-canonical pair for any compact subset  $K$  of  $X$ . Then  $X$  is a regular space by Lemma 1 putting  $K$  in the assumption a singleton. Next, it will be shown that each compact subset  $K$  of  $X$  has a countable  $X$ -base. Then  $X$  is the compact-covering open image of a metric space by Theorem M-N.

To complete the proof, let  $K$  be a compact subset of  $X$ . By the assumption, there exists a semi-canonical cover  $\mathcal{V}$  for  $(X \times I, K \times \{0\})$ . Put  $\mathcal{V}_n$  the finite subcollection of  $\mathcal{V}$  which covers  $K \times \{1/n\}$ , and put  $\mathcal{U}_n = \pi(\mathcal{V}_n|X_n)$  for  $n = 1, 2, \dots$ .

Then it is easy to show that the collection  $\bigcup_{n=1}^{\infty} \mathcal{U}_n$  is the required  $X$ -base for  $K$ , by the same technique as in the proof of Theorem 1.

**Sufficiency.** It is easy to check that, if  $X$  is the compact-covering open image of a metric space, then so is  $X \times I$ . Hence, for any compact subset  $K$  of  $X$ ,  $K \times \{0\}$  has a countable  $X \times I$ -base by Theorem M-N, and thus  $(X \times I, K \times \{0\})$  is a semi-canonical pair by Lemma 3, which completes the proof.

**PROOF OF THEOREM 3. Necessity.** By Lemma 1,  $X$  is a regular space. The first countability of  $X$  is proved by the same technique as in the proof of the necessity in Theorem 2, replacing  $K$  by a singleton.

**Sufficiency.** If  $X$  is a regular  $(T_1)$  first countable space, then so is  $X \times I$ . In general, it is easily seen that, in any regular  $(T_1)$  first countable space  $Y$ , the pair  $(Y, \{y\})$  is always semi-canonical for each point  $y \in Y$ . This completes the proof.

**5. Comments.** 1. From the proofs of Theorems 1, 2 and 3, it is easy to see that, in the conditions of these theorems, the closed interval  $I$  may be

<sup>3</sup> The fact (2)  $\rightarrow$  (3) was proved by M. M. Čoban [5]; for completely regular space  $X$ , it had previously been obtained by A. V. Arhangel'skii [1].

<sup>4</sup> A set  $K \subset X$  is of countable character in  $X$  if there is a countable outer base  $\{U_n: n = 1, 2, \dots\}$  at  $K$  in  $X$  (i.e. each  $U_n$  is open and contains  $K$ , and every open set containing  $K$  contains some  $U_n$ ) (cf. [9]).

replaced by any space containing a convergent sequence. By such replacement in Theorem 1, one obtains a slight modification of the proof of the following theorem due to D. M. Hyman [7], remembering two facts: (1) The closed image of a metric space is a Fréchet-Urysohn space (cf. [8]); and (2) any pair  $(X, A)$  is semi-canonical if  $X$  is the closed image of a metric space (cf. [7]).

**THEOREM (D. HYMAN).** *If  $X$  and  $Y$  are nondiscrete spaces and if  $X \times Y$  is the closed image of a metric space, then  $X$  and  $Y$  are metrizable.*

2. The semi-canonical property need not be two-productive. For example, let  $X = N \cup \{p\}$  be a subspace of Stone-Čech compactification  $\beta N$  of  $N$  ( $= \{1, 2, \dots\}$ ) with  $p \in \beta N - N$ . Then it is well known that  $X$  is not first countable at  $p$ , and thus  $(X \times I, \{(p, 0)\})$  is not semi-canonical by Theorem 3. However, it is easy to see that any pair  $(X, A)$  is always semi-canonical.

This example also shows that, in the conditions of Theorems 1 and 2,  $X \times I$  cannot be replaced by  $X$ . Clearly, then, the semi-canonical property in  $X$  is very different from the semi-canonical property in  $X \times I$ .

#### REFERENCES

1. A. V. Arhangel'skiĭ, *Bicomact sets and the topology of spaces*, Trudy Moskov. Mat. Obšč. **13** (1965), 3–55 = Trans. Moscow Math. Soc. **13** (1965), 1–62.
2. C. J. R. Borges, *On stratifiable spaces*, Pacific J. Math. **17** (1966), 1–16.
3. M. R. Cauty, *Une généralisation du théorème de Borsuk-Whitehead-Hanner aux espaces stratifiables*, C. R. Acad. Sci. Paris Sér. A-B **275** (1972), A271–A274.
4. J. G. Ceder, *Some generalizations of a metric space*, Pacific J. Math. **11** (1961), 105–126.
5. M. M. Čoban, *Perfect mappings and spaces of countable type*, Vestnik Moskov. Univ. Ser. I Mat. Meh. **22** (1967), no. 6, 87–93.
6. D. M. Hyman, *A generalization of the Borsuk-Whitehead-Hanner theorem*, Pacific J. Math. **23** (1967), 263–271.
7. D. M. Hyman, *A note on closed maps and metrizability*, Proc. Amer. Math. Soc. **21** (1969), 109–112.
8. N. Lasnev, *Closed images of metric spaces*, Dokl. Akad. Nauk SSSR **170** (1966) = Soviet Math. Dokl. **7** (1966), 1219–1221.
9. E. Michael and K. Nagami, *Compact-covering images of metric spaces*, Proc. Amer. Math. Soc. **37** (1973), 260–266.
10. K. Morita, *On the simple extension of a space with respect to a uniformity*. IV, Proc. Japan Acad. **27** (1951), 632–636.
11. S. San-ou, *A note on  $\Xi$ -product*, J. Math. Soc. Japan **29** (1977), 281–285.

DEPARTMENT OF MATHEMATICS, OSAKA KYOIKU UNIVERSITY, TENNOJI, OSAKA, JAPAN