

k -REGULAR MAPPINGS OF 2^n -DIMENSIONAL EUCLIDEAN SPACE

MICHAEL E. CHISHOLM

ABSTRACT. A map $f: X \rightarrow R^n$ is said to be k -regular if whenever x_1, \dots, x_k are distinct points of X , then $f(x_1), \dots, f(x_k)$ are linearly independent. Using configuration spaces and homological methods, it is shown that there does not exist a k -regular map from R^n into $R^{n(k-\alpha(k))+\alpha(k)-1}$ where $\alpha(k)$ denotes the number of ones in the dyadic expansion of k and n is a power of 2.

A continuous map $f: X \rightarrow R^n$ is said to be k -regular if whenever x_1, \dots, x_k are distinct elements of X , then $f(x_1), \dots, f(x_k)$ are linearly independent. The study of k -regular maps is prompted by the theory of Čebyšev approximation. The reader is referred to [12, pp. 237–242] for the relationship between these two concepts.

Results on existence and nonexistence of k -regular maps can be found in [1], [2], [4]–[7]. In [4], David Handel and Fred Cohen, using algebraic-topological methods, obtained a nonexistence theorem about k -regular mappings of the plane. The object of the present paper is to generalize their result to k -regular mappings of R^n where n is a power of 2. We obtain an improvement upon the following result, for the case n a power of 2.

THEOREM 1 (BOLTJANSKIĬ-RYŠKOV-ŠAŠKIN). *If a $2k$ -regular map of R^n into R^n exists, then $N \geq (n+1)k$.*

I wish to thank David Handel for his help and advice while writing this paper and throughout my graduate study.

THEOREM 2. *There does not exist a k -regular map of R^n into $n(k - \alpha(k)) + \alpha(k) - 1$ dimensional Euclidean space where $\alpha(k)$ denotes the number of ones in the dyadic expansion of k and n is a power of 2.*

In the proof we utilize algebraic-topological properties of the configuration space of R^n , denoted $F(R^n, k)$, is the subspace of $(R^n)^k$ consisting of ordered k -tuples of distinct points in R^n . The symmetric group Σ_k acts freely on $F(R^n, k)$ and orthogonally on R^k by permuting factors. Let $P_{n,k}$ denote the k -plane bundle $F(R^n, k) \times_{\Sigma_k} R^k \rightarrow F(R^n, k)/\Sigma_k$.

LEMMA 3. *If n and k are powers of 2, then $\bar{\omega}_{(n-1)(k-1)}(P_{n,k}) \neq 0$.*

Received by the editors March 21, 1978.

AMS (MOS) subject classifications (1970). Primary 55G99; Secondary 41A50.

Key words and phrases. k -regular maps, configuration spaces, Dyer-Lashof operations, Stiefel-Whitney classes.

© 1979 American Mathematical Society
0002-9939/79/0000-0183/\$02.00

PROOF. All homology and cohomology groups are with \mathbb{Z}_2 coefficients. Let \mathcal{C}_n denote the little n -cubes operad with associated monad C_n as constructed by J. P. May in [8]. By [8, Theorem 4.8], $F(R^n, k)$ is Σ_k -equivariantly homotopy equivalent to $\mathcal{C}_n(k)$. So, as in [4] replace $F(R^n, k)/\Sigma_k$ by $\mathcal{C}_n(k)/\Sigma_k$ in $P_{n,k}$.

The following composition is a classifying map for $P_{n,k}$;

$$\mathcal{C}_n(k)/\Sigma_k \xrightarrow{\sigma_k} \mathcal{C}_\infty(k)/\Sigma_k \cong B\Sigma_k \xrightarrow{\rho} BO(k)$$

where ρ is induced from the regular representation $\Sigma_k \rightarrow O(k)$ and σ_k is the direct limit of $\sigma_{m,k}$ where $\sigma_{m,k}$ is given in [8, p. 31].

Let $[1]$ denote the element of $H_0(S^0)$ determined by the nonbase point of S^0 . By [3, §3], $H_*(C_n S^0)$ is given in terms of the Dyer-Lashof operations on $[1]$.

Suppose $k = 2^r$, then $I = (2^{r-1}(n-1), \dots, 2(n-1), n-1)$ is an admissible sequence of degree $(n-1)(k-1)$ and excess $n-1$. By [3, §§1.4], $Q^I[1]$ is an element of $H_*(C_n S^0)$ and by filtration arguments given there it follows that $Q^I[1] \in H_*(\mathcal{C}_n(k)/\Sigma_k)$. Since σ_k is the restriction of a map of \mathcal{C}_n -spaces, we have

$$\sigma_k^*(Q^I[1]) = Q^I[1].$$

Thus it suffices to show that $\langle \rho^* \bar{\omega}_{(n-1)(k-1)}, Q^I[1] \rangle \neq 0$, where $\langle \cdot, \cdot \rangle$ denotes the Kronecker index and $\bar{\omega}_{(n-1)(k-1)}$ is the $(n-1)(k-1)$ -universal dual Stiefel-Whitney class.

As a first step we now show

$$\langle \rho^* \bar{\omega}_{(n-1)(k-1)}, Q^I[1] \rangle = \langle \rho^* \omega_{k-1}^{n-1}, Q^I[1] \rangle.$$

By [10, p. 220], $\rho^* \bar{\omega}_{(n-1)(k-1)} = \rho^*(\sum_j c_j \omega_1^{j_1} \cdots \omega_{k-1}^{j_{k-1}})$ where j runs over all (j_1, \dots, j_{k-1}) with $j_i \geq 0$, $\sum_{i=1}^{k-1} j_i = (n-1)(k-1)$ and $c_j = (j_1 + \cdots + j_{k-1})! / j_1! \cdots j_{k-1}!$, $\rho^* \omega_k = 0$ since every k -plane bundle with structural group Σ_k admits a nowhere zero section.

Suppose $1 \leq i < k-1$ is odd, $k = 2^r$, $u \in H^* B\Sigma_k$, $L = (s_1, \dots, s_r)$ and $\psi(Q^L[1]) = \Sigma Q^A[1] \otimes Q^B[1]$ where ψ is the diagonal map in homology. See [9, p. 6]. Thus

$$\langle \rho^* \omega_i u, Q^L[1] \rangle = \sum \langle \rho^* \omega_i, Q^A[1] \rangle \langle \rho^* u, Q^B[1] \rangle = 0.$$

This follows since for each A , the length of A is r and $Q^A = \Sigma Q^M$ where each M is admissible and of the same degree and length as A , and since we may assume the degree of each A is i . Hence each A has positive excess and by [11, Theorem 4.7], $\langle \rho^* \omega_i, Q^A[1] \rangle = 0$.

Thus

$$\langle \rho^* \bar{\omega}_{(n-1)(k-1)}, Q^I[1] \rangle = \sum_j \langle c_j \rho^* \omega_2^{j_2} \omega_4^{j_4} \cdots \omega_{k-3}^{j_{k-3}} \omega_{k-1}^{j_{k-1}}, Q^I[1] \rangle$$

where $j = (j_2, j_4, \dots, j_{k-2}, j_{k-1})$ and $2j_2 + 4j_4 + \cdots + (k-2)j_{k-2} + (k-1)j_{k-1} = (n-1)(k-1)$.

For such j we now show $c_j \equiv 0 \pmod{2}$ if $j_{k-1} \neq n-1$. Since j_{k-1} is odd, $j_{k-1} = 2^m S - 1$, S odd. Suppose $j_{k-1} \neq n-1$. Thus j_{k-1} can be written as $2^0 + 2^1 + \dots + 2^{m-1} + \text{other distinct powers of } 2$. Let $2 \leq i \leq k-2$ be even. If $j_i \neq 0$, then j_i can be written as $2^p + \text{other distinct powers of } 2$ where p is minimal in this expression of j_i . By writing c_j as a product of binomial coefficients, $c_j = y(j_i + j_{k-1})! / j_i! j_{k-1}!$, for some integer y . Thus if $p \leq m-1$, then $c_j \equiv 0 \pmod{2}$. We may thus assume that j_i is divisible by 2^m for i even. So we have;

$$2j_2 + \dots + (k-2)j_{k-2} = 2^m(k-1)(n' - s) \quad (1)$$

where $n' = n/2^m$ is even. Since the L.H.S. of (1) is divisible by 2^{m+1} and the R.H.S. of (1) is not divisible by 2^{m+1} we have a contradiction. Hence

$$\langle \rho^* \bar{\omega}_{(n-1)(k-1)}, Q^I[1] \rangle = \langle \rho^* \omega_{k-1}^{n-1}, Q^I[1] \rangle.$$

We now show that $\langle \rho^* \omega_{k-1}^m, Q^{mJ}[1] \rangle = 1$ where $mJ = (2^{r-1}m, \dots, 2m, m)$, $m \geq 1$, $k = 2^r$, concluding the proof of the lemma. It is easily seen, by induction on r , that $1J = J$ is the only admissible sequence of length r and degree $2^r - 1$ such that $Q^J \neq 0$. Next, using induction and the internal Cartan formula [9, p. 6], we see that if L is an admissible sequence of length r and degree $2^r - 1$, then $Q^L = 0$. By [11, Theorem 4.7], the diagonal Cartan formula [9, p. 6] and induction on m , $\langle \rho^* \omega_{k-1}^m, Q^{mJ}[1] \rangle = 1$.

In [4], the following result is proved.

THEOREM 4 (HANDEL-COHEN). *If a k -regular map of X into R^N exists, then $F(X, k) \times_{\Sigma_k} R^k \rightarrow F(X, k)/\Sigma_k$ admits an N - k -plane inverse.*

PROOF OF THEOREM 2. By Theorem 4, it suffices to show $\bar{\omega}_{(n-1)(k-\alpha(k))}(P_{n,k}) \neq 0$. Write $k = \sum_{i=1}^{\alpha(k)} j(i)$ where $j(i) = 2^{m(i)}$, $m(1) < m(2) < \dots < m(\alpha(k))$. We have a map of k -plane bundles

$$f: P_{n,j(1)} \times \dots \times P_{n,j(\alpha(k))} \rightarrow P_{n,k}$$

as follows: Choose pairwise disjoint open discs $E_1, \dots, E_{\alpha(k)}$ in R^n . Then we can regard $P_{n,j(i)}$ as $F(E_i, j(i)) \times_{\Sigma_{j(i)}} R^{j(i)} \rightarrow F(E_i, j(i))/\Sigma_{j(i)}$. Define f by

$$f((x_1, v_1), \dots, (x_{\alpha(k)}, v_{\alpha(k)})) = (x_1, \dots, x_{\alpha(k)}; v_1, \dots, v_{\alpha(k)}), \\ (x_i, v_i) \in F(E_i, j(i)) \times_{\Sigma_{j(i)}} R^{j(i)}.$$

Thus

$$f^* \bar{\omega}_{(n-1)(k-\alpha(k))}(P_{n,k}) = \bar{\omega}_{(n-1)(k-\alpha(k))}(P_{n,j(1)} \times \dots \times P_{n,j(\alpha(k))}),$$

which has as a nonzero component, by Lemma 3,

$$\bar{\omega}_{(n-1)(j(1)-1)}(P_{n,j(1)}) \times \dots \times \bar{\omega}_{(n-1)(j(\alpha(k))-1)}(P_{n,j(\alpha(k))}).$$

Thus $f^* \bar{\omega}_{(n-1)(k-\alpha(k))}(P_{n,k}) \neq 0$.

REFERENCES

1. V. G. Boltjanskii, S. S. Ryškov and Ju. A. Šaškin, *On k -regular imbeddings and their application to the theory of approximation of functions*, Uspehi Mat. Nauk **15** (1960), no. 6 (96), 125–132; Amer. Math. Soc. Transl. (2) **28** (1963), 211–219.
2. K. Borsuk, *On the k -independent subsets of the Euclidean space and of the Hilbert space*, Bull. Acad. Polon. Sci. Cl. III **5** (1957), 351–356.
3. F. R. Cohen, *The homology of \mathcal{C}_{n+1} -spaces, $n > 0$* , Lecture Notes in Math., vol. 533, Springer-Verlag, Berlin and New York, 1967, pp. 207–351.
4. F. R. Cohen and D. Handel, *k -regular embeddings of the plane*, Proc. Amer. Math. Soc. **72** (1978), 202–204.
5. D. Handel, *Obstructions to 3-regular embeddings*, Houston Math. J. (submitted).
6. ———, *Some existence and non-existence theorems for k -regular maps*, Fund. Math. (submitted).
7. D. Handel and J. Segal, *On k -regular embeddings of spaces in Euclidean space*, Fund. Math. (to appear).
8. J. P. May, *The geometry of iterated loop spaces*, Lecture Notes in Math., vol. 271, Springer-Verlag, Berlin and New York, 1972.
9. ———, *The homology of E_∞ -spaces*, Lecture Notes in Math., vol. 533, Springer-Verlag, Berlin and New York, 1967, pp. 1–68.
10. J. W. Milnor and J. D. Stasheff, *Characteristic classes*, Ann. of Math. Studies, no. 76, Princeton Univ. Press, Princeton, N.J., 1974.
11. S. Priddy, *Dyer-Lashof operations for the classifying spaces of certain matrix groups*, Quart. J. Math. Oxford Ser. (3) **26** (1975), 179–193.
12. I. Singer, *Best approximation in normed linear spaces by elements of linear subspaces*, Springer-Verlag, Berlin and New York, 1970.

DEPARTMENT OF MATHEMATICS, WAYNE STATE UNIVERSITY, DETROIT, MICHIGAN 48202