

## A CYCLE IS THE FUNDAMENTAL CLASS OF AN EULER SPACE

R. BENEDETTI AND M. DEDÒ<sup>1</sup>

**ABSTRACT.** We prove that every cycle in a closed P.L. manifold  $M$  can be regarded as the fundamental class of an Euler subpolyhedron of  $M$ .

Let  $V$  be a compact real analytic manifold without boundary. It is a long-standing problem to see which  $(\mathbf{Z}_2)$  homology classes of  $V$  can be represented as the fundamental class of an analytic subset of  $V$  (and, in fact, it is conjectured that this is true for any homology class). The analogous problem arises with real algebraic manifolds, although in this case the general statement is false (even if  $V$  is connected; see, for instance, [BT]).

D. Sullivan (in [S]) observed that every real analytic set can be regarded as an Euler space (see definition below); it is then natural to ask, first of all, if it is true that every homology class of a closed P.L. manifold  $M$  can be represented as the fundamental class of an Euler subpolyhedron of  $M$ .

In this note we prove that this in fact happens: actually, we give a construction to add lower-dimensional simplexes to a cycle in  $M$  until we get an Euler space (in  $M$ ).

The techniques used are entirely elementary and involve merely P.L. transversality (as stated for example in [RS]) and combinatorial results on Euler spaces (see [A]).

We shall work in the P.L. category. For notations and definitions we refer to [RS]. All cycles and manifolds are intended unoriented and compact.

By an  $n$ -cycle  $P$  we mean a polyhedron  $P = |K|$  such that

- (1)  $n = \max\{\dim A, \text{ for } A \text{ a simplex of } K\}$ ,
- (2) each  $(n - 1)$ -simplex of  $K$  is the face of an even number of simplexes of  $K$ .

By an  $n$ -cycle  $P$  with boundary  $\partial P$  we mean a pair of polyhedra  $(P, \partial P) = |K, \partial K|$  such that (1)  $n = \max\{\dim A, \text{ for } A \text{ a simplex of } K\}$ , (2)  $\partial P$  is an  $(n - 1)$ -cycle, (3) each  $(n - 1)$ -simplex of  $K \setminus \partial K$  is the face of an even number of  $n$ -simplexes of  $K$ , (4) each  $(n - 1)$ -simplex of  $\partial K$  is the face of an odd number of  $n$ -simplexes of  $K$ . A cycle (with boundary) in  $M$  is a subpolyhedron of  $M$  which is a cycle (with boundary).

A closed (P.L.) manifold is a compact (P.L.) manifold without boundary.

An Euler space is a polyhedron  $P$  such that, for each  $x \in P$ ,  $\chi(\text{lk}(x, P)) \equiv 0 \pmod{2}$ .

---

Received by the editors October 13, 1981 and, in revised form, March 11, 1982.  
1980 *Mathematics Subject Classification.* Primary 57Q99; Secondary 57R95.

<sup>1</sup>The authors are members of the G.N.S.A.G.A.

An Euler pair is a pair of polyhedra  $(P, Q)$  such that (1)  $\forall x \in P \setminus Q, \chi(\text{lk}(x, P)) \equiv 0 \pmod{2}$ ; (2)  $\forall x \in Q, \chi(\text{lk}(x, Q)) \equiv 0 \pmod{2}$ ; (3)  $\forall x \in Q, \chi(\text{lk}(x, P)) \equiv 1 \pmod{2}$ .

REMARKS. (1) An Euler space is a cycle (without boundary).

(2) An Euler pair  $(P, Q)$  is not, in general, a cycle with boundary (if  $\dim P = n, Q$  may not be of dimension  $n - 1$ ).

(3) Note that the definition of an  $n$ -cycle is slightly different from the usual one which requires also each simplex of  $K$  to be the face of an  $n$ -simplex of  $K$ . However, a cycle as we defined it naturally carries a fundamental class (which is a cycle in the usual sense) as follows:

Let  $P = |K|$  be an  $n$ -cycle. The fundamental class  $\tilde{P}$  of  $P$  is the polyhedron obtained by taking all the  $n$ -simplexes of  $K$  (together with their faces). Note that, if  $P$  is connected, then  $\tilde{P} \rightarrow P$  is a representative of the generator of  $H_n(P; \mathbf{Z}_2) \cong \mathbf{Z}_2$ .

In order to show the kind of arguments used, we first prove an "abstract" version of the stated result, that is

THEOREM 1. Let  $P$  be an  $n$ -cycle. Then there exists an Euler polyhedron  $P'$  such that  $P' \supset P$  and  $\dim(P' \setminus P) < n$ .

PROOF. Let  $P = |K|$  and assume that  $K = T^{(1)}$ , that is,  $K$  is the first barycentric subdivision of another triangulation  $T$  of  $P$ . Set

$$Q = \overline{\{A \in K: \chi(\text{lk}(A, K)) \equiv 1 \pmod{2}\}}.$$

$Q = |H|$  is a subpolyhedron of  $P$  and  $\dim Q < n - 1$  (as  $P$  is a cycle).

(a) Assume  $\dim Q = 0$ . Then  $Q$  consists of a finite number of points  $v_1, \dots, v_h$  and  $(P, Q)$  is an Euler pair. Let  $Z$  be the 1-skeleton of  $K$ ; then (for the properties of the barycentric subdivision)  $Z$  is a 1-cycle with boundary the 0-skeleton of  $H$ , that is,  $Q$  itself (see [A], Propositions 1 and 2, and the subsequent remark). Thus  $h$  is even and we can form  $P' = P \cup_Q \Gamma$ , where  $\Gamma$  is any 1-cycle with boundary  $Q$ .

(b) The general case. Let  $d = \dim Q$  ( $0 < d \leq n - 2$ ). We prove first of all that  $Q = |H|$  is a  $d$ -cycle. Let  $A$  be a  $(d - 1)$ -simplex of  $H$  and  $B_1, \dots, B_h$  the set of  $d$ -simplexes of  $H$  such that  $B_i > A$ . If  $C$  is a simplex of  $R = \text{lk}(A, K)$ , then  $C * A \in K$  and  $\text{lk}(C, R) = \text{lk}(C * A, K)$  (here  $*$  denotes the join operation). Since  $\dim(C * A) = \dim C + d, \chi(\text{lk}(C, R))$  is always even, except for the vertices  $v_1, \dots, v_h$  such that  $v_i * A = B_i$ . Then, by the case (a),  $h$  is even, which means that  $Q$  is a cycle.

Now we can form  $P_1 = P \cup_Q \Gamma$ , where  $\Gamma$  is any  $(d + 1)$ -cycle with boundary  $Q$ , for example the cone on  $Q$ .  $P_1$  is not necessarily an Euler space; however, if  $B$  is a  $d$ -simplex of  $H, \text{lk}(B, P_1) = \text{lk}(B, P) \amalg \{\text{odd number of points}\}$ , so that  $Q_1 = \overline{\{A \in P_1: \chi(\text{lk}(A, P_1)) \equiv 1\}}$  is a subpolyhedron of dimension  $\leq (d - 1)$  in  $P_1$ ; by iterating the argument we obtain the required Euler space  $P'$ .  $\square$

Note that the hypothesis that  $P$  is a cycle is necessary; see, for example, the following Figure 1.

The difficulty which arises in the general case is essentially to prove that  $Q$  is now a boundary in the ambient manifold.




FIGURE 1

**THEOREM 2.** *Let  $M$  be a closed  $m$ -manifold and  $P$  a cycle of dimension  $n < m$  in  $M$ . Then there exists a subpolyhedron  $P'$ ,  $P \subset P' \subset M$ , such that  $P'$  is an Euler space and  $\dim(P' \setminus P) < n$ .*

**PROOF.** Let  $Q$  be defined as in the previous theorem and  $(L, K, H)$  be a triangulation of  $(M, P, Q)$  which we assume, for the sake of simplicity, to be the first barycentric subdivision of another triangulation of  $(M, P, Q)$  (see remark below).

**CLAIM.**  $Q$  is a boundary in  $P$ .

(Note that this has already been proved in the case  $\dim Q = 0$ .) Let  $d = \dim Q$ ; let  $N$  be the simplicial neighbourhood of  $H^{(1)}$  in  $K^{(1)}$ ,  $\dot{N}$  the boundary of  $N$ ,  $p: N \rightarrow Q$  the simplicial retraction and  $\dot{p} = p|_{\dot{N}}$ .  $(\overline{P \setminus N}, \dot{N})$  is an Euler pair; therefore (again by [A, Proposition 1]), if  $Z$  denotes the  $(d + 1)$ -skeleton of  $\overline{P \setminus N}$  and  $S$  denotes the  $d$ -skeleton of  $\dot{N}$  (both with respect to  $K^{(1)}$ ), we have that  $Z$  is a  $(d + 1)$ -cycle with boundary  $S$ . Let  $f = \dot{p}|_S$ ;  $f$  is a simplicial map and we want to show that its degree is odd. Let  $\sigma \in H^{(1)}$  be a  $d$ -simplex and  $A \in H$  such that  $\sigma \subset A$ ; we must prove that  $\#\{\text{simplexes in } f^{-1}(\sigma)\} = \#\{d\text{-simplexes in } \dot{p}^{-1}(\sigma)\}$  is odd; as

$$\begin{aligned} \#\{B \in K: A < B\} &= \#\{\text{simplexes } C \text{ of } \text{lk}(A, K)\} \\ &\equiv \chi(\text{lk}(A, K)) \equiv 1 \pmod{2}, \end{aligned}$$

it is enough to show that, for each  $B > A$ ,  $\#\{d\text{-simplexes in } \dot{p}^{-1}(\sigma) \cap \dot{B}\}$  is odd. Let  $B > A$ ; then  $B = A * C$  and  $\dot{p}|_{\dot{N} \cap B}: \dot{N} \cap B \rightarrow A$  is obtained by the pseudoradial projection from  $C$ .




FIGURE 2




FIGURE 3




FIGURE 4

Note that, if  $\dim A = 0$ ,  $\#\{\text{vertices of } \hat{p}^{-1}(A) \cap \hat{B}\} = 1$  and  $\#\{\text{vertices of } \hat{p}^{-1}(A) \cap B\} = \#\{\text{vertices of } C \text{ in } K^{(1)}\} \equiv 1 \pmod{2}$  (see Figure 2); while, if  $\dim C = 0$  (so that  $B$  is a cone over  $A$  with vertex  $C$ ),  $\#\{d\text{-simplexes in } \hat{p}^{-1}(\hat{\sigma}) \cap \hat{B}\} = \#\{d\text{-simplexes in } \hat{p}^{-1}(\hat{\sigma}) \cap B\} = 1$  (see Figure 3). In general, if  $\sigma = \hat{A} * \tau$ , let  $A'$

be the face of  $A$  containing  $\tau$  and  $D = C * A'$ . Then, if  $\alpha$  is a  $d$ -simplex in  $\dot{p}^{-1}(\dot{\sigma}) \cap \dot{B}$ , necessarily  $\alpha = \dot{B} * \gamma$ , where  $\gamma$  is a  $(d - 1)$ -simplex in  $\dot{p}^{-1}(\tau) \cap D$  (see Figure 4). In order to conclude by induction, we have to show that also  $\# \{d\text{-simplexes in } \dot{p}^{-1}(\dot{\sigma}) \cap B\}$  is odd. But, if  $C'$  varies over the faces of  $C$ , and  $B' = A * C'$ , then

$$\begin{aligned} \# \{d\text{-simplexes in } \dot{p}^{-1}(\dot{\sigma}) \cap B\} &= \# \{d\text{-simplexes in } \dot{p}^{-1}(\dot{\sigma}) \cap \dot{B}\} \\ &+ \sum_{C' < C} \# \{d\text{-simplexes in } \dot{p}^{-1}(\dot{\sigma}) \cap \dot{B}'\}, \end{aligned}$$

By induction, all the terms of this sum are odd; moreover, their number equals  $\# \{C' : C' \leq C\} \equiv 1 \pmod{2}$ . Thus  $f: S \rightarrow Q$  is an odd degree map, so that the mapping cylinder  $C_f$  is a  $(d + 1)$ -cycle in  $P$  with boundary  $S \amalg Q$  and  $Q' = Z \cup_S C_f$  is the required cycle with boundary  $Q$ . This proves the claim.

In order to prove the theorem, it is enough now to put  $Q'$  transverse to  $P$  in  $M$  relatively to  $Q$  (see [RS, Theorem 5.3]). In this way we get a cycle  $Q''$  in  $M$  with boundary  $Q$  and such that  $\dim(Q'' \cap P) \leq d + 1 + n - m \leq d$ . Form  $P_1 = P \cup_Q Q''$ ;  $P_1$  is an  $n$ -cycle in  $M$  and, if  $A$  is a  $d$ -simplex in  $P_1$ , then

$$\text{lk}(A, P_1) = \begin{cases} \text{lk}(A, P) \amalg \{\text{odd number of points}\} & \text{if } A \in Q, \\ \text{lk}(A, P) & \text{if } A \in P \setminus Q'', \\ \text{lk}(A, Q'') & \text{if } A \in Q'' \setminus P, \\ \text{lk}(A, P) \amalg \{\text{even number of points}\} & \text{if } A \in Q'' \cap P. \end{cases}$$

In each case  $\chi(\text{lk}(A, P_1)) \equiv 0$ , so that  $Q_1 = \overline{\{A \in P_1 : \chi(\text{lk}(A, P_1)) \equiv 1\}}$  has dimension  $\leq (d - 1)$  and we can iterate the argument as before until we get an Euler space  $P'$ .  $\square$

REMARK. As regards the choice of the triangulation, what we need is only that the simplicial neighbourhood  $N$  of  $Q$  in  $P$  (with respect to  $K^{(1)}$ ) is in fact a regular neighbourhood; therefore, any triangulation  $(K, H)$  such that  $Q$  is full in  $P$  would be enough (see [RS] for a definition of full).

COROLLARY. Every homology class  $z \in H_n(M, \mathbf{Z}_2)$  can be represented as the fundamental class of an Euler subpolyhedron of dimension  $n$  in  $M$ .

ADDENDUM. With respect to the problem stated in the introduction (that is, to represent  $\mathbf{Z}_2$ -homology classes of a real algebraic manifold by algebraic subvarieties), since this paper was written we have proved the following (see [BD]):

For each  $d \geq 11$ , there exists a compact smooth manifold  $V$  and a class  $z \in H_{d-2}(V, \mathbf{Z}_2)$  such that, for any homeomorphism  $h: V \rightarrow V'$  between  $V$  and a real algebraic manifold  $V'$ ,  $h_*(z) \in H_{d-2}(V', \mathbf{Z}_2)$  cannot be represented by an algebraic subvariety of  $V'$ .

## REFERENCES

- [A] E. Akin, *Stiefel-Whitney homology classes and bordism*, Trans. Amer. Math. Soc. **205** (1975), 341–359.
- [BT] R. Benedetti and A. Tognoli, *Remarks and counterexamples in the theory of real algebraic vector bundles and cycles*, Lecture Notes, Springer, Berlin and New York (to appear).
- [RS] C. P. Rourke and B. J. Sanderson, *Introduction to piecewise-linear topology*, Ergebnisse der Math. und ihrer Grenzgebiete, Band 69, Springer-Verlag, Berlin and New York, 1972.
- [S] D. Sullivan, *Combinatorial invariants of analytic spaces*, Proc. Liverpool Singularities Symposium. I, Lecture Notes in Math., vol. 192, Springer-Verlag, Berlin and New York, 1971.
- [BD] R. Benedetti and M. Dedò, *Counterexamples to representing homology classes by real algebraic subvarieties up to homeomorphism* (to appear).

ISTITUTO MATEMATICO, VIA F. BUONARROTI 2, 56100 PISA, ITALY