

BETTER BOUNDS FOR PERIODIC SOLUTIONS OF DIFFERENTIAL EQUATIONS IN BANACH SPACES

STAVROS N. BUSENBERG, DAVID C. FISHER AND MARIO MARTELLI

ABSTRACT. Let f be Lipschitz with constant L in a Banach space and let $x(t)$ be a P -periodic solution of $x'(t) = f(x(t))$. We show that $P \geq 6/L$. An example is given with $P = 2\pi/L$, so the bound is nearly strict. We also give a short proof that $P \geq 2\pi/L$ in a Hilbert space.

I. Introduction. Suppose f is Lipschitz with constant L in a Banach space and $x(t)$ is a P -periodic solution of $x'(t) = f(x(t))$. How small can LP be?

This question was first addressed when Lasota and Yorke [3] proved $LP \geq 4$. Busenberg and Martelli [2] showed $LP \geq 4\frac{1}{2}$. We show that $LP \geq 6$.

The lowest known LP is 2π , e.g. for nonzero solutions of $u'(t) = v(t)$ and $v'(t) = -u(t)$. Also, in a Hilbert space, $LP \geq 2\pi$ (Lasota and Yorke [3], Yorke [4])—A short proof is given in §III). So the Hilbert space bound is strict, but the Banach space bound may not be strict.

II. In Banach spaces, $LP \geq 6$.

LEMMA 1. Let B be a Banach space and let $y: \mathbf{R} \rightarrow B$ be continuous and P -periodic with $\|y'(t)\|$ integrable. Then

$$\int_0^P \int_0^P \|y(t) - y(s)\| ds dt \leq \frac{P}{6} \int_0^P \int_0^P \|y'(t) - y'(s)\| ds dt.$$

PROOF.

$$\begin{aligned} A &\equiv \int_0^P \int_0^P \|y(t) - y(s)\| ds dt = \int_0^P \int_0^P \|y(s+t) - y(s)\| ds dt \\ &= \int_0^P \int_0^P \frac{(P-t)t}{P} \left\| \frac{y(s+t) - y(s)}{t} - \frac{y(s) - y(s+t-P)}{P-t} \right\| ds dt \\ &= \int_0^P \int_0^P \frac{(P-t)t}{P^2} \left\| \int_0^P \left(y' \left(s + \frac{tr}{P} \right) - y' \left(s + \frac{tr}{P} - r \right) \right) dr \right\| ds dt \\ &\leq \int_0^P \int_0^P \frac{(P-t)t}{P^2} \int_0^P \left\| y' \left(s + \frac{tr}{P} \right) - y' \left(s + \frac{tr}{P} - r \right) \right\| dr ds dt \\ &= \int_0^P \frac{(P-t)t}{P^2} \int_0^P \int_0^P \left\| y' \left(s + \frac{tr}{P} \right) - y' \left(s + \frac{tr}{P} - r \right) \right\| ds dr dt. \end{aligned}$$

Received by the editors March 21, 1986.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 34C45.

Key words and phrases. Periodic solutions, Lipschitz constant, Wirtinger's inequality, Banach space, Hilbert space, autonomous differential equations.

The first author was partially supported by NSF Grant No. DMS-8507193

Since the inner integral is over one period, it can be shifted by $tr/P - r$ to yield

$$\begin{aligned} A &\leq \int_0^P \frac{(P-t)t}{P^2} dt \int_0^P \int_0^P \|y'(s+r) - y'(s)\| ds dr \\ &= \frac{P}{6} \int_0^P \int_0^P \|y'(r) - y'(s)\| ds dr. \end{aligned}$$

THEOREM 1. *If f is Lipschitz with constant L in a Banach space and $x(t)$ is a nonconstant P -periodic solution of $x'(t) = f(x(t))$, then $LP \geq 6$.*

PROOF.

$$\begin{aligned} \int_0^P \int_0^P \|x(t) - x(s)\| ds dt &\leq \frac{P}{6} \int_0^P \int_0^P \|x'(t) - x'(s)\| ds dt \\ &= \frac{P}{6} \int_0^P \int_0^P \|f(x(t)) - f(x(s))\| ds dt \leq \frac{LP}{6} \int_0^P \int_0^P \|x(t) - x(s)\| ds dt. \end{aligned}$$

Solving for LP gives the result.

III. A short proof that $LP \geq 2\pi$ in Hilbert spaces.

LEMMA 2 (*a Hilbert space analog of Wirtinger's inequality*). *Let H be a Hilbert space and let $y: \mathbb{R} \rightarrow H$ be continuous and P -periodic with $\int_0^P y(t) dt = 0$ and $\|y'(t)\|^2$ integrable. Then*

$$\int_0^P \|y(t)\|^2 dt \leq \frac{P^2}{4\pi^2} \int_0^P \|y'(t)\|^2 dt.$$

PROOF. Since the path of $y(t)$ is compact, there is a countable orthonormal set e_1, e_2, \dots with $y(t) = \sum_{i=1}^\infty a_i(t)e_i$. Each $a_i(t)$ is P -periodic and $\int_0^P a_i(t) dt = 0$, so Wirtinger's inequality [1] gives $\int_0^P a_i(t)^2 dt \leq (P^2/4\pi^2) \int_0^P a_i'(t)^2 dt$. Then

$$\begin{aligned} \int_0^P \|y(t)\|^2 dt &= \int_0^P \left\| \sum_{i=1}^\infty a_i(t)e_i \right\|^2 dt = \int_0^P \sum_{i=1}^\infty a_i(t)^2 dt \\ &= \sum_{i=1}^\infty \int_0^P a_i(t)^2 dt \leq \frac{P^2}{4\pi^2} \sum_{i=1}^\infty \int_0^P a_i'(t)^2 dt = \frac{P^2}{4\pi^2} \int_0^P \sum_{i=1}^\infty a_i'(t)^2 dt \\ &= \frac{P^2}{4\pi^2} \int_0^P \left\| \sum_{i=1}^\infty a_i'(t)e_i \right\|^2 dt = \frac{P^2}{4\pi^2} \int_0^P \|y'(t)\|^2 dt. \end{aligned}$$

THEOREM 2 [3, 4]. *If f is Lipschitz with constant L in a Hilbert space and $x(t)$ is a nonconstant P -periodic solution of $x'(t) = f(x(t))$, then $LP \geq 2\pi$.*

PROOF. Pick h with $x(h) \neq x(0)$. Since $x(t)$ is P -periodic, $x(t+h) - x(t)$ is P -periodic and $\int_0^P (x(t+h) - x(t)) dt = 0$. Then from Lemma 2

$$\begin{aligned} \int_0^P \|x(t+h) - x(t)\|^2 dt &\leq \frac{P^2}{4\pi^2} \int_0^P \|x'(t+h) - x'(t)\|^2 dt \\ &= \frac{P^2}{4\pi^2} \int_0^P \|f(x(t+h)) - f(x(t))\|^2 dt \leq \frac{L^2 P^2}{4\pi^2} \int_0^P \|x(t+h) - x(t)\|^2 dt. \end{aligned}$$

Solving for LP gives the result.

NOTE ADDED IN PROOF. The bound in Theorem 1 is now known to be sharp. We have constructed an example in a Banach space with $LP = 6$.

REFERENCES

1. W. Blaschke, *Kries und Kugel*, Chelsea, New York, 1916.
2. S. Busenberg and M. Martelli, *Bounds for the period of periodic orbits of dynamical systems*, J. Differential Equations (to appear).
3. A. Lasota and J. Yorke, *Bounds for periodic solutions of differential equations in Banach spaces*, J. Differential Equations **10** (1971), 83–91.
4. J. Yorke, *Periods for periodic solutions and the Lipschitz constant*, Proc. Amer. Math. Soc. **22** (1969), 509–512.

DEPARTMENT OF MATHEMATICS, HARVEY MUDD COLLEGE, CLAREMONT, CALIFORNIA 91711 (Current address of S. N. Busenberg and D. C. Fisher)

DEPARTMENT OF MATHEMATICS, BRYN MAWR COLLEGE, BRYN MAWR, PENNSYLVANIA 19010 (Current address of Mario Martelli)