

## A NOTE ON QUADRICS THROUGH AN ALGEBRAIC CURVE

FERNANDO SERRANO

(Communicated by William C. Waterhouse)

**ABSTRACT.** In this note we describe the intersection of all quadric hypersurfaces containing a given linearly normal smooth projective curve of genus  $n$  and degree  $2n + 1$ .

Let  $C$  be an irreducible nonsingular curve of genus  $g$ , defined over an algebraically closed field of any characteristic. Let  $C$  be embedded in  $\mathbf{P}^r$  by a complete linear system  $|L|$ . Saint-Donat [5] has proved that if  $\deg L \geq 2g + 2$  then the homogeneous ideal  $I_C$  of  $C \subseteq \mathbf{P}^r$  is generated by quadrics, and if  $\deg L = 2g + 1$  then  $I_C$  is generated by quadrics and cubics (see also Fujita [1]). In [2], Green and Lazarsfeld have announced the following result: In case  $\deg L = 2g + 1$ ,  $I_C$  fails to be generated by quadrics if and only if  $C$  is hyperelliptic or  $L$  embeds  $C$  with a trisecant line, i.e.,  $H^0 \mathcal{O}_C(L - K_C) \neq 0$ , where  $K_C$  denotes the canonical divisor on  $C$ . In this note we describe the intersection of all quadric hypersurfaces passing through  $C \subseteq \mathbf{P}^r$  in the borderline situation  $\deg L = 2g + 1$ . The main ingredient of the proof is a theorem of Castelnuovo on the postulation of points.

A  $g_d^1$  on a curve is, by definition, a base-point free linear system of degree  $d$  and dimension 1. For the definition and properties of rational normal scrolls see [3].

Our result is the following.

**THEOREM.** *Let  $C \subseteq \mathbf{P}^{n+1}$  be a linearly normal smooth irreducible curve of genus  $n \geq 4$  and degree  $2n + 1$ . If  $W(C)$  denotes the intersection of all quadric hypersurfaces of  $\mathbf{P}^{n+1}$  which contain  $C$ , then either  $W(C)$  consists of  $C$  plus (possibly) a line and finitely many isolated points, or  $W(C)$  is a rational normal scroll of dimension 2. In case  $W(C)$  is a scroll, one of the following situations occurs:*

(i)  *$W(C)$  is smooth and  $C$  meets every fiber of  $W(C)$  at three points.  $C$  is trigonal and embedded by the linear system  $|K_C + g_3^1|$ .*

(ii)  *$W(C)$  is a cone with vertex  $P$ , and  $C$  passes through  $P$  and meets every fiber of  $W(C)$  at  $P$  plus two other points.  $C$  is hyperelliptic and embedded by  $|P + ng_2^1|$ .*

(iii)  *$W(C)$  is smooth and  $C$  is a divisor in  $W(C)$  of class  $2H + R$ , where  $H$  denotes a hyperplane and  $R$  a fiber of the ruling. In particular  $C$  is hyperelliptic, the  $g_2^1$  being given by restriction of the ruling of  $W(C)$ .*

**PROOF.** Throughout this proof we will assume that  $W(C)$  is not the union of  $C$  and (possibly) a line plus finitely many points. Consequently, there exists a curve  $G \subseteq W(C)$ ,  $G \neq C$ , with degree of  $G \geq 2$ .  $G$  is allowed to be a pair of distinct lines. Pick two distinct general points  $Q_1$  and  $Q_2$  on  $G$ , none of them on  $C$ . If  $G$  is a

Received by the editors October 20, 1986.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 14E25, 14H45, 14N05.

*Key words and phrases.* Rational scroll, hyperelliptic, trigonal.

union of two lines, then by a general pair we mean that  $Q_1$  is a general point on one of the lines and  $Q_2$  is general on the other line. Choose now a general hyperplane  $\mathbf{P}^n$  in  $\mathbf{P}^{n+1}$  passing through  $Q_1$  and  $Q_2$ , and set  $\Gamma = C \cap \mathbf{P}^n$ . Let  $W(\Gamma)$  be the intersection of all quadric hypersurfaces of  $\mathbf{P}^n$  which contain  $\Gamma$ . If  $I_C, I_\Gamma$  denote the ideal sheaves of  $C$  in  $\mathbf{P}^{n+1}$  and  $\Gamma$  in  $\mathbf{P}^n$  respectively, then the exact sequence

$$0 = H^0(\mathbf{P}^{n+1}, I_C(1)) \rightarrow H^0(\mathbf{P}^{n+1}, I_C(2)) \\ \rightarrow H^0(\mathbf{P}^n, I_\Gamma(2)) \rightarrow H^1(\mathbf{P}^{n+1}, I_C(1)) = 0$$

yields  $W(\Gamma) = W(C) \cap \mathbf{P}^n$ .

CLAIM 1.  $\Gamma$  consists of  $2n + 1$  points in general linear position (i.e., any subset of  $n + 1$  points of  $\Gamma$  spans  $\mathbf{P}^n$ ).

PROOF OF CLAIM 1. Let  $(\mathbf{P}^{n+1})^*$  be the space of hyperplanes of  $\mathbf{P}^{n+1}$ . It is a well-known fact that the set

$$\mathcal{U} = \{H \in (\mathbf{P}^{n+1})^* \mid H \cap C \text{ is in general linear position}\}$$

is dense in  $(\mathbf{P}^{n+1})^*$ . For  $i = 1, 2$ , the set  $M(Q_i) = \{H \in (\mathbf{P}^{n+1})^* \mid Q_i \in H\}$  is a hyperplane of  $(\mathbf{P}^{n+1})^*$ . Since degree of  $G \geq 2$  we have

$$\bigcup_{Q_1, Q_2 \in G} (M(Q_1) \cap M(Q_2)) = (\mathbf{P}^{n+1})^*$$

and thus  $M(Q_1) \cap M(Q_2) \cap \mathcal{U} \neq \emptyset$  for a generic choice of  $Q_1$  and  $Q_2$ . This proves Claim 1.

Choose linear subvarieties  $\hat{\mathbf{P}}^{n-1}, \hat{\mathbf{P}}^n$  of  $\mathbf{P}^{n+1}$  of dimensions  $n - 1$  and  $n$  respectively. Let  $\pi: C \rightarrow \hat{\mathbf{P}}^{n-1}$  be the projection of  $C$  from the line  $\overline{Q_1 Q_2}$  spanned by  $Q_1$  and  $Q_2$ , and let  $\pi_1: C \rightarrow \hat{\mathbf{P}}^n$  be the projection of  $C$  from  $Q_1$ .

CLAIM 2.  $\pi$  and  $\pi_1$  are generically one-to-one.

PROOF OF CLAIM 2. It suffices to prove the statement for  $\pi$ . Since  $n + 1 \geq 5$ , any hyperplane passing through  $Q_1$  and  $Q_2$  contains at least three fibers of  $\pi$ . If  $\pi$  has degree  $k \geq 2$  then those three fibers consist of  $3k \geq 6$  points which span a  $\mathbf{P}^3$  or a  $\mathbf{P}^4$ , so that they are not in general linear position. But this contradicts Claim 1.

CLAIM 3. A general hyperplane of  $\hat{\mathbf{P}}^n$  passing through the point  $\overline{Q_1 Q_2} \cap \hat{\mathbf{P}}^n$  cuts  $\pi_1(C)$  at a set of points in general linear position.

PROOF OF CLAIM 3. We argue as in Claim 1. The set

$$\mathcal{U}' = \{H \in (\hat{\mathbf{P}}^n)^* \mid H \cap \pi_1(C) \text{ is in general linear position}\}$$

is dense in  $(\hat{\mathbf{P}}^n)^*$ , and  $N(Q_2) = \{H \in (\hat{\mathbf{P}}^n)^* \mid \overline{Q_1 Q_2} \cap \hat{\mathbf{P}}^n \in H\}$  is a hyperplane of  $(\hat{\mathbf{P}}^n)^*$ . Fix  $Q_1$ . Since  $\deg G \geq 2$ , the points  $\overline{Q_1 Q_2} \cap \hat{\mathbf{P}}^n$  describe a curve in  $\hat{\mathbf{P}}^n$  as  $Q_2$  varies along  $G$ . Therefore

$$(\hat{\mathbf{P}}^n)^* = \bigcup_{Q_2 \in G} N(Q_2),$$

and thus  $N(Q_2) \cap \mathcal{U}' = \emptyset$  for at most finitely many  $Q_2$ 's.

CLAIM 4.  $\Gamma \cup \{Q_1, Q_2\}$  is in general linear position in  $\mathbf{P}^n$ .

PROOF OF CLAIM 4. Choose any subset  $\Omega$  of  $n + 1$  points in  $\Gamma \cup \{Q_1, Q_2\}$ . We have to show that  $\Omega$  spans  $\mathbf{P}^n$ .

Case 1.  $\Omega \subseteq \Gamma$ . The claim is obvious because  $\Gamma$  is in general linear position.

*Case 2.*  $\Omega = \{Q_1, T_1, \dots, T_n\}$  with  $\{T_1, \dots, T_n\} \subseteq \Gamma$ . By Claim 3, a general hyperplane  $\mathbf{P}^n \subseteq \mathbf{P}^{n+1}$  containing  $\overline{Q_1 Q_2}$  cuts  $\hat{\mathbf{P}}^n$  along an  $(n-1)$ -plane  $\tilde{\mathbf{P}}^{n-1}$  such that  $\tilde{\mathbf{P}}^{n-1} \cap \pi_1(C)$  is in general linear position. By Claim 2,  $\pi_1(T_1), \dots, \pi_1(T_n)$  are all distinct and belong to  $\tilde{\mathbf{P}}^{n-1} \cap \pi_1(C)$ . Since  $\{\pi_1(T_1), \dots, \pi_1(T_n)\}$  spans  $\tilde{\mathbf{P}}^{n-1}$ , it follows that  $\{Q_1, \pi_1(T_1), \dots, \pi_1(T_n)\}$  spans  $\mathbf{P}^n$ , and so does  $\Omega$ .

*Case 3.*  $\Omega = \{Q_1, Q_2, T_1, \dots, T_{n-1}\}$  with  $\{T_1, \dots, T_{n-1}\} \subseteq \Gamma$ . If  $\tilde{\mathbf{P}}^{n-2} = \hat{\mathbf{P}}^{n-1} \cap \mathbf{P}^n$  then  $\tilde{\mathbf{P}}^{n-2} \cap \pi(C)$  is in general linear position. The points  $\pi(T_1), \dots, \pi(T_{n-1})$  are all distinct because of Claim 2, and they belong to  $\tilde{\mathbf{P}}^{n-2} \cap \pi(C)$ . Inasmuch as  $\{\pi(T_1), \dots, \pi(T_{n-1})\}$  spans  $\tilde{\mathbf{P}}^{n-2}$  we get that  $\{Q_1, Q_2, \pi(T_1), \dots, \pi(T_{n-1})\}$  spans  $\mathbf{P}^n$ , and so does  $\Omega$ .

Let us summarize the results obtained so far. For a general hyperplane section  $\Gamma = C \cap \mathbf{P}^n$  of  $C$  we can find two points  $Q_1, Q_2 \in W(\Gamma)$  such that  $\Gamma \cup \{Q_1, Q_2\}$  is in general linear position. Since  $\Gamma$  imposes exactly  $2n + 1$  conditions on quadrics [3, p. 36], so does  $\Gamma \cup \{Q_1, Q_2\}$ . Hence  $\Gamma \cup \{Q_1, Q_2\}$  is a set of  $2n + 3$  points in general linear position in  $\mathbf{P}^n$  which imposes  $2n + 1$  conditions on quadrics. Here we use the main ingredient of the proof: a lemma of Castelnuovo states that  $\Gamma \cup \{Q_1, Q_2\}$  must lie on a rational normal curve  $B \subseteq \mathbf{P}^n$  [3, p. 36].

Pick a quadric  $R$  in  $\mathbf{P}^n$  which contains  $\Gamma$ . If  $B$  is not contained in  $R$  then  $2n + 1 = \text{cardinal of } \Gamma \leq \text{cardinal of } (R \cap B) = 2n$ , absurd. Hence  $B \subseteq R$ . Since the ideal of  $B$  is generated by quadrics we get  $W(\Gamma) = B$ . Now recall that  $W(\Gamma) = W(C) \cap \mathbf{P}^n$ . Notice that the above considerations hold for a general hyperplane  $\mathbf{P}^n$  of  $\mathbf{P}^{n+1}$ . It follows that  $W(C)$  is a surface of minimal degree.  $W(C)$  cannot be the Veronese surface in  $\mathbf{P}^5$  because  $C$  has odd degree and is contained in  $W(C)$ . Therefore  $W(C)$  is a rational normal scroll of dimension 2 [3, p. 51]. The homogeneous ideal of  $C$  in  $\mathbf{P}^{n+1}$  is generated by quadrics and cubics [5] and thus  $C$  meets every fiber of  $W(C)$  at no more than three points. Next we are going to classify the possible configurations  $(W(C), C)$ .

Assume first that  $W(C)$  is a cone. The vertex  $P$  of  $W(C)$  must belong to  $C$  (otherwise  $C$  would have degree  $2n$  or  $3n$ ), and  $C$  meets every fiber of  $W(C)$  at two other points. Now it is obvious that  $C$  is hyperelliptic, and that any hyperplane section of  $C$  passing through  $P$  belongs to the system  $|P + ng_2^1|$ .

Suppose that  $W(C)$  is nonsingular, and denote by  $F$  a general fiber of  $W(C)$ . If  $C$  meets  $F$  at three points then  $C$  is trigonal, and an easy application of the Riemann-Roch formula shows that the divisors of the  $g_3^1$  span lines only when the hyperplane divisor belongs to the system  $|K_C + g_3^1|$ . In case  $C$  meets  $F$  at two points and  $H$  denotes a hyperplane divisor of  $W(C)$  we have  $H^2 = n$ ,  $CH = 2n + 1$  and  $C$  is linearly equivalent to  $2H + bF$ . One concludes that  $b = 1$ .

REMARK. By Green-Lazarsfeld's claim, quoted in the Introduction, it follows that in case  $W(C)$  is not a scroll and  $W(C)$  contains a line, then this line is a trisecant of  $C$ .

REFERENCES

1. T. Fujita, *Defining equations for certain types of polarized varieties*, Complex Analysis and Algebraic Geometry, Cambridge Univ. Press, 1977.
2. M. Green and R. Lazarsfeld, *On the projective normality of complete linear series on an algebraic curve*, Invent. Math. **83** (1985), 73-90.

3. J. Harris, *A bound on the geometric genus of projective varieties*, Ann. Scuola Norm. Sup. Pisa Ser. (4) **8** (1981), 35–68.
4. P. Maroscia and W. Vogel, *On the defining equations of points in general position in  $\mathbf{P}^n$* , Math. Ann. **269** (1984), 183–189.
5. B. Saint-Donat, *Sur les équations définissant une courbe algébrique*, C. R. Acad. Sci. Paris Ser. A **274** (1972), 324–327 and 487–489.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CALIFORNIA 94720

*Permanent address:* Departament d'Algebra i Geometria, Facultat de Matemàtiques, Universitat de Barcelona, Gran Via 585, 08007 Barcelona, Spain