

CORRIGENDUM TO
"WEAK SEQUENTIAL DENSENESS IN BANACH SPACES"

J. HOWARD

(Communicated by John B. Conway)

In [2] the proof of the theorem for $(a) \Rightarrow (b)$ is incorrect. For separable Banach spaces X , the implication $(a) \Rightarrow (b)$ is given in [3]. The nonseparable case may be reduced to the separable case based on the fact that if S is a subset of a Banach space X and $x \in \overline{S}^w$, then S has a countable subset S_0 with $x \in \overline{S_0}^w$ (Part a, Theorem 8.12.4, p. 549 of [1]). Hence the Theorem of this paper remains correct.

I wish to thank Alfred D. Andrew for this correction.

REFERENCES

1. R. E. Edwards, *Functional analysis*, Holt, Rinehart, and Winston, New York, 1965.
2. J. Howard, *Weak sequential denseness in Banach spaces*, Proc. Amer. Math. Soc. **99** (1987), 351–352.
3. H. P. Rosenthal, *Some recent discoveries in the isomorphic theory of Banach spaces*, Bull. Amer. Math. Soc. **84** (1978), 803–831.

DEPARTMENT OF MATHEMATICS, NEW MEXICO HIGHLANDS UNIVERSITY, LAS VEGAS,
NEW MEXICO 87701

Received by the editors June 18, 1987.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 46B20; Secondary 46B22.

Key words and phrases. Weakly sequentially dense, weakly angelic.

©1988 American Mathematical Society
0002-9939/88 \$1.00 + \$.25 per page