

THE DENSITY OF ALTERNATION POINTS IN RATIONAL APPROXIMATION

P. B. BORWEIN, * A. KROÓ, * R. GROTHMANN * AND E. B. SAFF **

(Communicated by Irwin Kra)

ABSTRACT. We investigate the behavior of the equioscillation (alternation) points for the error in best uniform rational approximation on $[-1, 1]$. In the context of the Walsh table (in which the best rational approximant with numerator degree $\leq m$, denominator degree $\leq n$, is displayed in the n th row and the m th column), we show that these points are dense in $[-1, 1]$, if one goes down the table along a ray above the main diagonal ($n = [cm], c < 1$). A counterexample is provided showing that this may not be true for a sub-diagonal of the table. In addition, a Kadec-type result on the distribution of the equioscillation points is obtained for asymptotically horizontal paths in the Walsh table.

1. STATEMENT OF RESULTS

Denote by $\mathcal{R}_{m,n}$ the rational functions with numerator in Π_m , the set of algebraic real polynomials of degree at most m , and denominator in Π_n . Then the best approximation $r_{m,n}^* = p_{m,n}^*/q_{m,n}^*$ in $\mathcal{R}_{m,n}$ to $f \in C[-1, 1]$ with respect to the uniform norm

$$(1.1) \quad \|g\|_{[-1,1]} := \sup\{|g(x)| : x \in [-1, 1]\}$$

is unique and is characterized by an equioscillation property [M], i.e., there are $m + n + 2 - d$ points

$$(1.2) \quad -1 \leq x_1^{(m,n)} < \cdots < x_{m+n+2-d}^{(m,n)} \leq 1,$$

where

$$(1.3) \quad d := d(m, n) := \min\{m - \deg p_{m,n}^*, n - \deg q_{m,n}^*\},$$

such that for a $\sigma = \pm 1$ and all $k = 1, \dots, m + n + 2 - d$

$$(1.4) \quad f(x_k^{(m,n)}) - r_{m,n}^*(x_k^{(m,n)}) = \sigma(-1)^k \|f - r_{m,n}^*\|_{[-1,1]}.$$

Received by the editors June 9, 1987 and, in revised form, March 28, 1988.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 41A20.

Key words and phrases. Rational approximation, extreme points, best approximants.

* The research of these authors was done while visiting the Institute for Constructive Mathematics, University of South Florida, Tampa.

** The research of this author was supported, in part, by the National Science Foundation under grant DMS-8620098.

(Here and below, we assume that $p_{m,n}^*$ and $q_{m,n}^*$ do not have a common factor.)

Much is known about the behavior of alternation points for best polynomial approximation ($n = 0$). For this case, Lorentz [L] and Kroó and Saff [KS] give examples showing that for a subsequence $\{m_k\}$ the alternation points (1.2), with $m = m_k$, $n = 0$, may avoid a subinterval of $[-1, 1]$. However, Kadec [K] proved that there is always a subsequence such that the alternation points behave like the extremal points of the Chebyshev polynomial of degree $m + 1$, that is, like $\{\cos[k\pi/(m+1)]\}_{k=0}^{m+1}$. For polynomial approximation, this implies the denseness of the alternation points in $[-1, 1]$.

For rational approximation, given m and n , we pick any alternation set (1.2) and write

$$(1.5) \quad \rho_{m,n}(f) := \sup_{x \in [-1, 1]} \min_k |x - x_k^{(m,n)}|$$

as a measure for the density of the alternation set in $[-1, 1]$. We shall prove

Theorem 1.1. *Let $n = n(m)$ satisfy*

$$(1.6) \quad n(m) \leq n(m+1) \leq n(m) + 1, \quad n(m) \leq m,$$

for $m = 0, 1, \dots$. If $f \in C[-1, 1]$, $f \notin \mathcal{R}_{m, n(m)}$, $m = 0, 1, \dots$, then

$$(1.7) \quad \liminf_{m \rightarrow \infty} \left(\frac{m - n(m)}{\log m} \right) \rho_{m, n(m)}(f) < \infty.$$

The proof of Theorem 1.1 will be given in §2.

Remark. Theorem 1.1 applies in the case $n(m) = [cm]$ for any constant $c \leq 1$, where $[\cdot]$ denotes the greatest integer function. If $c < 1$, we deduce from (1.7) that

$$(1.8) \quad \liminf_{m \rightarrow \infty} \rho_{m, [cm]}(f) = 0,$$

which implies that the alternation points are dense in $[-1, 1]$ for such a “ray sequence” of best approximants. On the other hand, we show in Theorem 1.3 below that this density may not hold when $m/n(m) \rightarrow 1$.

Our second result is similar to Kadec’s result [K] on polynomial approximation. We write for $-1 \leq \alpha < \beta \leq 1$ (with $x_k^{(m,n)}$ as in (1.2))

$$(1.9) \quad N_{m,n}(\alpha, \beta) := \#\{x_k^{(m,n)} : \alpha \leq x_k^{(m,n)} \leq \beta, k = 1, \dots, m+n+2-d\}.$$

Theorem 1.2. *Assume, in addition to the hypotheses of Theorem 1.1, that*

$$(1.10) \quad \lim_{m \rightarrow \infty} \frac{n(m)}{m} = 0.$$

Then there exists a subsequence Ω of \mathbb{N} such that for all $[\alpha, \beta] \subseteq [-1, 1]$,

$$(1.11) \quad \lim_{\substack{m \rightarrow \infty \\ m \in \Omega}} \frac{N_{m, n(m)}(\alpha, \beta)}{N_{m, n(m)}(-1, 1)} = \frac{\arccos \alpha - \arccos \beta}{\pi}.$$

Finally, we give a counterexample, which shows that Theorems 1.1 and 1.2 cannot be proved for a subdiagonal of the Walsh table. Indeed, for approximation in $\mathcal{R}_{n-1, n}$ it is possible that, for all n , the extremal points all reside in an arbitrarily small interval.

Theorem 1.3. *For every $2 > \varepsilon > 0$, there is a function $f \in C[-1, 1]$ such that for each $n = 1, 2, \dots$ the error $f - r_{n-1,n}^*(f)$ has no alternation points in $(-1 + \varepsilon, 1]$.*

The proof of Theorem 1.3 will be given in §3. The results of this paper should be compared with those of Kroó and Peherstorfer [KP] for L_1 -approximation.

2. PROOFS OF THEOREMS 1.1 AND 1.2

We need the following lemma, which follows easily from classical results. We include the proof for the sake of completeness.

Lemma 2.1. *Given $-1 \leq \alpha < \beta \leq 1$ and $n \in \mathbb{N}$ there exists a $p_n \in \Pi_n$ with*

$$(2.1) \quad \|p_n\|_{[-1, \alpha] \cup [\beta, 1]} < 1,$$

and

$$(2.2) \quad \|p_n\|_{[\alpha, \beta]} > c_1 e^{c_2 n(\beta - \alpha)},$$

where $c_1, c_2 > 0$ are constants independent of α, β and n .

In (2.1) and (2.2) the norms are again the sup norms over the indicated set.

Proof. Let

$$(2.3) \quad T_m(x) := \cos(m \arccos x)$$

denote the Chebyshev polynomial of degree m . For $m := [n/2]$, $\tau := (\beta - \alpha)/2$, set

$$(2.4) \quad q_n(x) := \frac{1}{2} T_m \left(1 + \frac{\tau^2}{2} - \left(2 + \frac{\tau^2}{2} \right) \frac{x^2}{4} \right).$$

Since $\tau \leq 1$ and

$$(2.5) \quad T_m(1 + \eta) \geq \frac{1}{2} (1 + \sqrt{2\eta})^m, \quad \eta > 0,$$

we have for some constants $c_1, c_2 > 0$:

$$(2.6) \quad q_n(0) = \frac{1}{2} T_m \left(1 + \frac{\tau^2}{2} \right) \geq \frac{1}{4} (1 + \tau)^m > c_1 e^{nc_2(\beta - \alpha)}.$$

For $x \in [-2, -\tau] \cup [\tau, 2]$,

$$(2.7) \quad -1 \leq 1 + \frac{\tau^2}{2} - \left(2 + \frac{\tau^2}{2} \right) \frac{x^4}{4} < 1.$$

The lemma now follows with $p_n(x) := q_n(x - (\alpha + \beta)/2)$. \square

Proof of Theorem 1.1. Set $E_m(f) := \|f - r_{m,n(m)}^*\|_{[-1, 1]}$ for $m \in \mathbb{N}$. Since $f \notin \mathcal{R}_{m,n(m)}$, we have $E_m(f) > 0$ for all $m \in \mathbb{N}$. Also, from (1.6), it follows that $E_m(f) \downarrow 0$, and so from elementary theorems about series (cf. [K])

$$(2.8) \quad \sum_{m=0}^{\infty} \frac{E_m(f) - E_{m+1}(f)}{E_m(f) + E_{m+1}(f)} = \infty.$$

Thus there is a subsequence Ω of \mathbb{N} with $E_m(f) - E_{m+1}(f) \neq 0$ and

$$(2.9) \quad \frac{E_m(f) + E_{m+1}(f)}{E_m(f) - E_{m+1}(f)} < m^2$$

for all $m \in \Omega$.

For $m \in \Omega$, set

$$(2.10) \quad R_m := \frac{1}{E_m(f) - E_{m+1}(f)} (r_{m,n(m)}^* - r_{m+1,n(m+1)}^*).$$

At the alternation points

$$(2.11) \quad -1 \leq x_1^{(m)} < \cdots < x_{m+n(m)+2-d(m)}^{(m)} \leq 1$$

of $f - r_{m,n(m)}^*$ we have with $\sigma = \pm 1$

$$(2.12) \quad \sigma(-1)^k R_m(x_k^{(m)}) \geq 1, \quad k = 1, \dots, m+n(m)+2-d(m).$$

Moreover, from (1.6) and (2.12) it follows that $R_m = P_m/Q_m$ with

$$(2.13) \quad \deg P_m = m+n(m)+1-d(m),$$

$$(2.14) \quad \deg Q_m \leq 2n(m)+1-d(m).$$

Thus $R_m - q$ can have at most $m+n(m)+1-d(m)$ zeros, if $q \in \Pi_{m-n(m)}$.

Let c_1, c_2 be as in Lemma 2.1. For $m \in \Omega$, let $x_m^* \in [-1, 1]$ satisfy

$$(2.15) \quad \min_k |x_m^* - x_k^{(m)}| = \rho_{m,n(m)}(f) =: t_m.$$

If $x_m^* \in [-1, x_1^{(m)}]$, we let $p_{m-n(m)}$ be the polynomial that satisfies Lemma 2.1 with $\alpha = -1$ and $\beta = x_1^{(m)}$. From (2.12) and (2.1) it follows that $R_m \pm p_{m-n(m)}$ has $m+n(m)+1-d(m)$ zeros in $(x_1^{(m)}, 1]$ and hence is zero-free in $[-1, x_1^{(m)}]$. Thus

$$(2.16) \quad c_1 e^{c_2 t_m(m-n(m))} \leq \|R_m\|_{[-1,1]} < m^2,$$

where the last inequality follows from (2.9). If $x_m^* \in [x_{m+n(m)+2-d(m)}^{(m)}, 1]$, we use Lemma 2.1 with $\alpha = x_{m+n(m)+2-d(m)}^{(m)}$ and $\beta = 1$ and again we get (2.16). Otherwise denote the zeros of R_m by $y_k^{(m)}$, where

$$(2.17) \quad x_1^{(m)} < y_1^{(m)} < x_2^{(m)} < \cdots < y_{m+n(m)+1-d(m)}^{(m)} < x_{m+n(m)+2-d(m)}^{(m)},$$

and set $y_0^{(m)} := x_1^{(m)}$, $y_{m+n(m)+2-d(m)}^{(m)} := x_{m+n(m)+2-d(m)}^{(m)}$. Then $|y_k^{(m)} - y_{k+1}^{(m)}| \geq t_m$ for some $k = k^*$. As above, counting the zeros of $R_m \pm (p_{m-n(m)} - 1)/2$, where $p_{m-n(m)}$ satisfies Lemma 2.1 with $\alpha = y_{k^*}^{(m)}$ and $\beta = y_{k^*+1}^{(m)}$, yields

$$(2.18) \quad \frac{1}{2} (c_1 e^{c_2 t_m(m-n(m))} - 1) < m^2.$$

By (2.16) or (2.18) we get for a constant $c_3 > 0$

$$(2.19) \quad t_m(m - n(m)) \leq c_3 \log m,$$

which yields (1.7). \square

Proof of Theorem 1.2. It suffices to prove (1.11) for the case $\alpha = -1$. In fact, it is enough to show that

$$(2.20) \quad \limsup_{\substack{m \rightarrow \infty \\ m \in \Omega}} \frac{N_{m,n(m)}(-1, \beta)}{N_{m,n(m)}(-1, 1)} \leq \frac{\pi - \arccos \beta}{\pi},$$

since replacing x by $-x$ and β by $-\beta$, we get the corresponding lower estimate for \liminf . Let $m - n(m) = s(m) + l(m)$, where $s(m)$ is to be determined later. With the notations of the previous proof, set for $m \in \Omega$,

$$(2.21) \quad q_m(x) := \frac{1}{2} T_{s(m)}(x + 1 - \beta) T_{l(m)}(x),$$

$$(2.22) \quad N(m) := \#\{x \in (\beta, 1]: |T_{l(m)}(x)| = 1, |q_m(x)| > m^2\},$$

where T_k denotes the k th the Chebyshev polynomial, $\|T_k\|_{[-1,1]} = 1$. Then $R_m - q_m$ has at least $N(m) - 1$ zeros in $(\beta, 1]$. Thus it can have at most $m + n(m) + 2 - d(m) - N(m)$ zeros in $[-1, \beta]$. Hence

$$(2.23) \quad \limsup_{m \rightarrow \infty} \frac{N_{m,n(m)}(-1, \beta)}{N_{m,n(m)}(-1, 1)} \leq \limsup_{m \rightarrow \infty} \frac{m + n(m) + 2 - d(m) - N(m)}{m + n(m) + 2 - d(m)} \\ = 1 - \liminf_{m \rightarrow \infty} \frac{N(m)}{m},$$

since between two alternation points of R_m in $[-1, \beta]$ is one zero of $R_m - q_m$ and since $n(m)/m \rightarrow 0$ ($d(m) \leq n(m)$). In (2.23) and the rest of the proof, all limits are for $m \in \Omega$. Now choose $s(m)$ such that

$$(2.24) \quad \lim_{m \rightarrow \infty} \frac{s(m)}{\log m} = \infty, \quad \lim_{m \rightarrow \infty} \frac{l(m)}{m} = 1.$$

Then the first equation in (2.24) together with (2.5) yields

$$(2.25) \quad \lim_{m \rightarrow \infty} (\inf\{x \in (\beta, 1]: \frac{1}{2} T_{s(m)}(x + 1 - \beta) > m^2\}) = \beta.$$

Also, for $\beta < \tilde{\beta} < 1$, it follows from the second equation in (2.24) that

$$(2.26) \quad \lim_{m \rightarrow \infty} \frac{\#\{x \in (\tilde{\beta}, 1]: |T_{l(m)}(x)| = 1\}}{m} = \frac{\arccos \tilde{\beta}}{\pi}.$$

Finally, (2.25) and (2.26) yield (with (2.21) and (2.22))

$$(2.27) \quad \liminf_{m \rightarrow \infty} \frac{N(m)}{m} \geq \frac{\arccos \beta}{\pi},$$

which together with (2.23) gives (2.20). \square

3. PROOF OF THEOREM 1.3

For $\alpha_1 \leq \dots \leq \alpha_n < 0$ we denote by $V_{n-1}(\alpha_1, \dots, \alpha_n) = x^{n-1} + \dots$ the monic polynomial that minimizes

$$(3.1) \quad \left\| \frac{p_{n-1}(x)}{\prod_{k=1}^n (x - \alpha_k)} \right\|_{[0,1]}$$

among all monic polynomials p_{n-1} of degree $n-1$. Set

$$(3.2) \quad r_{n-1}(\alpha_1, \dots, \alpha_n)(x) := \frac{V_{n-1}(\alpha_1, \dots, \alpha_n)(x)}{\prod_{k=1}^n (x - \alpha_k)}.$$

Then from the Haar condition (cf. [M, §3.2]), $r_{n-1}(\alpha_1, \dots, \alpha_n)$ is uniquely determined and equioscillates n times in $[0, 1]$. Moreover, these n equioscillation points are the only extremal points of $r_{n-1}(\alpha_1, \dots, \alpha_n)$ in $[0, 1]$, since $r_{n-1}(\alpha_1, \dots, \alpha_n) - c$ can have at most n zeros for each $c \in \mathbb{R}$ and since $r_{n-1}(\alpha_1, \dots, \alpha_n)$ has all its $n-1$ zeros in $[0, 1]$. Also, zero must be one of the equioscillation points, since $|r_{n-1}(\alpha_1, \dots, \alpha_n)(x)|$ decreases between α_n and the first zero of $r_{n-1}(\alpha_1, \dots, \alpha_n)$.

We need the following lemmas.

Lemma 3.1. *Let $\alpha_1 \leq \dots \leq \alpha_n < 0$ and $\beta_1 \leq \dots \leq \beta_n < 0$ satisfy $\alpha_k \leq \beta_k$ for $k = 1, \dots, n$. Then the equioscillation points $x_1 < \dots < x_n$ of $r_{n-1}(\alpha_1, \dots, \alpha_n)$ and $y_1 < \dots < y_n$ of $r_{n-1}(\beta_1, \dots, \beta_n)$ satisfy $x_k \geq y_k$ for $k = 1, \dots, n$.*

Proof. It suffices to prove the lemma in the case $\alpha_k = \beta_k$ for $k \neq k^*$, $\alpha_{k^*} < \beta_{k^*}$, since we can transfer α_n to β_n, \dots, α_1 to β_1 successively. Define

$$(3.3) \quad C_\alpha := 1/\|r_{n-1}(\underline{\alpha})\|_{[0,1]},$$

$$(3.4) \quad C_\beta := 1/\|r_{n-1}(\underline{\beta})\|_{[0,1]},$$

where $\underline{\alpha} = (\alpha_1, \dots, \alpha_n)$, $\underline{\beta} = (\beta_1, \dots, \beta_n)$, and set

$$(3.5) \quad \begin{aligned} q(x) &:= C_\alpha(x - \beta_{k^*})V_{n-1}(\underline{\alpha})(x) - C_\beta(x - \alpha_{k^*})V_{n-1}(\underline{\beta})(x) \\ &= (C_\alpha r_{n-1}(\underline{\alpha})(x) - C_\beta r_{n-1}(\underline{\beta})(x))(x - \alpha_{k^*})(x - \beta_{k^*}) \prod_{k \neq k^*} (x - \alpha_k). \end{aligned}$$

By the equioscillation, we have ($y_1 = 0$)

$$(3.6) \quad q(y_n) \leq 0, q(y_{n-1}) \geq 0, \dots, q(0) = 0.$$

It is easy to see that $C_\alpha > C_\beta$. Thus there is a point $x > y_n$ with $q(x) > 0$. For the (necessarily real) zeros $\xi_1 \leq \dots \leq \xi_n$ of q (some zeros may be counted twice), this implies $\xi_k \geq y_k$ for $k = 1, \dots, n$. We also have ($x_1 = 0$)

$$(3.7) \quad q(x_n) \geq 0, q(x_{n-1}) \leq 0, \dots, q(0) = 0.$$

For every $\varepsilon > 0$, there is a polynomial $\tilde{q} \in \Pi_n$ with highest coefficient $C_\alpha - C_\beta$ and real zeros $\tilde{\xi}_1 \leq \dots \leq \tilde{\xi}_n$ such that

$$(3.8) \quad \tilde{q}(x_n) > 0, \tilde{q}(x_{n-1}) < 0, \dots, \tilde{q}(0) < 0 \quad \text{for } n \text{ even}, \\ \tilde{q}(0) > 0 \quad \text{for } n \text{ odd},$$

$$(3.9) \quad |\tilde{\xi}_k - \xi_k| < \varepsilon, \quad k = 1, \dots, n.$$

It follows that $\tilde{\xi}_1 < 0$ and thus $x_k > \tilde{\xi}_k$ for $k = 1, \dots, n$. Since $\varepsilon > 0$ is arbitrary, this implies $x_k \geq \xi_k \geq y_k$ for $k = 1, \dots, n$. \square

Lemma 3.2. *Given $0 < \varepsilon < 1$ there is an increasing sequence $a_1 < a_2 < \dots < 0$, such that $r_{n-1}(a_1, \dots, a_n)$ has no extremal points in $(\varepsilon, 1]$ for $n \geq 1$.*

Proof. Set $a_1 := -\varepsilon/4$. We will construct a_n by induction such that the function

$$(3.10) \quad f_n(x) := \frac{\prod_{k=2}^n (x + 2a_k)}{\prod_{k=1}^n (x - a_k)}$$

alternates in sign in the points

$$(3.11) \quad 0 = \delta_{1,n} < \dots < \delta_{n,n} < \varepsilon$$

and satisfies

$$(3.12) \quad |f_n(\delta_{k,n})| > |f_n(x)| \quad \text{for } k = 1, \dots, n, \quad x \in [\varepsilon, 1].$$

If we have this sequence, $r_{n-1}(a_1, \dots, a_n)$ cannot have an alternation point in $(\varepsilon, 1]$ for $n \geq 2$, since otherwise for a suitably chosen $\gamma \in \mathbb{R}$ the function $f_n - \gamma r_{n-1}(a_1, \dots, a_n)$ has a zero in each interval $(\delta_{k,n}, \delta_{k+1,n})$ and an additional zero in $(\delta_{n,n}, 1)$.

We observe now that $f_1(x) = 1/(x - a_1)$ is decreasing in $[0, 1]$ and satisfies (3.12) with $\delta_{1,1} = 0$. Having constructed a_1, \dots, a_{n-1} , we observe that

$$(3.13) \quad \lim_{a_n \rightarrow 0^-} \frac{x + 2a_n}{x - a_n} = 1 \text{ uniformly on } [\lambda, 1]$$

for all $\lambda > 0$. Thus, for $|a_n|$ sufficiently small, (3.12) will be satisfied for $\delta_{k,n+1} := \delta_{k-1,n}$, $k = 3, \dots, n+1$ and for $\delta_{1,n+1} := \delta_{1,n} = 0$. Thus it remains to show the existence of $\delta_{2,n+1}$. Since $f_{n+1}(0) = -2f_n(0)$, this follows from (3.13) by choosing $|a_n|$ small enough. \square

Proof of Theorem 1.3. We will prove the theorem on the interval $[0, 1]$. Choose a_1, a_2, \dots as in Lemma 3.2 with $\varepsilon/2$ replacing ε . Let, for $b_k > 0$,

$$(3.14) \quad S_n(x) := \sum_{k=1}^n \frac{b_k}{x - a_{2k}}.$$

We now use a result in [B] stating that the best approximation to S_n out of $\mathcal{R}_{n-2, n-1}$ has the form

$$(3.15) \quad r_{n-2, n-1}^*(S_n)(x) = \frac{p_{n-2}^*(x)}{\prod_{k=1}^{n-1} (x - a_{2k+1}^*)},$$

where p_{n-2}^* is of degree $n-2$ and

$$(3.16) \quad a_2 < a_3^* < a_4 < \cdots < a_{2n-1}^* < a_{2n} < 0.$$

Thus, by the equioscillation property (1.4), there is a constant c_n such that

$$(3.17) \quad S_n - r_{n-2, n-1}^*(S_n) = c_n r_{2n-2}(a_2, a_3^*, a_4, \dots, a_{2n}).$$

Since $r_{2n-2}(a_1, \dots, a_{2n-1})$ has no alternation point in $(\varepsilon/2, 1]$, Lemma 3.1 shows that $S_n - r_{n-2, n-1}^*(S_n)$ has no alternation point in $(\varepsilon/2, 1]$. We choose the b_k 's such that

$$(3.18) \quad \text{the series } f(x) = \sum_{k=1}^{\infty} \frac{b_k}{x - a_{2k}} \text{ converges uniformly on } [0, 1],$$

and

$$(3.19) \quad \begin{aligned} &r_{n-2, n-1}^*(f) \text{ is close enough to } r_{n-2, n-1}^*(S_n) \text{ to guarantee that} \\ &f - r_{n-2, n-1}^*(f) \text{ has no alternation point in } (\varepsilon, 1]. \end{aligned}$$

For (3.19) we used the fact (cf. [W]) that the best approximation operator is continuous in S_n , since $r_{n-2, n-1}^*(S_n)$ is nondegenerate (i.e. $d = 0$ in (1.3)).

□

REFERENCES

- [B] M. Barnsley, *The bounding properties of the multipoint Padé approximant to a series of Stieltjes on the real line*, J. Math. Phys. **14** (1973), 299–313.
- [K] M. I. Kadec, *On the distribution of points of maximal deviation in the approximation of continuous functions by polynomials*, Uspekhi Mat. Nauk. **15** (1960), 199–202.
- [KP] A. Kroó and F. Peherstorfer, *Interpolating properties of best rational L_1 -approximations*, Constr. Approx. **4** (1988), 97–106.
- [KS] A. Kroó and E. B. Saff, *The density of extreme points in complex polynomial approximation*, Proc. Amer. Math. Soc. **103** (1988), 203–209.
- [L] G. G. Lorentz, *Distribution of alternation points in uniform polynomial approximation*, Proc. Amer. Math. Soc. **92** (1984), 401–403.
- [M] G. Meinardus, *Approximation theory: Theory and numerical methods*, Springer tracts in natural philosophy, Vol. 13, Springer-Verlag, New York, 1967.
- [W] H. Werner, *On the rational Tschebyscheff operator*, Math. Z. **86** (1964), 317–326.

DEPARTMENT OF MATHEMATICS, DALHOUSIE UNIVERSITY, HALIFAX, NOVA SCOTIA, CANADA B3H 4H8

Current address (A. KROÓ): MATHEMATICAL INSTITUTE OF THE HUNGARIAN ACADEMY OF SCIENCES, BUDAPEST, REÁLTANODA U. 13-15, H-1053 HUNGARY

Current address (R. GROTHMANN): MATHEM. GEOGR. FAKULTÄT, KATHOLISCHE UNIVERSITÄT EICHSTÄTT, OSTENSTRASSE 18, D-8078 EICHSTÄTT, WEST GERMANY

Current address (E. B. SAFF): INSTITUTE FOR CONSTRUCTIVE MATHEMATICS, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTH FLORIDA, TAMPA, FLORIDA 33620 USA