

## ON THE ENTROPY NORM SPACES AND THE HARDY SPACE $\text{Re } H^1$

W. C. LANG

(Communicated by J. Marshall Ash)

**ABSTRACT.** R. Dabrowski introduced certain natural multiplier operators which map from the entropy norm spaces of B. Korenblum into the Hardy space  $\text{Re } H^1$ . We show that the images of the entropy norm spaces in  $\text{Re } H^1$  do not include all of that space.

### 1. INTRODUCTION

We consider the entropy norm spaces of Korenblum [4]. He defined an entropy function  $\kappa : [0, 1] \rightarrow [0, 1]$  to be a concave, continuous, increasing function with  $\kappa(0) = 0$ . We denote by  $K_0$  the set of such functions such that  $\kappa'(0) = \lim_{x \rightarrow 0^+} \kappa(x)/x = \infty$ . According to Dabrowski [1] to each  $\kappa \in K_0$  there is a unique probability measure  $\mu = \mu_\kappa$  such that

$$\kappa(x) = \int_0^x \int_t^1 \frac{d\mu(u)}{u} dt.$$

Then the entropy norm of a continuous 1-periodic function  $f \in C(T)$  (where  $T = \mathbb{R} \bmod 1$ ) is given by

$$\|f\|_\kappa = \int_0^1 \int_T \Omega_I(f) dt \frac{d\mu(s)}{s}$$

where  $I = [t - s/2, t + s/2]$  and where  $\Omega_I(f) = \sup\{|f(u) - f(v)| : u, v \in I\}$ . (This norm was introduced by Korenblum [4]; this formula for the norm is due to Dabrowski [4].) We denote by  $C_\kappa \subseteq C(T)$  the space of continuous 1-periodic functions of finite entropy norm.

In [2], Dabrowski introduced an operator  $T_\kappa : C_\kappa \rightarrow \text{Re } H^1$ , given by

$$T_\kappa f(t) = \int_T \int_0^1 \frac{\chi_I(t)}{s^2} (f(t) - f(I)) d\mu(s) dx$$

where  $I = [x - s/2, x + s/2]$ ,  $f(I) = \frac{1}{|I|} \int_I f(t) dt$  is the average of  $f$  over  $I$ , and  $\chi_I$  is the usual characteristic function of  $I$ . He showed that  $T_\kappa$  is a

---

Received by the editors November 6, 1991; presented March 20, 1992 in the Special Session on Harmonic Analysis of the 873rd Meeting of the AMS, at Southwest Missouri State University.

1991 *Mathematics Subject Classification.* Primary 42A20, 46E15.

*Key words and phrases.* Entropy norm spaces, real Hardy space.

multiplier with coefficients

$$\beta_n = \beta_n(\kappa) = \frac{1}{2\pi^2 n^2} \int_{(0,1]} (\cos(2\pi ns) - 1 + 2\pi^2 n^2 s^2) \frac{1}{s^3} d\mu_\kappa(s)$$

(for  $n > 0$  we set  $\beta_{-n} = \beta_n$  and  $\beta_0 = 0$ ). In [3], Dabrowski asked the question: given  $f \in \text{Re } H^1$ , are there  $\kappa \in K_0$  and  $g \in C_\kappa$  such that  $f = T_\kappa g$ ? (One reason why this question is of interest is because, as Dabrowski remarks, a positive answer would imply the Fefferman duality  $(\text{Re } H^1(0))^* = \text{BMO}$ .)

2. THE MAIN RESULT

We are ready to give a negative answer to this question.

**Theorem.** *There is a function  $f \in \text{Re } H^1$  such that there are no  $\kappa \in K_0$  and  $g \in C_\kappa$  with  $f = T_\kappa g$ .*

*Proof.* We construct  $f$  as follows. Let  $h$  be the function with Fourier series  $\sum_{n=1}^\infty (\sqrt{n} \log(n+1))^{-1} e_n$ , where  $e_n = e^{2\pi i n t}$ . Then  $h \in H^2$ . So  $h^2 \in H^1$  (see, e.g., Zygmund [6, VII (7.22), p. 275]). We let  $f = \text{Re}(h^2)$ . So of course  $f \in \text{Re } H^1$ . We have

$$h^2 \sim \sum_{n=1}^\infty \left( \sum_{j=1}^{n-1} b_j b_{n-1} \right) e_n$$

where  $b_j = (\sqrt{j} \log(j+1))^{-1}$ . It is not hard to show that  $f$  has Fourier series  $\sum_{n=1}^\infty a_n \cos(2\pi n t)$  where  $a_n \geq \text{const.}(\log(n+1))^{-2}$  for  $n = 1, 2, 3, \dots$ .

Now we suppose that there is a  $\kappa \in K_0$  and a  $g \in C_\kappa$  such that  $T_\kappa g = f$ . We write  $g$  as  $\sum c_n e_n$ . Then since  $T_\kappa g = f$  we have  $c_n = a_n / \beta_n$ ,  $n \geq 1$ . This enables us to write  $g$  as  $\sum_1^\infty c_n \cos(2\pi n t)$  where  $c_n \geq 0$  for all  $n > 0$ .

We assume that  $g \in C_\kappa$  which implies that  $g$  is bounded. Consequently (since  $g$  has a cosine series with positive coefficients), we must have  $\sum c_n < \infty$  or  $\sum a_n / \beta_n < \infty$ . Therefore

$$(1) \quad \sum_{n=1}^\infty \left( \frac{1}{\log(n+1)} \right)^2 \frac{1}{\beta_n} < \infty.$$

We must also have

$$(2) \quad \sum_{n=1}^\infty \frac{1}{n^2} \beta_n < \infty.$$

[By Lang [5],  $\beta_n$  compares with  $n\kappa(1/n) - n^2 \int_0^{1/n} \kappa(t) dt = \bar{\kappa}'(1/n)$  where  $\bar{\kappa}(x) = \frac{1}{x} \int_0^x \kappa(t) dt$ . We have  $\bar{\kappa}(x) = \int_0^x \bar{\kappa}'(t) dt$ , so this integral must be convergent; we may estimate this integral by the sum

$$\sum_{n=1}^\infty \left( \frac{1}{n} - \frac{1}{n+1} \right) \bar{\kappa}' \left( \frac{1}{n} \right) \approx \sum_{n=1}^\infty \frac{1}{n^2} \beta_n.$$

(Note that  $\bar{\kappa}'(x) = (1/x^2)(x\kappa(1/x) - \int_0^x \kappa(t) dt)$  is the product of  $1/x^2$  and a function which goes to 0 monotonically as  $x \rightarrow 0$ . So the integral and the sum compare.)]

But (1) and (2) are not compatible. Indeed, suppose the sums (1) and (2) are both finite. Then by the Cauchy-Schwarz inequality

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{1}{n} \frac{1}{\log(n+1)} &= \sum_{n=1}^{\infty} \left( \frac{1}{n} \sqrt{\beta_n} \right) \left( \frac{1}{\log(n+1)} \frac{1}{\sqrt{\beta_n}} \right) \\ &\leq \left( \sum_{n=1}^{\infty} \frac{1}{n^2} \beta_n \right)^{1/2} \left( \sum_{n=1}^{\infty} \left( \frac{1}{\log(n+1)} \right)^2 \frac{1}{\beta_n} \right)^{1/2} < \infty, \end{aligned}$$

which is nonsense. So there cannot be  $\kappa \in K_0$ ,  $g \in C_\kappa$  such that  $T_\kappa g = f$ , and we are done.  $\square$

### REFERENCES

1. R. Dabrowski, *Probability measure representation of norms associated with the notion of entropy*, Proc. Amer. Math. Soc. **90** (1984), 263–268.
2. ———, *On Fourier coefficients of a continuous periodic function of bounded entropy norm*, Bull. Amer. Math. Soc. (N.S.) **18** (1988), 49–51.
3. ———, *On a natural connection between the entropy spaces and Hardy space  $\text{Re } H^1$* , Proc. Amer. Math. Soc. **104** (1988), 812–818.
4. B. Korenblum, *On a class of Banach spaces associated with the notion of entropy*, Trans. Amer. Math. Soc. **290** (1985), 527–553.
5. W. C. Lang, *A growth condition for Fourier coefficients of a function of bounded entropy norm*, Proc. Amer. Math. Soc. **112** (1991), 433–439.
6. A. Zygmund, *Trigonometric series*, 2nd ed., vol. I, Cambridge Univ. Press, Cambridge, 1959.

DEPARTMENT OF MATHEMATICS AND STATISTICS, MISSISSIPPI STATE UNIVERSITY, MISSISSIPPI STATE, MISSISSIPPI 39762

*E-mail address*: lang@math.msstate.edu