

## SIEGEL'S THEOREM FOR COMPLEX FUNCTION FIELDS

JOSÉ FELIPE VOLOCH

(Communicated by William Adams)

**ABSTRACT.** We give a short proof of the finiteness of the set of integral points on an affine algebraic curve of genus at least one, defined over a function field of characteristic zero.

Siegel [Si] has shown that an affine algebraic curve of genus at least one defined over a number field has only finitely many integral points. Lang [L] has proven an analogous result for curves defined over a function field of characteristic zero not defined over the constant field. For curves of genus at least two, one even has the Mordell conjecture (proved by Faltings [F] in the number field case and by Manin [M] in the function field case) that there are only finitely many rational points.

For genus one, Manin [M] gave a proof of a strengthening of Lang's result as a by-product of his work on the Mordell conjecture. Mason [Ms] then gave an effective proof by more elementary considerations. In this note, we give a short proof of Manin's (and hence Lang's) result for genus one. The proof can be adapted to higher genus as well (see the remark below).

Let  $K$  be a function field with constant field of characteristic zero and  $E/K$  an elliptic curve with nonconstant  $j$ -invariant. The reader may consult [S] for definitions and results about elliptic curves. In particular, we shall use the following results. The group  $E(K)$  is a finitely generated abelian group by the Mordell-Weil theorem, and there is a height function  $h : E(K) \rightarrow \mathbb{R}$  with the property that there are only finitely many points of bounded height ([L], Proposition 2). The height can be written as a sum of local heights  $\sum \lambda_v(P)$ , where  $v$  ranges through the places of  $K$ . The local heights satisfy  $\lambda_v(P) = \max\{0, v(t(P))\} + \beta_v(P)$ , where  $\beta_v$  is bounded for all  $v$  and is identically zero for all but finitely many  $v$ , and  $t$  is a uniformizer at  $0 \in E$ . For example,  $t = x/y$ , where  $x, y$  are coordinates of a Weierstrass equation for  $E$ .

Now, Lang's result for genus one can be reduced to the case of a Weierstrass equation ([S], Corollary IX.3.2.2) and in this case we argue as follows. Let  $S$  be a finite set of places of  $K$ . Then  $\sum_{v \notin S} \lambda_v(P)$  is bounded independently of  $P$ , if  $P$  is  $S$ -integral and, since there are only finitely many points of bounded

---

Received by the editors October 5, 1992 and, in revised form, March 20, 1993.

1991 *Mathematics Subject Classification*. Primary 11G05, 11G30, 14G25.

*Key words and phrases*. Algebraic curves, function fields, integral points.

©1994 American Mathematical Society  
0002-9939/94 \$1.00 + \$.25 per page

height, if there are infinitely many  $S$ -integral points, then  $\lambda_v$  is unbounded for some  $v \in S$ . It suffices thus to prove the following result:

**Theorem (Manin).** *Let  $K$  be a function field with constant field of characteristic zero and  $E/K$  an elliptic curve with nonconstant  $j$ -invariant and  $v$  a place of  $K$ . Then the local height function  $\lambda_v$  is bounded on  $E(K)$ .*

*Proof.* The points on  $E(K_v)$  that reduce to  $0 \pmod{v}$  form a subgroup  $E_1(K_v)$  isomorphic to the group of points of a formal group. Choosing an uniformizer  $t$ , as above, on  $E$  at 0, then  $E_1(K_v) = \{P \in E(K_v) | t(P) \in \mathcal{M}_v\}$ , where  $\mathcal{M}_v$  is the maximal ideal of the local ring at  $v$ . Moreover,  $\lambda_v(P)$  differs from  $v(t(P))$  by a bounded amount. Hence, it suffices to show that  $v(t(P))$  is bounded above on  $E(K)$ . Suppose not and choose  $P_n, n = 1, 2, \dots$ , in  $E(K)$  such that  $v(t(P_{n+1})) > v(t(P_n)) > 0$ . We claim that  $P_1, P_2, \dots$  are linearly independent over  $\mathbb{Z}$ . Recall that  $t$  induces a group isomorphism between  $E_r/E_{r+1}$ , where  $E_r = E_r(K_v) = \{P \in E(K_v) | t(P) \in \mathcal{M}_v^r\}$ , and  $\mathcal{M}_v^r/\mathcal{M}_v^{r+1}$ . If  $n_i P_i = \sum_{j>i} n_j P_j$ ,  $n_i \neq 0$ , and  $r = v(t(P_i))$  then  $n_i P_i$  is 0 in  $E_r/E_{r+1}$ , but  $t(n_i P_i) \equiv n_i t(P_i) \not\equiv 0 \pmod{\mathcal{M}_v^{r+1}}$ , which proves the claim. On the other hand, the claim contradicts the Mordell-Weil theorem and this completes the proof.

*Remark.* On a curve of genus greater than one, if a sequence of points  $P_1, P_2, \dots$  approaches rapidly a point  $P_\infty$ , then a similar argument shows that the  $P_i - P_\infty$  are linearly independent over  $\mathbb{Z}$  in the Jacobian of the curve, and Lang's result follows from this. The author and A. Buium [BV] have recently proved a conjecture of Lang to the effect that an affine open subset of an abelian variety of any dimension over a function field of characteristic zero has finitely many integral points.

## REFERENCES

- [BV] A. Buium and J. F. Voloch, *Integral points of abelian varieties over function fields of characteristic zero*, Math. Ann. **297** (1993), 303–307.
- [F] G. Faltings, *Endlichkeitssätze für abelsche Varietäten über Zahlkörpern*, Invent. Math. **73** (1983), 349–366.
- [L] S. Lang, *Integral points on curves*, Inst. Hautes Études Sci. Publ. Math. **6** (1960), 27–43.
- [M] Yu. I. Manin, *Rational points on an algebraic curve over function fields*, Izv. Acad. Nauk USSR (1963) and Letter to the editor, Math. USSR-Izv. **34** (1990), 465–466; English transl., Transl. Amer. Math. Soc. Ser. 2, vol. 50, Amer. Math. Soc., Providence, RI, 1966, pp. 189–234.
- [Ms] R. C. Mason, *Diophantine equations over function fields*, London Math. Soc. Lecture Note Ser., vol. 96, Cambridge Univ. Press, Cambridge, 1984.
- [S] J. H. Silverman, *The arithmetic of elliptic curves*, Graduate Texts in Math., vol. 106, Springer, New York, 1986.
- [Si] C. L. Siegel, *Einige Anwendungen diophantischer Approximationen*, Abh. Preuss. Akad. Wiss. Phys. Math. Kl. **1** (1929), 41–69.