

REMOVABILITY OF THE SINGULAR SET OF THE HEAT FLOW OF HARMONIC MAPS

YUNMEI CHEN AND LIVIO FLAMINIO

(Communicated by Peter Li)

ABSTRACT. We show that the singular set of a weak stationary solution u of the heat flow of harmonic maps between Riemannian manifolds M and N , with N compact, is removable if it has “parabolic codimension” greater than two and the initial energy $E(u_0)$ is sufficiently small.

1. INTRODUCTION AND PRELIMINARIES

Let (M, g) and (N, h) be smooth Riemannian manifolds of dimensions m and n , respectively, with N compact without boundary. Using the Nash-Moser Theorem, we embed N isometrically into \mathbb{R}^K .

We recall that the energy $E(u)$ of a smooth map $u: M \rightarrow N$ is defined as

$$E(u) = \frac{1}{2} \int_M |du|^2 dm_g,$$

where dm_g is the volume form induced by g and $|du|$ is the Hilbert-Schmidt norm of du considered as a section of $TM^* \otimes u^*TN$. A smooth function $u: M \rightarrow N$ is said to be harmonic if it is a critical point of the energy, i.e. if $d^*du = 0$ (see [5]). The *heat flow* for harmonic maps, with initial data $u_0: M \rightarrow N$, is the solution to the evolution equation corresponding to the harmonic map problem:

$$(\partial_t + d^*d)u = 0, \quad u(0, \cdot) = u_0(\cdot).$$

Using the embedding $N \rightarrow \mathbb{R}^K$ the above equation can be rewritten as

$$(1.1) \quad \partial_t u + \Delta_M u = \text{Tr}_g A_u(du, du), \quad u(0, \cdot) = u_0(\cdot),$$

where Δ_M is the Laplace-Beltrami operator of (M, g) ; the expression $A_p(\cdot, \cdot)$ denotes the second fundamental form at $p \in N$ of the embedding $N \rightarrow \mathbb{R}^K$, and $\text{Tr}_g A_u(du, du)$ at (t, x) is the trace of $A_{u(t, x)}(du(t, x), du(t, x))$ with respect to g_x , which is a bilinear form on $T_x M$ with values in \mathbb{R}^K . The advantage of the form (1.1) for the heat flow equation is that it allows us to formulate the problem in a

Received by the editors September 4, 1994.

1991 *Mathematics Subject Classification*. Primary 35B65, 35D10, 49N60, 35Kxx, 58E20, 58G11.

Key words and phrases. Heat flow, harmonic maps.

The first author was supported by NSF grant #DMS-9101911.

weak sense. Define the space

$$H(\mathbb{R}^+ \times M, N) = \{u \in L^\infty(\mathbb{R}^+ \times M, \mathbb{R}^K) : u(t, x) \in N \text{ a.e.}, \\ \partial_t u \in L^2(\mathbb{R}^+ \times M, \mathbb{R}^K), du \in L^\infty(\mathbb{R}^+, L^2(M, \mathbb{R}^K))\}.$$

Then a *weak solution of the heat flow* (1.1) is a function $u \in H(\mathbb{R}^+ \times M, N)$ that satisfies

$$(1.2) \quad \int_{\mathbb{R}^+ \times M} \{\langle \partial_t u, \eta \rangle + \langle du, d\eta \rangle - \langle \eta, \text{Tr}_g A(du, du) \rangle\} dm_g dt$$

for any $\eta \in C_0^\infty(\mathbb{R}^+ \times M, \mathbb{R}^K)$ as well as the initial condition $u(0, \cdot) = u_0(\cdot)$.

The global existence of a weak solution to the problem (1.1) is known; see for example [3]. The weak solution obtained in [3] is smooth outside a set $\Sigma(u) \subseteq \mathbb{R}^+ \times M$, the *singular set*, which is closed and has locally finite m -dimensional Hausdorff measure (here and hereafter the Hausdorff measure on $\mathbb{R}^+ \times M$ is defined using the parabolic metric defined in §2). However, Coron showed in [4] that the problem (1.1) may have infinitely many distinct weak solutions. Therefore the partial regularity of weak solutions becomes an interesting problem.

In analogy with the notion of stationary harmonic map (see for example [12], [13]), Feldman introduced in [7] the notion of *stationary solution* of the heat flow. The stationarity condition can be formulated in the following way.

A vector field X on $\mathbb{R}^+ \times M$ is *future directed* if $dt(X) \geq 0$, where t is the first coordinate function on $\mathbb{R}^+ \times M$.

If $Z = (T, X)$ is a C^1 vector field of compact support on $\mathbb{R}^+ \times M$, define $\tilde{u}_{Z, \varepsilon} : \mathbb{R}^+ \times M \rightarrow N$ as

$$\tilde{u}_{Z, \varepsilon}(t, x) = u(\exp_{t, x}(\varepsilon Z(t, x))) = u(t + \varepsilon T(t, x), \exp_x(\varepsilon X(t, x))).$$

1.1 Definition ([7]). A weak solution u of the heat flow (1.1) is said to be *stationary* if for any C^1 future directed vector field Z on $\mathbb{R}^+ \times M$ with compact support we have

$$(1.3) \quad \int_{\mathbb{R}} \int_M \left\langle \partial_t u, \frac{\partial}{\partial \varepsilon} u_{Z, \varepsilon} \Big|_{\varepsilon=0} \right\rangle dm_g dt + \frac{\partial}{\partial \varepsilon} \left(\int_{\mathbb{R}^+} E(u_{Z, \varepsilon}) dt \right) \Big|_{\varepsilon=0} \leq 0.$$

In [7], Feldman proved that if the target manifold N is the standard n -dimensional sphere, then the singular set of a weak stationary solution u of (1.1) has zero m -dimensional Hausdorff measure. Independently, Chen, Li and Lin [2] obtained the same result by assuming, instead of stationarity, the monotonicity inequality and energy inequality (see (2.1) and (2.7) in [2]).

Extending previous partial regularity results of Hélein [8], [9] and Evans [6], Bethuel [1] proved that for stationary harmonic maps the singular set has zero m -dimensional Hausdorff measure, provided that the solution is stationary.

Thus one would expect that for the heat flow the singular set of a weak stationary solution of the heat flow into arbitrary target manifolds also has zero m -dimensional Hausdorff measure. The purpose of this note is a partial result in this direction.

Roughly speaking, we will show that if a weak stationary solution u of the heat flow has small initial energy $E(u_0)$ and if its singular set $\Sigma(u)$ has co-dimension greater than 2, then u is smooth. More precisely we have

1.2 Theorem. *Let $u \in H(\mathbb{R}^+ \times M, N)$ be a weak solution of the heat flow (1.1). Assume that:*

- (1) *The weak solution u is stationary.*

- (2) The singular set $\Sigma(u)$ of u satisfies the following regularity conditions:
 - (a) for some strictly positive $\delta_0 < 1$, the set $\Sigma(u)$ is contained in $M \times [\delta_0^2, \infty)$;
 - (b) the set $\Sigma(u)$ has bounded uniform (m, q) -density with respect to the parabolic distance for $q > 2$.
- (3) For some $\varepsilon = \varepsilon(m, N, \delta_0)$ we have $E(u_0) < \varepsilon$.

Then u is smooth in M .

For the definition of (m, q) -density see Definition 2.3 below. The condition of stationarity can be replaced by assuming the monotonicity inequality (4.5) below and the energy inequality [2, (2.7)]. In fact, in [2] it was shown that these inequalities imply the estimates (4.2) and (4.4) below from which Theorem 1.2 follows.

In §3 we shall prove that the condition of stationarity is automatically satisfied if we assume that the codimension q of the singular set is strictly greater than 2 and that $|du|$ has higher integrability. See Theorem 3.1 below for a precise statement.

Our assumptions on the singular set are similar to Mou’s [11] who proved analogous results for p -harmonic maps.

1.3 *Notation.* Let $\mathcal{L}^{m+1} = dm_g dt$. Whenever not explicitly indicated, integration on subsets of M and of $\mathbb{R}^+ \times M$ will always be intended with respect to the measures dm_g and \mathcal{L}^{m+1} . By an *absolute constant* we intend a constant which may depend only on (M, g) and (N, h) . Occasionally we shall write $a \ll b$ to mean that for some absolute constant C we have $a < Cb$.

1.4 *Notation.* For $u \in H(\mathbb{R}^+ \times M, N)$ we set $u(t) = u(t, \cdot)$.

1.5 *Notation.* The metric $dt^2 \otimes g$ induces a connection $\vec{\nabla}$ on $\mathbb{R}^+ \times M$, for which the hypersurfaces $\{t\} \times M$ are totally geodesics. In particular the pull back of g to $\mathbb{R}^+ \times M$ (denoted g again) is parallel. We shall denote by $\vec{d}u$ the differential of $u: \mathbb{R}^+ \times M \rightarrow N$, while reserving the symbol du to denote the restriction of $\vec{d}u$ to $\{t\} \times TM$. Similarly $\vec{\delta}Z$ denotes the divergence $\vec{\delta}Z = -\text{Tr } \vec{\nabla}Z$.

1.6 *Remark.* The condition of stationarity yields immediately that $u \in H(\mathbb{R}^+ \times M, N)$ is a stationary solution of the heat flow iff for any C^1 -future directed vector field Z on $\mathbb{R}^+ \times M$ with compact support we have:

$$(1.4) \quad \int_{\mathbb{R}^+ \times M} \left\{ \langle \partial_t u, \vec{d}u(Z) \rangle + \langle du, \vec{d}u(\vec{\nabla}Z) \rangle + \frac{1}{2} \langle du, du \rangle \vec{\delta}Z \right\} \leq 0,$$

where $\langle \cdot, \cdot \rangle$ denotes the inner product on any of the various tensor bundles $T^{(r,s)}M \otimes u^*TN$.

1.7 **Lemma.** *If $u \in H(\mathbb{R}^+ \times M, N)$ is a weak solution of the heat flow and $u \in H_{loc}^2(\Omega_0, N)$ for some open subset $\Omega_0 \subseteq \mathbb{R}^+ \times M$, then u is stationary on Ω_0 .*

Proof (Feldman [7]). A weak solution of the heat flow in $H_{loc}^2(\Omega_0, N)$ is a strong solution of the heat equation (1.1) on Ω_0 .

Let Z be a vector field of compact support in Ω_0 . Then, denoting by \approx equality up to divergence terms we have

$$\begin{aligned} \frac{1}{2} \|du\|^2 \vec{\delta}Z &\approx \frac{1}{2} \vec{\nabla}_Z (\|du\|^2) = \langle \vec{\nabla}_Z du, du \rangle \\ &= \langle \nabla(\vec{d}u(Z)) - \vec{d}u(\vec{\nabla}Z), du \rangle = \langle \vec{d}u(Z), \nabla^* du \rangle - \langle \vec{d}u(\vec{\nabla}Z), du \rangle \\ &= -\langle \vec{d}u(Z), \partial_t u \rangle - \langle \vec{d}u(\vec{\Delta}Z), du \rangle, \end{aligned}$$

where in the last line we have used the heat equation (1.1). After integration we obtain the claim. \square

2.

2.1 Notation. For $z_1 = (t_1, x_1), z_2 = (t_2, x_2)$ in $\mathbb{R}^+ \times M$, the parabolic distance of z_1 and z_2 is defined as $\rho(z_1, z_2) = \max(|t_1 - t_2|^{1/2}, |x_1 - x_2|)$. We set $\rho(z, \Sigma) = \inf\{\rho(z, z') | z' \in \Sigma\}$. For $r > 0$, denote

$$\begin{aligned} \Sigma_r &= \{z \in \mathbb{R}^+ \times M | \rho(z, \Sigma) < r\}, \\ P_r(z) &= \{z' \in \mathbb{R} \times M | \rho(z, z') < r\}. \end{aligned}$$

Recall that m denotes the dimension of M .

2.2 Definition. The upper Minkowski q -content of a set $\Sigma \in \mathbb{R}^+ \times M$ with respect to the parabolic metric ρ is the number

$$\overline{\mathcal{M}}_\rho^q(\Sigma) = \limsup_{r \rightarrow 0} \frac{\mathcal{L}^{m+1}(\Sigma_r)}{r^{m+2-q}}.$$

We say that a set $\Sigma \subseteq \mathbb{R}^+ \times M$ has locally finite upper Minkowski k -content, i.e. for any compact set K we have $\overline{\mathcal{M}}_\rho^k(\Sigma \cap K) < \infty$.

2.3 Definition. We say that a set $\Sigma \subseteq \mathbb{R}^+ \times M$ has bounded uniform (m, q) -density with respect to the parabolic distance if for some $r_0 > 0$ the quantity

$$\Phi_{r_0}^{(m,q)}(\Sigma) = \sup\{\mathcal{L}^{m+1}(\Sigma_{r_1} \cap P_{r_2}(z)) / (r_1^{m+2-q} r_2^q) : r_1 < r_0, r_2 < r_0, z \in \mathbb{R}^+ \times M\}$$

is finite.

Clearly if $\Sigma \subseteq \mathbb{R}^+ \times M$ has bounded uniform (m, q) -density, then it has locally finite upper Minkowski q -content. Both these notions are not vacuous. To see this we give the following definition:

2.4 Definition. A closed set $\Sigma \in \mathbb{R}^+ \times M$ is *rectifiable of parabolic dimension q* if, for any compact set $K \subseteq \mathbb{R}^+ \times M$, there exist finitely many Lipschitzian maps $f_i: \mathbb{R}^{q-1} \rightarrow \mathbb{R}^+ \times M$ and $g_j: \mathbb{R}^q \rightarrow \{t_j\} \times M$ and compact sets $C_i \subseteq \mathbb{R}^{q-1}$ and $D_j \subseteq \mathbb{R}^q$ such that $\Sigma \cap K \subseteq \bigcup_{i,j} f_i(C_i) \cup g_j(D_j)$.

Then, it is not difficult to see that we have:

2.5 Proposition. *A rectifiable closed set $\Sigma \in \mathbb{R}^+ \times M$ of parabolic dimension q has locally finite upper Minkowski q -content. Similarly a compact smooth set $\Sigma \in \mathbb{R}^+ \times M$ of parabolic dimension q has bounded uniform (m, q) -density.*

3.

The goal of this section is to show that weak solutions of the heat flow are stationary if the singular set is “sufficiently small and regular” and the derivative du blows-up “slowly” near the singular set.

3.1 Theorem. *If $u \in H(\mathbb{R}^+ \times M, N) \cap C^\infty(\mathbb{R}^+ \times M \setminus \Sigma, N)$ is a weak solution of the heat flow, then u is a stationary solution provided that*

(a) *The singular set Σ has locally finite upper Minkowski p -content for some $p < m$.*

(b) *We have $du \in L^q(\mathbb{R}^+ \times M, TN)$ with $q = 2(m + 2 - p)/(m - p)$.*

We also reach the same conclusion if

- (a') The singular set Σ has null upper Minkowski $(m - 2)$ -content.
- (b') We have $|du(z)| \ll 1/\rho(z, \Sigma)$ for all $z \in \mathbb{R}^+ \times M \setminus \Sigma$.

We shall make use of the following observation, whose proof is standard.

3.2 Remark. For each compact set K and positive r , there exists a smooth cut-off function $\psi_r: \mathbb{R}^+ \times M \rightarrow [0, 1]$ with compact support such that $\psi_r = 0$ on $\Sigma_r \cap K$, $\psi_r = 1$ on $(\Sigma_{2r})^c \cap K$, $|\nabla\psi_r| < 2/r$ and $|\partial_t\psi_r| < 1/r^2$.

Proof of Theorem 3.1. Set $p_1 = (m + 2 - p)/(m - p)$ and $p_2 = (1 - 1/p_1)^{-1} = (m + 2 - p)/2$. From (a) we have $\limsup_{r \rightarrow 0} \mathcal{L}^{m+1}(\Sigma_r)/r^{m+2-p} < \infty$, which implies $\limsup_{r \rightarrow 0} \mathcal{L}^{m+1}(\Sigma_r)^{1/p_2}/r^2 < \infty$. Since the measure of Σ_r goes to zero with r , from (b) we have $\lim_{r \rightarrow 0} \int_{\Sigma_r} |du|^{2p_1} = 0$ and therefore, applying Hölder's inequality, we obtain

$$\begin{aligned}
 (3.1) \quad & \limsup_{r \rightarrow 0} \frac{1}{r^2} \int_{\Sigma_r} |du|^2 d\mathcal{L}^{m+1} \\
 & \leq \limsup_{r \rightarrow 0} \frac{1}{r^2} \left(\int_{\Sigma_r} |du|^{2p_1} d\mathcal{L}^{m+1} \right)^{1/p_1} (\mathcal{L}^{m+1}(\Sigma_r))^{1/p_2} = 0.
 \end{aligned}$$

If we assume (a') and (b') instead, we obtain

$$\int_{\Sigma_r \setminus \Sigma_{2r}} |du|^2 d\mathcal{L}^{m+1} \ll r^{-2} \mathcal{L}^{m+1}(\Sigma_{2r})$$

and $\lim_{r \rightarrow 0} \mathcal{L}^{m+1}(\Sigma_{2r})/r^4 = 0$. Thus under both sets of hypotheses we have that

$$(3.2) \quad \limsup_{r \rightarrow 0} \frac{1}{r^2} \int_{\Sigma_{2r} \setminus \Sigma_r} |du|^2 d\mathcal{L}^{m+1} = 0.$$

Since $\partial_t u \in L^2(\mathbb{R}^+ \times M, TN)$, equation (3.2) implies that

$$(3.3) \quad \limsup_{r \rightarrow 0} \frac{1}{r} \int_{\Sigma_{2r} \setminus \Sigma_r} |du| |\partial_t u| d\mathcal{L}^{m+1} = 0.$$

Now, let Z be a C^1 future-oriented vector field of compact support K on $\mathbb{R}^+ \times M$. Fix $r > 0$ and let $\psi_r: \mathbb{R}^+ \times M \rightarrow [0, 1]$ be a smooth cut-off function as in Remark 3.2. Then $\psi_r Z$ is also future-oriented and C^1 , and supported in a region on which u is smooth. Assuming that $K \subseteq [t_1, t_2] \times M$, by Lemma 1.7, we have

$$(3.4) \quad \int_{t_1}^{t_2} \int_M \left\{ \langle \partial_t u, \vec{d}u(\psi_r Z) \rangle + \langle du, \vec{d}u(\vec{\nabla}\psi_r Z) \rangle + \frac{1}{2} \langle du, du \rangle \vec{\delta}(\psi_r Z) \right\} \leq 0,$$

from which we derive

$$\begin{aligned}
 (3.5) \quad & \int_{t_1}^{t_2} \int_M \psi_r \left\{ \langle \partial_t u, \vec{d}u(Z) \rangle + \langle du, \vec{d}u(\vec{\nabla}Z) \rangle + \frac{1}{2} \langle du, du \rangle \vec{\delta}Z \right\} \\
 & \leq \int_{\Sigma_{2r} \setminus \Sigma_r} \left\{ \langle du, \vec{d}u(Z) \otimes d\psi_r \rangle - \frac{1}{2} \langle du, du \rangle \vec{d}\psi_r(Z) \right\}.
 \end{aligned}$$

As we let r tend to 0 the left-hand side of (3.5) converges to

$$\int_{t_1}^{t_2} \int_M \left\{ \langle \partial_t u, \vec{d}u(Z) \rangle + \langle du, \vec{d}u(\vec{\nabla}Z) \rangle + \frac{1}{2} \langle du, du \rangle \vec{\delta}Z \right\}.$$

Thus, in order to establish the proposition, it is sufficient to show that the right-hand side of (3.5) has a non-positive upper limit.

Writing $Z = T\partial_t + X$ with $dt(X) = 0$, the right-hand side of (3.5) is equal to

$$(3.6) \quad \int_{\Sigma_{2r} \setminus \Sigma_r} \left\{ T \langle du, \partial_t u \otimes d\psi_r \rangle + \langle du, du(X) \otimes d\psi_r \rangle - \frac{1}{2} \langle du, du \rangle (T\partial_t \psi_r + d\psi_r(X)) \right\},$$

and, using $|d\psi_r| < 2/r$ and $|\partial_t \psi_r| < 1/r^2$, we obtain that the expression (3.6) is dominated by

$$(3.7) \quad \int_{\Sigma_{2r} \setminus \Sigma_r} \left\{ \frac{2}{r} |T| |du| |\partial_t u| + \frac{2}{r} |X| |du|^2 + \frac{1}{2} \frac{1}{r^2} |T| |du|^2 + \frac{1}{2} \frac{2}{r} |X| |du|^2 \right\} d\mathcal{L}^{m+1} < 2 \|Z\|_\infty \int_{\Sigma_{2r} \setminus \Sigma_r} \left\{ \frac{1}{r^2} |du|^2 + \frac{1}{r} |du| |\partial_t u| \right\} d\mathcal{L}^{m+1},$$

which by (3.2)–(3.3) goes to zero as $r \rightarrow 0$. \square

4.

Henceforth we assume, for simplicity, that $M = \mathbb{R}^m$ with the ordinary Euclidean metric.

4.1 Lemma. *If $u \in H(\mathbb{R}^+ \times M, N)$ is a weak stationary solution of the heat flow, then*

(1) *The energy is decreasing at almost all times:*

$$(4.1) \quad E(u(t)) < E(u(0)) \quad \text{for a.e. } t \in \mathbb{R}^+.$$

(2) *There is an absolute constant C_1 such that:*

$$(4.2) \quad r^{2-m} \int_{P_{r/2}(z)} |\partial_t u|^2 < C_1 r^{-m} \int_{P_r(z)} |du|^2$$

for any ball $P_r(z) \subset \mathbb{R}^+ \times M$.

Proof. These are Propositions 10–12 in [7]. \square

4.2 Notation. For $u \in H(\mathbb{R}^+ \times M, N)$ and any ball $P_r(z) \subseteq \mathbb{R}^+ \times M$ we define the normalized energy in the r -ball about z as

$$(4.3) \quad N_r(u, z) = N(u, P_r(z)) = r^{-m} \int_{P_r(z)} |du|^2.$$

4.3 Lemma (Monotonicity property). *There is an absolute constant C_2 such that, for any weak stationary solution of the heat flow $u \in H(\mathbb{R}^+ \times M, N)$ and for any ball $P_r(z) \subseteq \mathbb{R}^+ \times M$, we have:*

$$(4.4) \quad N_{r'}(u, z') < C_2 N_r(u, z) \quad \text{for all } z' \in P_{r/4}(z) \text{ and } r' < r/8.$$

Proof. Let $\beta \in (0, 1)$ be a fixed constant and let $\phi: \mathbb{R}^m \rightarrow [0, 1]$ be a C^∞ function with support in $(B_{1/4}(0))$ and with $\phi \equiv 1$ on $B_{1/8}(0)$. For any $z_0 \in \mathbb{R}^+ \times M$, and any $R \in (0, \frac{\sqrt{t_0}}{2\beta})$, define

$$\Phi_\beta(R, u, z_0) = \frac{\beta^2 R^2}{2} \int_{\{t_0 - \beta^2 R^2\} \times M} |du|^2 G_{z_0} \phi_\beta^2 dm_g,$$

and

$$\Psi_\beta(R, u, z_0) = \frac{1}{2} \int_{t_0-4\beta^2 R^2}^{t_0-\beta^2 R^2} \int_M |du|^2 G_{z_0} \phi_\beta^2 d\mathcal{L}^{m+1},$$

where $G_{z_0} : (-\infty, t_0] \times M \rightarrow R^+$ is the solution of the equation $(\partial_t + \Delta)G_{z_0} = 0$ of the backward heat flow on M which satisfies the final condition $G_{z_0}(t_0) = \delta_{x_0}$, and where we have set $\phi_\beta(x) = \phi(\frac{x-x_0}{\beta})$.

We claim that there exists an absolute constant $C > 0$ such that for any $z_0 \in M \times R_+$, and $0 < R_1 \leq R_2 \leq \min(\frac{\sqrt{t_0}}{2\beta}, 1/4)$, we have

$$(4.5) \quad \Psi_\beta(R_1, u, z_0) \leq \Psi_\beta(R_2, u, z_0) + C\beta^{-m} \int_{P_{\beta/2}(z_0)} |du|^2 d\mathcal{L}^{m+1}.$$

Since the equation (1.1) is invariant under the translation $z \rightarrow z - z_0$, we may assume, to simplify the notation, that $z_0 = (0, 0)$. We denote $G = G_{(0,0)}$, $P_r = P_r(0, 0)$ and $B_r = B_r(0)$.

By the argument in [7] (see the derivation of (24) and (25) in [7]), it is not difficult to obtain the following estimates (take $\theta = tG\phi_\beta^2$, $\zeta = xG\phi_\beta^2$):

$$(4.6) \quad \begin{aligned} & \int_{\{t_2\} \times M} - \int_{\{t_1\} \times M} t|du|^2 G\phi_\beta^2 dm_g \\ & \geq \int_{t_1}^{t_2} \int_M \left[-|du|^2 \left(\frac{n-2}{2} + \frac{|x|^2}{4t} \right) \right. \\ & \quad \left. - \langle \partial_t u \otimes x, du \rangle - 2|\partial_t u|^2 t \right] G\phi_\beta^2 d\mathcal{L}^{m+1} + D_1 \end{aligned}$$

and

$$(4.7) \quad \int_{t_1}^{t_2} \int_M \left[\langle \partial_t u \otimes x, du \rangle - |du|^2 \left(\frac{n-2}{2} + \frac{|x|^2}{4t} \right) + \frac{|du \cdot x|^2}{2t} \right] G\phi_\beta^2 d\mathcal{L}^{m+1} + D_2 = 0,$$

where

$$\begin{aligned} D_1 &= \int_{t_1}^{t_2} \int_M \frac{4t}{\beta} \langle \partial_t u \otimes d\phi(x/\beta), du \rangle \phi_\beta G d\mathcal{L}^{m+1} \\ &= \int_{t_1}^{t_2} \int_M \frac{4t}{\beta} \partial_t u^\alpha \partial_j \phi(x/\beta) \partial_j u^\alpha \phi_\beta G d\mathcal{L}^{m+1} \end{aligned}$$

and

$$D_2 = \int_{t_1}^{t_2} \int_M \left[-\frac{1}{\beta} |du|^2 x^j \partial_j \phi(x/\beta) \phi_\beta G + \frac{2}{\beta} \partial_j u^\alpha \partial_k u^\alpha x^k \phi_\beta \partial_j \phi(x/\beta) G \right] d\mathcal{L}^{m+1}.$$

Let $t_1 = -R_2^2 \beta^2$, $t_2 = -R_1^2 \beta^2$; we have

$$\begin{aligned} |D_1| &\leq \int_{-R_2^2 \beta^2}^{-R_1^2 \beta^2} \int_M \beta^{2-m} |t|^{m/2} |\partial_t u|^2 G \phi_\beta^2 d\mathcal{L}^{m+1} \\ &\quad + C \int_{-R_2^2 \beta^2}^{-R_1^2 \beta^2} \int_M \beta^{m-4} |t|^{(4-m)/2} |du|^2 G |d\phi|^2 \left(\frac{x}{\beta} \right) d\mathcal{L}^{m+1} \\ &= D_{11} + D_{12}. \end{aligned}$$

By Lemma 4.1,

$$\begin{aligned}
 (4.8) \quad |D_{11}| &\leq C \int_{-R_2^2\beta^2}^{-R_1^2\beta^2} \int_{B_{\beta/4}} \beta^{2-m} |\partial_t u|^2 d\mathcal{L}^{m+1} \\
 &\leq C \int_{P_{\beta/4}} \beta^{2-m} |\partial_t u|^2 d\mathcal{L}^{m+1} \leq C\beta^{-m} \int_{P_{\beta/2}} |du|^2 d\mathcal{L}^{m+1}.
 \end{aligned}$$

Noticing that on $\{\beta/8 \leq |x| \leq \beta/4\} \times \{-R_2^2\beta^2 \leq t \leq -R_1^2\beta^2\}$,

$$|t|^{(4-m)/2} G \leq C|t|^{2-m} \exp\left\{-\frac{\beta^2}{256|t|}\right\} \leq C\beta^{4-2m},$$

we obtain

$$(4.9) \quad |D_{12}| \leq C\beta^{-m} \int_{P_{\beta/2}} |du|^2 d\mathcal{L}^{m+1}.$$

Formulas (4.8) and (4.9) lead to

$$(4.10) \quad |D_1| \leq C\beta^{-m} \int_{P_{\beta/2}} |du|^2 d\mathcal{L}^{m+1}.$$

Similarly, we obtain

$$(4.11) \quad |D_2| \leq C\beta^{-m} \int_{P_{\beta/2}} |du|^2 d\mathcal{L}^{m+1}.$$

Subtracting (4.7) from (4.6), and taking $t_1 = -R_2^2\beta^2$, $t_2 = -R_1^2\beta^2$, we obtain

$$\begin{aligned}
 (4.12) \quad &\Phi_\beta(R_2, u, 0) - \Phi_\beta(R_1, u, 0) \\
 &\geq \int_{-R_2^2\beta^2}^{-R_1^2\beta^2} \int_M - \left[2\partial_t u du \cdot x + \frac{|du \cdot x|^2}{2t} + 2|\partial_t u|^2 t \right] G \phi_\beta^2 d\mathcal{L}^{m+1} + D_1 - D_2.
 \end{aligned}$$

From (4.10)–(4.11) and the inequality

$$|2\langle \partial_t u \otimes x, du \rangle| \leq \frac{|\partial_j u x^j|^2}{2t} + 2|t| |\partial_t u|^2,$$

we see that (4.12) yields

$$(4.13) \quad \Phi_\beta(R_1, u, 0) \leq \Phi_\beta(R_2, u, 0) + C\beta^{-m} \int_{P_{\beta/2}} |du|^2 d\mathcal{L}^{m+1}.$$

Observing that

$$\Psi_\beta(R, u, 0) = \frac{2}{R} \int_R^{2R} \Phi_\beta(r, u, 0) dr,$$

we conclude from (4.13) that we have

$$\Psi_\beta(R_1, u, 0) \leq \Psi_\beta(R_2, u, 0) + C\beta^{-m} \int_{P_{\beta/2}} |du|^2 d\mathcal{L}^{m+1}.$$

Thus the claim (4.5) is proved.

As in the proof of Lemma 2.2 in [2], we have for any $0 < r' \leq r/8$

$$\begin{aligned} r'^{-m} \int_{P_{r'}(z_0)} |du|^2 d\mathcal{L}^{m+1} &\leq C\Psi_r(r'/r, u, (x_0, t_0 + 2r'^2)) \\ &\leq C\Psi_r(1/4, u, (x_0, t_0 + 2r'^2)) + Cr^{-m} \int_{P_{r/2}(x_0, t_0 + 2r'^2)} |du|^2 d\mathcal{L}^{m+1} \\ &\leq Cr^{-m} \int_{P_{r/2}(x_0, t_0 + 2r'^2)} |du|^2 d\mathcal{L}^{m+1} \leq Cr^{-m} \int_{P_r(z_0)} |du|^2 d\mathcal{L}^{m+1}. \end{aligned}$$

□

To prove Theorem 1.2 we will require two lemmata. The first lemma states that in a region where u is smooth we can estimate $|du|^2$ by its average.

4.4 Lemma. *Let $u \in H(\mathbb{R}^+ \times M, N)$ be a weak stationary solution of the heat flow. There are absolute constants ε_1 and C_3 such that if u is smooth on $P_r(z_0) \subseteq \mathbb{R}^+ \times M$ and $N_r(u, z_0) < \varepsilon_1$, then*

$$(4.14) \quad \sup_{P_{r/16}(z_0)} |du|^2 < C_3 r^{-m-2} \int_{P_r(z)} |du|^2.$$

Proof. We follow Schoen and Uhlenbeck [12, Theorem 2.2]. Bochner’s formula for $\Delta|du|^2$ (see [5]) implies the estimate

$$(\partial_t - \Delta)|du|^2 \leq C_N |du|^4,$$

where C_N is the norm of the Riemannian curvature of (N, h) .

Let $z' \in P_r(z_0)$ and $P_\alpha(z') \subseteq P_r(z_0)$. If on $P_\alpha(z')$ we have $|du|^2 < B$, then

$$(\partial_t - \Delta)|du|^2 \leq C_N B |du|^2, \quad \text{on } P_\alpha(z'),$$

and, applying Moser’s mean value inequality [10, Formula (3.12)] for parabolic equations in the rectangle $P_\alpha(z')$, we obtain

$$(4.15) \quad |du|^2(z') < \gamma e^{C_N B \alpha^2} \alpha^{-2} N_\alpha(u, z'),$$

where the constant γ depends only on m .

Choose $\tau_0 \in (0, r/8)$ so that

$$(4.16) \quad (r/8 - \tau_0)^2 \sup_{P_{\tau_0}(z_0)} |du|^2 = \max_{0 \leq \tau \leq r/8} \left[(r/8 - \tau)^2 \sup_{P_\tau(z_0)} |du|^2 \right]$$

and pick $\bar{z} \in P_{\tau_0}(z_0)$ so that

$$(4.17) \quad |du|^2(\bar{z}) = \max_{P_{\tau_0}(z_0)} |du|^2 = e_0.$$

Set $\rho_0 = \frac{1}{2}(r/8 - \tau_0)$. Then $\tau_0 + \rho_0 \leq r/8$ and therefore

$$(4.18) \quad \max_{P_{\rho_0}(\bar{z})} |du|^2 \leq \max_{P_{\rho_0 + \tau_0}(z)} |du|^2 \leq \frac{(r/8 - \tau_0)^2}{(r/8 - \rho_0 - \tau_0)^2} |du|^2(\bar{z}) = 4e_0.$$

If $\rho_0 > (e_0 C_N)^{-1/2}$, then applying (4.15) with $z' = \bar{z}$ and $\alpha = (e_0 C_N)^{-1/2}$ we obtain

$$e_0 = |du|^2(\bar{z}) < \gamma e^4 e_0 C_N N_{(e_0 C_N)^{-1/2}}(u, \bar{z}) < \gamma e^4 e_0 C_N C_1 N_r(u, z_0).$$

In the last step, we have used the monotonicity inequality and the fact that $(e_0 C_N)^{-1/2} < \rho_0 < r/16$ and $\bar{z} \in P_{r/8}(z_0)$. The last inequality is a contradiction

if $N_r(u, z_0)$ is smaller than $\varepsilon_1 = \gamma^{-1}e^{-4}C_N^{-1}$. Thus we can assume that $\rho_0 < (e_0C_N)^{-1/2}$. Applying (4.15) with $z' = \bar{z}$ and $\alpha = \rho_0$ we obtain

$$e_0 = |du|^2(\bar{z}) < \gamma e^4 \rho^{-2} N_{\rho_0}(u, \bar{z}) < \gamma e^4 \rho^{-2} C_1 N_r(u, z_0)$$

and therefore

$$(r/8 - \tau)^2 \sup_{P_\tau(z_0)} |du|^2 \leq 4\rho^2 e_0 \leq \gamma e^4 C_1 N_r(u, z_0).$$

Choosing $\tau = r/16$ and $C_3 = 64\gamma e^4 C_1$ we conclude that:

$$\sup_{P_\tau(z_0)} |du|^2 \leq C_3 r^{-m-2} \int_{P_\tau(z)} |du|^2.$$

□

4.5 Lemma. *Assume that $u \in H(\mathbb{R}^+ \times M, N)$ is a weak stationary solution of the heat flow with singular set $\Sigma \subseteq \mathbb{R}^+ \times M$ satisfying the condition (2) of Theorem 1.2. There exist constants $\sigma = \sigma(m, N, \delta_0) \in (0, 1/4)$ and $\varepsilon_1 = \varepsilon_1(m, N, \delta_0)$ such that for any $z \in M \times [\delta_0^2, \infty)$ and $r < \min(1/4, \delta_0/4)$ the inequality $N_r(u, z) < \varepsilon_1$ implies*

$$(4.19) \quad N_{\sigma r}(u, z) < \frac{1}{2} N_r(u, z).$$

Proof. Arguing by contradiction, we fix sequences ε_i and $\sigma_i < 1/4$ decreasing to zero and we assume that there exist sequences of points $z_i \in [\delta_0^2, \infty) \times M$ and $r_i < \min(1/4, \delta_0/4)$ such that

$$(4.20) \quad \lambda_i^2 := \frac{1}{2} N_{r_i}(u, z_i) < \varepsilon_i \quad \text{for all } i$$

and

$$(4.21) \quad N_{\sigma_i r_i}(u, z_i) \geq \frac{1}{2} N_{r_i}(u, z_i) \quad \text{for all } i.$$

Define

$$a_i = \frac{1}{\text{Vol}(P_{r/2}(z_i))} \int_{P_{r/2}(z_i)} u$$

and set

$$(4.22) \quad v_i(t, x) = \lambda_i^{-1} u(r_i^2 t + t_i, r_i x + x_i).$$

Then $v_i: P_1 \subseteq \mathbb{R} \times M \rightarrow N$ is a stationary solution of the heat flow satisfying

$$(4.23) \quad N_1(v_i, 0) = \lambda_i^{-2} N_{r_i}(u, z_i) = 1$$

and

$$(4.24) \quad N_{\sigma_i}(v_i, 0) = \lambda_i^{-2} N_{\sigma_i r_i}(u, z_i) \geq \frac{\lambda_i^{-2}}{2} N_{r_i}(u, z_i) = \frac{1}{2}.$$

From (4.2) and (4.23) we obtain

$$\int_{P_{1/2}} |\partial_t v_i|^2 < C_1$$

and therefore, using the Poincaré inequality,

$$\int_{P_{1/2}} |v_i|^2 < C_2.$$

We can assume then, by possibly passing to a subsequence, that as $i \rightarrow \infty$ we have

$$(4.25) \quad v_i \rightarrow \bar{v} \text{ strongly in } L^2(P_{1/2}, \mathbb{R}^K),$$

$$(4.26) \quad dv_i \rightarrow d\bar{v} \text{ weakly in } L^2(P_1, \mathbb{R}^K),$$

$$(4.27) \quad \partial_t v_i \rightarrow \partial_t \bar{v} \text{ weakly in } L^2(P_{1/2}, \mathbb{R}^K).$$

Choose any smooth map $\omega: \mathbb{R} \times M \rightarrow \mathbb{R}^K$ with support in $P_{1/2}$. Then since v_i are weak solutions of the heat equation (1.1) we have

$$\int_{\mathbb{R} \times M} \langle \partial_t v_i, \omega \rangle + \langle dv_i, d\omega \rangle = \lambda_i \int_{\mathbb{R} \times M} A(a_i + \lambda_i v_i)(dv_i, dv_i)\omega.$$

By the compactness of N the right-hand side is bounded by $C_3 \lambda_i N_{1/2}(v_i) \|\omega\|_\infty$, and therefore as $i \rightarrow \infty$ we obtain from (4.26), (4.27)

$$\int_{\mathbb{R} \times M} \langle \partial_t \bar{v}, \omega \rangle + \langle d\bar{v}, d\omega \rangle = 0.$$

Hence the limit \bar{v} solves the heat equation $(\partial_t + \Delta)\bar{v} = 0$ weakly and also classically. From the local estimate of the parabolic equations and (4.23), (4.26) we obtain that

$$\|d\bar{v}\|_{L^\infty(P_{1/4})} \leq C_4 N_{1/2}(\bar{v}).$$

□

4.6 *Claim.*

$$dv_i \rightarrow d\bar{v} \text{ strongly in } L^2(P_{1/4}, \mathbb{R}^K).$$

Fix σ so that $N_\sigma(\bar{v}) < 1/4$. From Claim 4.6 and (4.24) we obtain

$$\frac{1}{4} > N_\sigma(\bar{v}) = \lim_i N_\sigma(v_i) \geq \lim_i N_{\sigma_i}(v_i) \geq \frac{1}{2},$$

which is a contradiction. Thus in order to conclude our proof, it suffices to prove Claim 4.6.

Proof of Claim 4.6. Recall that $\Sigma(u)$ denotes the singular set of u . For simplicity we define for $z = (t, x) \in P_1$

$$\phi_i(t, x) = (r_i^2 t + t_i, r_i x + x_i).$$

Thus $v_i = \lambda_i^{-1} u \circ \phi_i$ on P_1 .

Given an integer $\nu = 4^l$ with $l > 3$, let \mathcal{P}_ν be the partition P_1 into the parallelepipeds with vertices on the lattice $\{(2\nu^{-2}j_0, 2\nu^{-1}j_1, \dots, 2\nu^{-1}j_m) : (j_0, j_1, \dots, j_m) \in \mathbb{Z}^m\}$. We say that a parallelepiped $P \in \mathcal{P}_\nu$ is ν_i -good if the function v_i is smooth on P and on all parallelepipeds contiguous to P .

By passing to a subsequence of v_i , still denoted by v_i , we have that each P is either ν_i -good for all i or ν_i -bad for all i . Thus we can simply say that $P \in \mathcal{P}$ is good or bad.

Denote by \mathcal{G}_ν the collection of good parallelepipeds meeting $P_{1/4}$ and let $G_\nu = \bigcup_{P \in \mathcal{G}_\nu} P$ be their union. Let us also denote the collection of bad parallelepipeds meeting $P_{1/4}$ by \mathcal{B}_ν and their union by B_ν .

Let $P \in \mathcal{G}_\nu$. Clearly any point in P has parabolic distance greater than ν^{-1} from the singular set $\Sigma_i = \phi_i^{-1}(\Sigma(u))$ of v_i . This implies that $\phi_i(P)$ has parabolic distance greater than $r_i \nu^{-1}$ from $\Sigma(u)$. Therefore we obtain that for $z \in P$,

$$(4.28) \quad |dv_i|^2(z) = \lambda_i^{-2} r_i^2 |du|^2(\phi_i(z)).$$

By the monotonicity formula (4.4), we have

$$N_{16r_i\nu^{-1}}(u, \phi_i(z)) < C_2 N_{r_i/8}(u, \phi_i(z)) \leq C_2 N_{r_i}(u, z_i) = C_2 \lambda_i^2,$$

and therefore for large i we have $N_{16r_i\nu^{-1}}(u, \phi_i(z)) < \varepsilon_0$, where ε_0 is the constant appearing in Lemma 4.4. Applying now this lemma we obtain that

$$(4.29) \quad \begin{aligned} |du|^2(\phi_i(z)) &< C_3(r_i\nu^{-1})^{-2} N_{r_i\nu^{-1}}(u, \phi_i(z)) \\ &< C_2 C_3(r_i\nu^{-1})^{-2} N_{16r_i\nu^{-1}}(u, \phi_i(z)) \\ &< C_2 C_3(r_i\nu^{-1})^{-2} \lambda_i^2. \end{aligned}$$

Therefore (4.28) and (4.29) imply that for $z \in P$ and $P \in \mathcal{G}_\nu$

$$|dv_i|^2(z) < C_2 C_3 \nu^2$$

and by passing to a subsequence we obtain that dv_i converges strongly in L^2 on G_ν .

Now we estimate $\int_{B_\nu} |dv_i|^2$. Observe that if $z \in P_{1/4}$ belongs to a bad parallelepiped P , then there is a point of Σ_i in P or in a parallelepiped contiguous to P . Then we have $\rho(z, \Sigma_i) < 2\nu^{-1}$ and therefore $\rho(\phi_i(z), \Sigma(u)) < 2r_i\nu^{-1}$. Then

$$\mathcal{L}^{m+1}(B_\nu) < \mathcal{L}^{m+1}(P_{1/4} \cap (\Sigma_i)_{2\nu^{-1}}) < r_i^{-m-2} \mathcal{L}^{m+1}(P_{r_i/4}(z_i) \cap \Sigma(u)_{2r_i\nu^{-1}}).$$

By our assumption on $\Sigma(u)$ we have that

$$\mathcal{L}^{m+1}(P_{r_i/4}(z_i) \cap \Sigma(u)_{2r_i\nu^{-1}}) < C(r_i/4)^{m+2-q} (2r_i\nu^{-1})^q = C' r_i^{m+2} \nu^{-q}$$

and we conclude that

$$\mathcal{L}^{m+1}(B_\nu) < C' \nu^{-q}.$$

Since each parallelepiped in \mathcal{P} has \mathcal{L} -measure equal to $2^{m+1} \nu^{-m-2}$, we obtain that \mathcal{B}_ν has at most $C' \nu^{m+2-q}$ elements. Thus, denoting by $c(P, i)$ the centre of $\phi_i(P)$ we have

$$\begin{aligned} \int_{B_\nu} |dv_i|^2 &= \sum_{P \in \mathcal{B}_\nu} \int_P |dv_i|^2 = \sum_{P \in \mathcal{B}_\nu} (\mathcal{L}^{m+1}(P))^{m/(m+2)} N(v_i, P) \\ &= 2^{m(m+1)/(m+2)} \nu^{-m} \sum_{P \in \mathcal{B}_\nu} N(v_i, P) \\ &= 2^{m(m+1)/(m+2)} \nu^{-m} \sum_{P \in \mathcal{B}_\nu} \lambda_i^{-2} N(u, \phi_i(P)) \\ &= 2^{m(m+1)/(m+2)} \nu^{-m} \sum_{P \in \mathcal{B}_\nu} \lambda_i^{-2} N_{r_i\nu^{-1}}(u, c(P, i)) \\ &< 2^m \nu^{-m} \sum_{P \in \mathcal{B}_\nu} C 2^m \lambda_i^{-2} N_{r_i}(u, z_i) \\ &= C 2^{2m} \nu^{-m} \sum_{P \in \mathcal{B}_\nu} 1 \\ &< C 2^{2m} \nu^{-m} C' \nu^{m+2-q} = 2^{2m} C C' \nu^{2-q} = C''' \nu^{2-q}. \end{aligned}$$

We conclude that

$$\limsup_i \int_{P_{1/4}} |dv_i|^2 \leq \int_{G_\nu} |d\bar{v}|^2 + C''' \nu^{2-q}.$$

Since $q > 2$, letting $\nu \rightarrow \infty$ we obtain that

$$\limsup_i \int_{P_{1/4}} |dv_i|^2 \leq \int_{G_\nu} |d\bar{v}|^2.$$

This implies our claim. \square

Proof of Theorem 1.2. Since u is a weak stationary solution of the heat equation (1.1), from the monotonicity inequality (4.4) and the energy inequality (4.1) we have

$$N_r(u, z) < C_2 N_{r_0}(u, z) < C_2 r_0^{2-m} E(u_0)$$

for any $z = (x, t) \in M \times [\delta_0^2, \infty)$ and any $r < r_0/8 = \delta_0/8$.

Let ε_1 be determined as in Lemma 4.5. Then, if $E(u_0)$ is smaller than $\varepsilon_1 C_2^{-1} r_0^{m-2}$, we obtain that $N_r(u, z) < \varepsilon_1$ for any $z = (x, t) \in M \times [\delta_0^2, \infty)$ and any $r < \delta_0/8$. Using Lemma 4.5 and a standard iteration argument (see [2, Lemma 5.1], for example), we obtain the Morrey type estimate

$$\int_{P_r(z)} (|du|^2 + r^2 |\partial_t u|^2) \ll r^{m+\alpha}$$

which implies that u is $C^{\alpha/2}$ in some ball around z . The fact that u is C^∞ follows as in [7, Lemma 21]. \square

REFERENCES

1. F. Bethuel *On the singular set of stationary harmonic maps*, Manuscripta Math. **78** (1993), 417–443. MR **94a**:58047
2. Y. Chen, J. Li, and F. H. Lin *Partial regularity for weak heat flows into spheres*, Comm. Pure Appl. Math. (to appear).
3. Y. Chen and M. Struwe *Existence and partial regularity for heat flow for harmonic maps*, Math. Z. **201** (1989), 83–103. MR **90i**:58031
4. J. M. Coron *Nonuniqueness for the heat flow of harmonic maps*, Ann. Inst. H. Poincaré Anal. Non Linéaire **7** (1990), 335–344. MR **91g**:58058
5. J. Eells and J. Sampson *Harmonic mappings of Riemannian manifolds*, Amer. J. Math. **86** (1964), 133–147. MR **29**:1603
6. L. C. Evans *Partial regularity for stationary harmonic maps into spheres*, Arch. Rational Mech. Anal. **116** (1991), 101–113. MR **93m**:58026
7. M. Feldman *Partial regularity for harmonic maps of evolution into spheres*, preprint (1994).
8. F. Hélein *Régularité des applications faiblement harmoniques entre une surface et une sphere*, C. R. Acad. Sci. Paris. Sér I Math. **311** (1990), 519–524. MR **92a**:58034
9. ———, *Régularité des applications faiblement harmoniques entre une surface et une variété riemannienne*, C. R. Acad. Sci. Paris Sér. I Math. **312** (1991), 591–596. MR **92e**:58055
10. J. Moser *A Harnack inequality for parabolic differential equations*, Comm. Pure Appl. Math. **17** (1964), 101–134. MR **28**:2357
11. L. Mou *Removability of singular sets of harmonic maps*, preprint, 1993.
12. R. Schoen *Analytical aspects of the harmonic map problem*, Seminar on Nonlinear Partial Differential Equations (S. S. Chern, ed.), vol. 2, Math. Sci. Res. Inst. Publ., Springer-Verlag, New York, Berlin, Heidelberg, and Tokyo, 1984, pp. 321–358. MR **86b**:58032
13. L. Simon *Singularities of geometric variational problems*, Graduate Summer School Lecture at Stanford, 1993.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF FLORIDA, GAINESVILLE, FLORIDA 32611

E-mail address: `yunmei@math.ufl.edu`

E-mail address: `flaminio@math.ufl.edu`