

## $E_\infty$ -RING STRUCTURES FOR TATE SPECTRA

J. E. McCLURE

(Communicated by Thomas Goodwillie)

### 1. INTRODUCTION

Let  $G$  be a compact Lie group and  $k_G$  a  $G$  spectrum (as defined in [3, Section I.2]). Greenlees and May ([2]) have defined an associated  $G$ -spectrum  $t(k_G)$  called the *Tate spectrum* of  $k_G$ . They observe that if  $k_G$  is a ring  $G$ -spectrum, then there is an induced ring  $G$ -spectrum structure on  $t(k_G)$ , and that if  $k_G$  is homotopy-commutative, then  $t(k_G)$  will also be homotopy-commutative (see [2, Proposition 3.5]). It is therefore natural to ask whether an equivariant  $E_\infty$  ring structure on  $k_G$  induces an equivariant  $E_\infty$  ring structure on  $t(k_G)$  (we will recall the definition in a moment). We offer both positive and negative answers to this question.

On the positive side, we show that  $t(k_G)$  inherits a structure which is somewhat weaker than an equivariant  $E_\infty$  ring structure, but which should be adequate for most practical purposes. To explain this we first recall that, given a  $G$ -universe  $U$ , there is an equivariant operad  $\mathcal{L}(U)$  whose  $j$ th space consists of the (nonequivariant) linear isometries from  $U^{\oplus j}$  to  $U$ . From now on we fix a complete  $G$ -universe  $U$ . By definition, an *equivariant  $E_\infty$  operad*  $\mathcal{C}$  is an equivariant operad which is equivariantly equivalent to  $\mathcal{L}(U)$ , and an *equivariant  $E_\infty$  ring structure* is an action of an equivariant  $E_\infty$  operad (see [3, Definition VII.2.1]). Next let us define an  *$E'_\infty$  operad* to be a nonequivariant  $E_\infty$  operad provided with trivial  $G$ -action. For example, if  $\mathcal{C}$  is an equivariant  $E_\infty$  operad, then its  $G$ -fixed points form an  $E'_\infty$  operad  $\mathcal{C}^G$  by [3, Example VII.1.4]. We define an *equivariant  $E'_\infty$  ring structure* to be an action of an  $E'_\infty$  operad (in other words, it is an action of a nonequivariant  $E_\infty$  operad through  $G$ -maps). Since any action by  $\mathcal{C}$  restricts to an action by  $\mathcal{C}^G$ , we see that an equivariant  $E_\infty$  ring structure always includes an equivariant  $E'_\infty$  ring structure.

The reason why  $E'_\infty$  ring structures are interesting is that if  $k_G$  is an  $E'_\infty$  ring spectrum, then the fixed-point spectra  $(k_G)^H$  have (nonequivariant)  $E_\infty$  ring structures which are consistent as  $H$  varies (see Remark VII.2.5 of [3]); this is likely to be the property one needs for applications.

Our positive result is:

**Theorem 1.** *If  $k_G$  is an equivariant  $E'_\infty$  ring spectrum, then so is  $t(k_G)$ ; in particular all fixed-point spectra  $(t(k_G))^H$  are nonequivariant  $E_\infty$  ring spectra.*

---

Received by the editors June 24, 1994 and, in revised form, November 10, 1994.

1991 *Mathematics Subject Classification.* Primary 55P91.

The author was partially supported by National Science Foundation grant 9207731-DMS.

The proof of Theorem 1 will show that the diagram in Proposition 3.5 of [2] is a diagram of equivariant  $E'_\infty$  ring spectra.

To state our negative result we need to recall the definition of  $t(k_G)$ . Let  $EG$  be a contractible free  $G$ -CW complex and let  $\tilde{E}G$  denote the  $G$ -space defined by the cofiber sequence

$$EG_+ \rightarrow S^0 \rightarrow \tilde{E}G$$

(here  $+$  denotes a disjoint basepoint). Let  $F(EG_+, k_G)$  be the function spectrum of maps from  $EG_+$  to  $k_G$  ([3, Definition I.3.2]). Then  $t(k_G)$  is defined to be the  $G$ -spectrum

$$F(EG_+, k_G) \wedge \tilde{E}G.$$

Let us write  $\iota$  for the natural map  $S^0 \rightarrow \tilde{E}G$ .

**Theorem 2.** *Let  $G$  be a finite group and let  $k_G$  be any  $G$ -spectrum. Suppose that  $t(k_G)$  has an equivariant  $E_\infty$  ring structure whose unit factors (up to equivariant homotopy) through  $\sum_G^\infty \iota$ . Then  $t(k_G)$  must be equivariantly contractible.*

This implies that if  $k_G$  is a ring  $G$ -spectrum for which  $t(k_G)$  is not equivariantly contractible, then  $t(k_G)$  cannot have an equivariant  $E_\infty$  ring structure whose underlying ring  $G$ -spectrum structure is compatible with that of  $k_G$  under the natural map  $k_G \rightarrow t(k_G)$ . In particular, the underlying ring  $G$ -spectrum structure of  $t(k_G)$  cannot be that defined in [3, Proposition 3.5]. Thus it seems that there is no natural way to give  $t(k_G)$  an equivariant  $E_\infty$  ring structure.

I would like to thank Mike Hopkins for suggesting this problem to me and the referee for clarifying a point in the proof.

## 2. PROOF OF THEOREM 1

Theorem 1 is an immediate consequence of the following two lemmas, of which the second is well-known. Let us recall from [3, Definition VII.2.7] that, given an equivariant operad  $\mathcal{C}$ , a  $\mathcal{C}_0$  space is an action of  $\mathcal{C}$  in the category of based  $G$ -spaces; that is, it is a based  $G$ -space  $X$  with based  $G$ -maps

$$(\mathcal{C}_j)_+ \wedge_{\Sigma_j} X^{(j)} \rightarrow X$$

(here  $^{(j)}$  denotes  $j$ -fold smash product) satisfying the same compatibility conditions that are used to define an equivariant  $\mathcal{C}$ -space. In particular, this definition makes sense if  $\mathcal{C}$  is a nonequivariant operad provided with the trivial  $G$ -action; it then says that  $\mathcal{C}$  acts on  $X$  through  $G$ -maps.

**Lemma 3.** *There is a nonequivariant  $E_\infty$  operad  $\mathcal{D}$  for which  $\tilde{E}G$  is an equivariant  $\mathcal{D}_0$  space.*

**Lemma 4.** *Let  $\mathcal{C}$  be any equivariant operad.*

(a) *If  $k_G$  is a  $\mathcal{C}$  ring spectrum (that is, if it has an equivariant action of  $\mathcal{C}$ ), then so is  $F(Y_+, k_G)$  for any  $G$ -space  $Y$ .*

(b) *If  $h_G$  is a  $\mathcal{C}$  ring spectrum and  $X$  is a  $\mathcal{C}_0$ -space, then  $h_G \wedge X$  is a  $\mathcal{C}$  ring spectrum.*

*Proof of Theorem 1.* Suppose that  $k_G$  has an action of an  $E'_\infty$  operad  $\mathcal{C}'$ . Let  $\mathcal{C} = \mathcal{C}' \times \mathcal{D}$ , where  $\mathcal{D}$  is the operad of Lemma 3. Then  $\mathcal{C}$  is an  $E'_\infty$  operad, and it acts on  $k_G$  (via the projection  $\mathcal{C}' \times \mathcal{D} \rightarrow \mathcal{C}'$ ) and on  $\tilde{E}G$  (via the projection

$\mathcal{C}' \times \mathcal{D} \rightarrow \mathcal{D}$ ). Now Lemma 4(a) implies that  $\mathcal{C}$  acts on  $F(EG_+, k_G)$ , and the theorem follows from Lemma 4(b) if we take  $h_G$  to be  $F(EG_+, k_G)$  and  $X$  to be  $\widetilde{EG}$ .  $\square$

*Proof of Lemma 4.* In each case, we specify the structural maps which constitute the  $\mathcal{C}$ -action; the fact that they satisfy the necessary compatibility relations is a straightforward application of the methods of [3, Sections VI.1–VI.3].

For part (a) the structural map

$$\xi_j : \mathcal{C}_j \times F(Y_+, k_G)^{(j)} \rightarrow F(Y_+, k_G)$$

is the adjoint of the composite

$$\begin{aligned} Y_+ \wedge \mathcal{C}_j \times F(Y_+, k_G)^{(j)} &\xrightarrow{\Delta \wedge 1} (Y_+)^{(j)} \wedge \mathcal{C}_j \times F(Y_+, k_G)^{(j)} \\ &\xrightarrow{\cong} \mathcal{C}_j \times \left( (Y_+)^{(j)} \wedge F(Y_+, k_G)^{(j)} \right) \xrightarrow{1 \times e} \mathcal{C}_j \times k_G^{(j)} \xrightarrow{\xi'_j} k_G; \end{aligned}$$

here  $\Delta$  is the diagonal map of  $Y$ , the isomorphism is that of [3, Proposition VI.1.5],  $e$  is the evaluation map, and  $\xi'_j$  is the structural map of  $k_G$ .

For part (b) the structural map

$$\xi_j : \mathcal{C}_j \times (h_G \wedge X)^{(j)} \rightarrow h_G \wedge X$$

is the composite

$$\mathcal{C}_j \times (h_G \wedge X)^{(j)} = \mathcal{C}_j \times (h_G^{(j)} \wedge X^{(j)}) \xrightarrow{\delta} (\mathcal{C}_j \times h_G^{(j)}) \wedge (\mathcal{C}_{j+} \wedge X^{(j)}) \xrightarrow{\xi'_j \wedge \xi''_j} h_G \wedge X,$$

where  $\delta$  is the map given in Definition VI.3.5 of [3] and  $\xi'_j, \xi''_j$  are the structural maps for  $h_G$  and  $X$ .  $\square$

*Proof of Lemma 3.* First let us observe that  $\widetilde{EG}$  is nonequivariantly contractible and that for any nontrivial subgroup  $H$  of  $G$  the  $H$ -fixed set  $(\widetilde{EG})^H$  is exactly  $S^0$ ; the same is true for  $(\widetilde{EG})^{(j)}$  since the smash product of spaces commutes with  $H$ -fixed sets.

Let  $\text{Map}_*^G$  denote based  $G$ -maps. Restriction to the  $G$ -fixed set gives a map

$$\phi : \text{Map}_*^G(\widetilde{EG}^{(j)}, \widetilde{EG}) \rightarrow \text{Map}_*(S^0, S^0)$$

which we claim is a weak equivalence. Assuming this for the moment, let  $\mathcal{D}'_j$  be the space  $\phi^{-1}(\text{id})$ . Then the spaces  $\mathcal{D}'_j$  with the evident composition operations  $\gamma$  form an operad  $\mathcal{D}'$  and  $\widetilde{EG}$  is a  $\mathcal{D}'_0$ -space. The only thing preventing  $\mathcal{D}'$  from being a nonequivariant  $E_\infty$  operad is that the action of  $\Sigma_j$  on  $\mathcal{D}'_j$  may not be free. To remedy this let  $\mathcal{C}$  be any nonequivariant  $E_\infty$  operad and define  $\mathcal{D}$  to be  $\mathcal{D}' \times \mathcal{C}$ , acting on  $\widetilde{EG}$  via the projection  $\mathcal{D}' \times \mathcal{C} \rightarrow \mathcal{D}'$ .

It only remains to prove the claim that  $\phi$  is a weak equivalence. First we observe that the reduced diagonal map

$$\Delta : \widetilde{EG} \rightarrow \widetilde{EG}^{(j)}$$

is a weak equivalence on each fixed-point set, and is therefore a  $G$ -homotopy equivalence by the equivariant Whitehead theorem. It follows that

$$\Delta^* : \text{Map}_*^G(\widetilde{EG}, \widetilde{EG}) \rightarrow \text{Map}_*^G(\widetilde{EG}^{(j)}, \widetilde{EG})$$

is a homotopy equivalence, so it suffices to verify the claim when  $j = 1$ .

To handle this case, we map the cofiber sequence

$$EG_+ \rightarrow S^0 \rightarrow \tilde{E}G$$

into  $\tilde{E}G$  to get a fiber sequence

$$\text{Map}_*^G(\tilde{E}G, \tilde{E}G) \rightarrow \text{Map}_*^G(S^0, \tilde{E}G) \rightarrow \text{Map}_*^G(EG_+, \tilde{E}G).$$

The middle term is equal to  $S^0$ , so it suffices to show that the third term is weakly contractible. For this we recall that the functor  $\text{Map}_*^G(EG_+, -)$  takes  $G$ -maps which are nonequivariant weak equivalences to weak equivalences (for example, this follows from [1, XI.5.6] since  $\text{Map}_*^G(EG_+, -)$  is a special case of the holim construction). Since  $\tilde{E}G$  is nonequivariantly contractible, we see that  $\text{Map}_*^G(EG_+, \tilde{E}G)$  is weakly contractible and we are done.  $\square$

### 3. PROOF OF THEOREM 2

As motivation for the proof of Theorem 2, we first explain why the operad  $\mathcal{D}'$  constructed in the proof of Lemma 3 is not equivalent to the linear isometries operad  $\mathcal{L}U$ . Let  $G = Z/2$  for simplicity and consider the  $G \times \Sigma_2$ -spaces  $\mathcal{L}U_2$  and  $\mathcal{D}'_2$  (recall that  $\mathcal{D}'_2$  has trivial  $G$ -action). Let  $H$  be the diagonal copy of  $Z/2$  in  $G \times \Sigma_2 = Z/2 \times Z/2$ . We claim that  $\mathcal{L}U_2$  has  $H$  fixed points but  $\mathcal{D}'_2$  has none; this certainly implies that  $\mathcal{L}U_2$  and  $\mathcal{D}'_2$  are not  $G \times \Sigma_2$ -equivalent. To see that  $\mathcal{L}U_2$  has  $H$ -fixed points we need only show that there is an  $H$ -equivariant linear isometry from  $U \oplus U$  to  $U$ ; but this is obvious since as an  $H$ -representation  $U \oplus U$  is a complete  $H$ -universe, and is therefore  $H$ -isomorphic to  $U$ . (We note for later use that  $(\mathcal{L}U_2)^H$  is in fact contractible by [3, Lemma II.1.5].) On the other hand, if  $\mathcal{D}'_2$  had an  $H$ -fixed point, then there would be a  $G \times \Sigma_2$ -equivariant map

$$\tilde{E}G^{(2)} \rightarrow \tilde{E}G$$

(with  $\Sigma_2$  acting trivially on the target) which extends the identity map of  $S^0$ , and passing to  $H$ -fixed points would give a (nonequivariant) map  $(\tilde{E}G^{(2)})^H \rightarrow S^0$  which extends the identity map of  $S^0$ . But this is impossible since  $(\tilde{E}G^{(2)})^H$  is contractible: there is a (nonequivariant) homeomorphism

$$\tilde{E}G \rightarrow (\tilde{E}G^{(2)})^H$$

which takes  $x$  to  $x \wedge gx$ , where  $g$  is the generator of  $G$ .

The proof of Theorem 2 is a variant of the same idea. For simplicity, we begin with the case  $G = Z/2$ . Suppose that  $t(k_G)$  has an equivariant  $E_\infty$  ring structure whose unit  $\eta$  factors through  $\Sigma_G^\infty \iota$ . Then there is a  $G$ -homotopy commutative diagram of  $G$ -spectra

$$\begin{array}{ccc} \mathcal{L}U_2 \times_{\Sigma_2} (S_G^0)^{(2)} & \xrightarrow{1 \times \Sigma_G^\infty \iota^{(2)}} & \mathcal{L}U_2 \times_{\Sigma_2} \left( \Sigma_G^\infty \tilde{E}G \right)^{(2)} & \longrightarrow & \mathcal{L}U_2 \times_{\Sigma_2} t(k_G)^{(2)} \\ \downarrow \xi_2 & & & & \downarrow \xi'_2 \\ S_G^0 & \xrightarrow{\eta} & & & t(k_G) \end{array}$$

where  $\xi_2$  and  $\xi'_2$  are the structural maps for  $S_G^0$  and  $t(k_G)$ . Next we recall that the upper-left corner of this diagram is an equivariant suspension spectrum, so that

we may pass to the adjoint to get a  $G$ -homotopy commutative diagram of spaces. More precisely, [3, Proposition VI.5.3] gives an isomorphism

$$\mathcal{L}U_2 \times_{\Sigma_2} (S^0)^{(2)} \cong \Sigma_G^\infty(\mathcal{L}U_{2+} \wedge_{\Sigma_2} (S^0)^{(2)})$$

which carries  $\xi_2$  to the composite

$$\Sigma_G^\infty(\mathcal{L}U_{2+} \wedge (S^0)^{(2)}) = \Sigma_G^\infty(\mathcal{L}U_2/\Sigma_2)_+ \xrightarrow{\Sigma_G^\infty \pi} \Sigma_G^\infty S^0;$$

here  $\pi$  is the evident projection  $(\mathcal{L}U_2/\Sigma_2)_+ \rightarrow S^0$ . Thus the adjoint of the diagram above has the form

$$\begin{array}{ccc} \mathcal{L}U_{2+} \wedge_{\Sigma_2} (S^0)^{(2)} & \xrightarrow{1 \wedge_{\Sigma_2} \iota^{(2)}} & \mathcal{L}U_{2+} \wedge \tilde{E}G^{(2)} \longrightarrow \Omega_G^\infty(\mathcal{L}U_2 \times_{\Sigma_2} t(k_G)^{(2)}) \\ \downarrow = & & \\ (\mathcal{L}U_2/\Sigma_2)_+ & & \downarrow \Omega_G^\infty \xi'_2 \\ \downarrow \pi & & \\ S^0 & \xrightarrow{\tilde{\eta}} & \Omega_G^\infty t(k_G) \end{array}$$

For our purposes, the important thing about this diagram is that  $\tilde{\eta} \circ \pi$  factors, up to  $G$ -homotopy, through  $\mathcal{L}U_{2+} \wedge \tilde{E}G^{(2)}$ . Precomposing with the projection

$$\mathcal{L}U_{2+} \wedge (S^0)^{(2)} \rightarrow \mathcal{L}U_{2+} \wedge_{\Sigma_2} (S^0)^{(2)},$$

we see that the composite

$$(1) \quad \mathcal{L}U_{2+} \wedge (S^0)^{(2)} = (\mathcal{L}U_2)_+ \xrightarrow{\pi} S^0 \xrightarrow{\tilde{\eta}} \Omega_G^\infty t(k_G)$$

(where we have again written  $\pi$  for the evident projection) factors up to  $G \times \Sigma_2$ -homotopy through  $\mathcal{L}U_{2+} \wedge \tilde{E}G^{(2)}$ . Now let  $H$  be the diagonal copy of  $Z/2$  in  $G \times \Sigma_2$ . Passing to the  $H$ -fixed points of (1) (and noting that the  $H$ -fixed points of  $\Omega_G^\infty t(k_G)$  are the same as the  $G$ -fixed points since  $\Sigma_2$  acts trivially), we see that the composite

$$(2) \quad (\mathcal{L}U_2^H)_+ \xrightarrow{\pi^H} S^0 \xrightarrow{\tilde{\eta}^G} (\Omega_G^\infty t(k_G))^G$$

factors up to (nonequivariant) homotopy through

$$\mathcal{L}U_{2+}^H \wedge (\tilde{E}G^{(2)})^H.$$

But we have shown in the first paragraph of this section that  $(\tilde{E}G^{(2)})^H$  is contractible, so composite (2) is (nonequivariantly) homotopy trivial. We also showed in the first paragraph that  $\mathcal{L}U_2^H$  is contractible, so  $\pi^H$  is an equivalence, and we conclude that

$$\tilde{\eta}^G : S^0 \rightarrow (\Omega_G^\infty t(k_G))^G$$

is homotopy trivial. This means that  $\tilde{\eta}$  is  $G$ -homotopy trivial, and passing to the adjoint, we see that  $\eta$  itself is  $G$ -homotopy trivial. But  $\eta$  is the unit of the equivariant  $E_\infty$  ring  $t(k_G)$ , so  $t(k_G)$  must be equivariantly contractible, as was to be shown.

So far we have assumed that  $G$  is  $Z/2$ . When  $G$  is finite of order  $n$  the action of  $G$  on itself by multiplication induces a homomorphism  $\rho : G \rightarrow \Sigma_n$ , and one can repeat the argument given above with  $\mathcal{L}U_2$  replaced by  $\mathcal{L}U_n$ ,  $\Sigma_2$  replaced by  $\Sigma_n$ , and  $H$  replaced by the subgroup of  $G \times \Sigma_n$  consisting of elements of the form  $(g, \rho(g))$ .  $\square$

## REFERENCES

1. A. K. Bousfield and D. M. Kan, *Homotopy limits, completions and localizations*, Lecture Notes in Math., vol. 304, Springer-Verlag, Berlin and New York, 1972. MR **51**:1825
2. J. P. C. Greenlees and J. P. May, *Generalized Tate, Borel and co Borel cohomology*, preprint.
3. L. G. Lewis, J. P. May, and M. Steinberger, *Equivariant stable homotopy theory*, Lecture Notes in Math., vol. 1213, Springer-Verlag, Berlin and New York, 1986. MR **88e**:55002

DEPARTMENT OF MATHEMATICS, MATHEMATICAL SCIENCES BUILDING, PURDUE UNIVERSITY,  
WEST LAFAYETTE, INDIANA 47907-1395