

ON C^* -ALGEBRAS ASSOCIATED WITH LOCALLY COMPACT GROUPS

M. B. BEKKA, E. KANIUTH, A. T. LAU, AND G. SCHLICHTING

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. Let G be a locally compact group, and let G_d denote the same group G with the discrete topology. There are various C^* -algebras associated to G and G_d . We are concerned with the question of when these C^* -algebras are isomorphic. This is intimately related to amenability. The results can be reformulated in terms of Fourier and Fourier-Stieltjes algebras and of weak containment properties of unitary representations.

INTRODUCTION

Let G be a locally compact group and G_d the group G equipped with the discrete topology. Denote by λ_G and λ_{G_d} the left regular representation of G and G_d on $L^2(G)$ and $\ell^2(G_d)$, respectively. Let $C_\delta^*(G)$ and $C_r^*(G_d)$ be the C^* -subalgebras of $\mathcal{L}(L^2(G))$ and $\mathcal{L}(\ell^2(G_d))$ generated by $\lambda_G(G)$ and $\lambda_{G_d}(G_d)$, respectively.

When G is abelian, $C_r^*(G_d)$ via Fourier transform is isomorphic to the algebra $C(\widehat{G_d})$ of all continuous functions on the compact dual group $\widehat{G_d}$ of G_d . Similarly, $C_\delta^*(G)$ is isomorphic to the norm closed subalgebra of $L^\infty(\widehat{G})$ generated by the elements of G_d considered as functions on $\Gamma = \widehat{G}$ by means of the canonical isomorphism between \widehat{G} and $\widehat{\Gamma}$. By a classical result, $\widehat{G_d}$ is equal to the Bohr compactification $b\Gamma = (\widehat{\Gamma})_d$ of Γ . Moreover, the algebra $C(b\Gamma)$ of almost periodic functions on Γ considered as a subalgebra of $L^\infty(\Gamma)$ is generated by the set $G_d = \widehat{\Gamma}_d = \widehat{b\Gamma}$ of continuous characters on Γ . Hence, $C_\delta^*(G)$ is isomorphic to $C_r^*(G_d)$.

In this paper, we shall be concerned with possible extensions of this result to non-abelian groups.

Relationships between the two unital C^* -algebras $C_\delta^*(G)$ and $C_r^*(G_d)$ were studied by Dunkl and Ramirez [DuR] and by Bédos [Béd]. It was shown in [DuR], Theorem 2.5 (see also [Béd], Lemma 2) that λ_{G_d} extends to a (surjective) $*$ -homomorphism

$$\Phi : C_\delta^*(G) \rightarrow C_r^*(G_d).$$

Received by the editors January 27, 1995 and, in revised form, March 8, 1995 and April 3, 1995.

1991 *Mathematics Subject Classification*. Primary 43A30, 22D10.

Key words and phrases. Amenable group, connected Lie group, group C^* -algebra, weak containment, Fourier algebra.

This research is supported by a NATO collaborative research grant 940184.

This means that λ_G , when viewed as a unitary representation of G_d , weakly contains λ_{G_d} (concerning the notion of weak containment, we refer to [Dix], Chap. 18). For the convenience of the reader, we reproduce below (Proposition 1) a very short proof of this fact appearing in [BeV], Proposition 1.

A natural question is whether Φ is an isomorphism (that is, whether Φ is injective). This is easily settled in case G is amenable (see also [DuR], [Béd]): Φ is an isomorphism if and only if G_d is amenable. Indeed, in this case, by Hulanicki's theorem (see [Pat], Theorem 4.21 or [Pie], Theorem 8.9) the trivial one-dimensional representation 1_G is weakly contained in λ_G (as representations of G and, a fortiori, as representations of G_d). The claim now follows, as G_d is amenable if and only if 1_G is weakly contained in λ_{G_d} .

Our main result in this paper is a complete answer to the question raised above.

Theorem 1. *Let G be a locally compact group. Then $\Phi : C_\delta^*(G) \rightarrow C_r^*(G_d)$ is an isomorphism if and only if G contains an open subgroup H such that H_d is amenable.*

It is well-known that, for a connected Lie group G , G_d is amenable if and only if G is solvable. So, the following is an immediate consequence of Theorem 1.

Corollary 1. *Let G be a connected Lie group. Then Φ is an isomorphism if and only if G is solvable.*

In terms of weak containment, Theorem 1 states that λ_G is weakly contained in λ_{G_d} if and only if G contains an open subgroup H such that H_d is amenable. In fact, we shall prove the following stronger result:

Theorem 2. *Let G be a locally compact group. The following are equivalent:*

- (i) λ_{G_d} weakly contains λ_G ;
- (ii) λ_{G_d} weakly contains some continuous unitary representation of G , viewed as representation of G_d ;
- (iii) G contains an open subgroup H such that H_d is amenable.

As is to be expected, the difficult part in the proof of this theorem is to show that (ii) implies (iii). This will require several steps, the main one being the case where G is a connected Lie group. The proof in this situation is based on the existence, for G non-solvable, of many non-abelian free subgroups of G and on estimates for the norm of convolution operators on free groups, similar to those appearing in [Kes], [Lei] and [AkO].

We now reformulate our results in terms of the Fourier and Fourier-Stieltjes algebras of G and G_d . Recall that the Fourier-Stieltjes algebra $B(G)$ of a locally compact group G is the linear span of all continuous positive definite functions on G . Recall also that $B(G)$ may be identified, in a natural way, with the dual of $C^*(G)$, the C^* -algebra of G . The Fourier algebra $A(G)$ of G is the closed subalgebra, generated by all functions in $B(G)$ with compact support. $A(G)$ may also be described as the set of all matrix coefficients of the regular representation λ_G of G . More details on $A(G)$ and $B(G)$ are to be found in [Eym] where these spaces are extensively studied.

Let $\overline{A(G_d)}^{w^*}$ denote the closure of $A(G_d)$ in $B(G_d)$, with respect to the weak* topology $\sigma(B(G_d), C^*(G_d))$.

Then $\overline{A(G_d)}^{w^*}$ coincides with the space, denoted by $B_{\lambda_d}(G_d)$ in [Eym], of all matrix coefficients of the unitary representations of G_d which are the linear span

of the positive definite functions associated with the unitary representations which are weakly contained in λ_{G_d} (see [Eym], (2.1) Proposition). Hence, the equality $\overline{A(G_d)}^{w*} = B_{\lambda_d}(G_d)$ follows from [Eym], (1.2.1) Proposition.

As any continuous function in $B(G_d)$ actually lies in $B(G)$ ([Eym], (2.24) Corollaire 1), it is now clear that Theorem 2 may be reformulated as follows.

Theorem 2'. *Let G be a locally compact group. The following are equivalent:*

- (i) $\overline{A(G_d)}^{w*}$ contains $A(G)$;
- (ii) $\overline{A(G_d)}^{w*}$ contains a non-zero continuous function on G ;
- (iii) G contains an open subgroup H such that H_d is amenable.

This paper is organized as follows. In Section 1, we treat the case of connected Lie groups. The proof of Theorem 2 is then completed in Section 2. Section 3 contains some remarks about other C^* -algebras associated with G .

1. THE CONNECTED LIE GROUP CASE

In this section, we show that (ii) implies (iii) in Theorem 2 when G is a connected Lie group.

As mentioned above, we first reproduce the proof given in [BeV] for the existence of Φ .

Proposition 1. *λ_{G_d} is weakly contained in λ_G , for any locally compact group G .*

Proof. Since λ_{G_d} is cyclic, it suffices to show that δ_e , the Dirac function at the group unit e , is the pointwise limit of positive definite functions associated to λ_G . Let F be a finite subset of $G \setminus \{e\}$. Choose a neighbourhood K of e such that $gK \cap K = \emptyset$ for all $g \in F$. Set

$$\varphi(g) = \frac{1}{\mu(K)} \langle \lambda_G(g) \chi_K, \chi_K \rangle, \quad g \in G,$$

where χ_K is the characteristic function of K , and μ is a left Haar measure on G . Then φ is a positive definite function associated to λ_G such that $\varphi(g) = \delta_e(g)$ for all $g \in F \cup \{e\}$. □

We shall use the following estimate for the norm of convolution operators on free groups. This may easily be deduced from more general results appearing in [AkO], [Kes] and [Lei]. For the sake of completeness, we prefer to give an independent, short proof (compare also [BCH], Proof of Lemma 2.2).

Proposition 2. *Let Γ be a non-abelian free group on the generators x and y , and let λ denote the left regular representation of Γ on $\ell^2(\Gamma)$. Then*

$$\left\| \sum_{n=1}^{\infty} a_n \lambda(y^n x y^{-n}) \right\| \leq 2 \|a\|_2$$

for all sequences $a = (a_n)_{n \in \mathbb{N}}$ in $\ell^2(\mathbb{N})$.

Proof. Let W_0 be the subset of Γ consisting of the words which do not begin with a nontrivial power of y . Let $W_n = y^n W_0$ for $n \in \mathbb{Z}$. Then $W_n \cap W_m = \emptyset$ for all $n \neq m$. For any $f, g \in \ell^2(\Gamma)$ and $n \in \mathbb{Z}$, the following holds

$$\begin{aligned} |\langle \lambda(y^n x y^{-n}) f, g \rangle| &= |\langle \lambda(y^n x y^{-n}) \chi_{W_n} f, g \rangle| + |\langle \lambda(y^n x y^{-n}) \chi_{\Gamma \setminus W_n} f, g \rangle| \\ &\leq \|\chi_{W_n} f\| \|g\| + \|f\| \|\chi_{y^n x y^{-n} \Gamma \setminus W_n} g\|, \end{aligned}$$

where χ_A denotes the characteristic function of $A \subseteq \Gamma$. Since $y^n xy^{-n}(\Gamma \setminus W_n) \subseteq W_n$, we get

$$|\langle \lambda(y^n xy^{-n})f, g \rangle| \leq \|\chi_{W_n} f\| \|g\| + \|f\| \|\chi_{W_n} g\|$$

and therefore

$$\begin{aligned} |\langle \sum_{n=1}^{\infty} a_n \lambda(y^n xy^{-n})f, g \rangle| &\leq \sum_{n=1}^{\infty} |a_n| (\|\chi_{W_n} f\| \|g\| + \|f\| \|\chi_{W_n} g\|) \\ &\leq 2\|a\|_2 \|f\| \|g\|, \end{aligned}$$

by the Cauchy-Schwarz inequality. □

Proposition 3. *Let G be a connected non-solvable Lie group. Then no continuous unitary representation of G , viewed as a representation of G_d , is weakly contained in λ_{G_d} .*

Proof. It is well-known that such a group contains a non-abelian free subgroup F on two generators a and b (see [Pat], Theorem 3.9). For any finite set of integers $i_1, \dots, i_n \in \mathbb{Z} \setminus \{0\}$, let $p_{i_1 \dots i_n} : G \rightarrow G$ denote the word function

$$p_{i_1 \dots i_n}(x) = \begin{cases} a^{i_1} x^{i_2} \dots a^{i_{n-1}} x^{i_n}, & \text{if } n \text{ is even,} \\ a^{i_1} x^{i_2} \dots x^{i_{n-1}} a^{i_n}, & \text{if } n \text{ is odd.} \end{cases}$$

Then, the set

$$G_{i_1 \dots i_n} = \{x \in G, p_{i_1 \dots i_n}(x) \neq e\}$$

is open. It is also nonempty since $b \in G_{i_1 \dots i_n}$. Moreover, because $p_{i_1 \dots i_n}$ is an analytic function on G , $G_{i_1 \dots i_n}$ is dense. So, by Baire's category theorem, the intersection

$$X = \bigcap \{G_{i_1 \dots i_n}; i_1, \dots, i_n \in \mathbb{Z} \setminus \{0\}\}$$

is dense in G .

By the definition of X , for any $x \in X$, the subgroup Γ_x generated by a and x is a free group. Hence, by Proposition 2, for any $x \in X$ and $N \in \mathbb{N}$, we have the following estimate:

$$(*) \quad \left\| \frac{1}{N} \sum_{n=1}^N \lambda_{\Gamma_x}(a^n x a^{-n}) \right\| \leq \frac{2}{\sqrt{N}}$$

where λ_{Γ_x} denotes the regular representation of the discrete group Γ_x . But, since the restriction of λ_{G_d} to Γ_x is a multiple of λ_{Γ_x} , we may replace λ_{Γ_x} by λ_{G_d} in the above inequality (*).

Now suppose, by contradiction, that there exists a continuous unitary representation π of G which is weakly contained in λ_{G_d} . Then, by (*),

$$\left\| \frac{1}{N} \sum_{n=1}^N \pi(a^n x a^{-n}) \right\| \leq \frac{2}{\sqrt{N}}$$

for any $x \in X$ and any $N \in \mathbb{N}$. Therefore, for any unit vector ξ in the Hilbert space of π , we have

$$(**) \quad \left| \frac{1}{N} \sum_{n=1}^N \langle \pi(a^n x a^{-n}) \xi, \xi \rangle \right| \leq \frac{2}{\sqrt{N}}$$

for all $x \in X$ and $N \in \mathbb{N}$. Since X is dense and π is (strongly) continuous, (**) holds for any $x \in G$.

Taking $x = e$ and $N \geq 5$, we reach the contradiction

$$1 = \frac{1}{N} \sum_{n=1}^N \langle \pi(a^n e a^{-n}) \xi, \xi \rangle \leq \frac{2}{\sqrt{5}}.$$

□

2. THE GENERAL CASE

We now proceed with the proof of Theorem 2. This will require two results related to Proposition 3.

Proposition 4. *Let G be an amenable locally compact group. Assume that λ_{G_d} weakly contains some continuous unitary representation π of G . Then G_d is amenable.*

Proof. Let $\bar{\pi}$ denote the representation conjugate to π . Then the (inner) tensor product $\pi \otimes \bar{\pi}$ is weakly contained in $\lambda_{G_d} \otimes \bar{\lambda}_{G_d}$ and, hence, in λ_{G_d} , since $\lambda_{G_d} \otimes \bar{\lambda}_{G_d}$ is a multiple of λ_{G_d} .

On the other hand, because G is amenable, the trivial representation 1_G is weakly contained in $\pi \otimes \bar{\pi}$, by [Bek], Theorems 2.2 and 5.1. Hence, 1_G is weakly contained in λ_{G_d} . Therefore, G_d is amenable. □

Next, we extend Proposition 3 to all connected groups.

Proposition 5. *Let G be a connected locally compact group. Assume that λ_{G_d} weakly contains some continuous unitary representation π of G . Then G_d is amenable.*

Proof. By the structure theory for connected groups, G contains a compact normal subgroup K such that G/K is a Lie group (see [MoZ], p. 175).

We first claim that K_d is amenable. Indeed, since the restriction $\lambda_{G_d}|_K$ of λ_{G_d} to K is a multiple of λ_{K_d} , λ_{K_d} weakly contains $\pi|_K$. As K is amenable, the claim follows from Proposition 4.

Suppose, by contradiction, that $(G/K)_d$ is not amenable. Then, by the proof of Proposition 3, there exist an element $\dot{a} \in G/K$ and a dense subset \dot{X} of G/K such that, for any $\dot{x} \in \dot{X}$, the subgroup generated by \dot{a} and \dot{x} is free.

Let $p : G \rightarrow G/K$ denote the canonical projection. Choose any $a \in G$ with $p(a) = \dot{a}$, and set $X = p^{-1}(\dot{X})$. Then, for any $x \in X$, the subgroup of G generated by a and x is free. As in the proof of Proposition 3, this, together with the fact that X is dense in G , yields a contradiction.

Therefore, $(G/K)_d$ and K_d are amenable. It follows that G_d is amenable. □

Proof of Theorem 2. That (i) implies (ii) is obvious. Suppose (ii) holds, that is, λ_{G_d} weakly contains some continuous representation π of G . Let G^0 denote the connected component of e in G . As $\lambda_{G_d^0}$ weakly contains $\pi|_{G^0}$, G_d^0 is amenable by

Proposition 5. Since G/G^0 is totally disconnected, we may choose an open subgroup H of G containing G^0 such that H/G^0 is compact (see [HeR], Theorem 7.7). We claim that H_d is amenable. Indeed since H is amenable, the claim follows immediately from Proposition 4. This completes the proof that (ii) implies (iii).

Suppose (iii) holds, that is, G contains an open subgroup H such that H_d is amenable. Then λ_{H_d} weakly contains λ_H (in fact, λ_{H_d} weakly contains any unitary representation of H_d).

Now, since G/H is discrete, the induced representation $\text{ind}_{H_d}^{G_d} \lambda_H$ is equivalent to $\text{ind}_H^G \lambda_H = \lambda_G$. Therefore, by continuity of inducing, λ_G is weakly contained in $\text{ind}_{H_d}^{G_d} \lambda_{H_d} = \lambda_{G_d}$. This shows that (i) holds and completes the proof of Theorem 2. \square

3. SOME REMARKS ON OTHER C^* -ALGEBRAS ASSOCIATED WITH G

Let ω be the universal representation of the locally compact group G . Let $C^*(G_d)$ be the (maximal) C^* -algebra of G_d , and let $C_{\delta,\omega}^*(G)$ denote the C^* -algebra generated by all operators $\omega(x)$, $x \in G$. There is an obvious surjective $*$ -homomorphism

$$\Psi : C^*(G_d) \rightarrow C_{\delta,\omega}^*(G).$$

The main result in [BeV] may be reformulated as follows.

Theorem 3 ([BeV]). *Let G be a connected Lie group. Then Ψ is an isomorphism if and only if G is solvable.*

Remark 1. We do not know how to extend this result to other locally compact groups. A reasonable conjecture seems to be: Ψ is an isomorphism if and only if G contains an open subgroup H such that H_d is amenable (that is, by Theorem 1, if and only if Φ is an isomorphism).

Let $\Lambda : C_{\delta,\omega}^*(G) \rightarrow C_\delta^*(G)$ be the surjective $*$ -homomorphism, defined in the natural way. The following result, for which we offer a simple proof, may also be deduced from [Béd, Theorem 1].

Theorem 4. *Let G be a locally compact group. Then Λ is an isomorphism if and only if G is amenable.*

Proof. If G is amenable, then λ_G is weakly equivalent to the universal representation ω (even as representations of G). This implies that Λ is an isomorphism. Conversely, assume Λ is injective. Then 1_G is weakly contained in λ_G , where both representations are viewed as representations of G_d . That is, G has Reiter's weak property (P_2^*) (see [Pie], p. 56) which is known to characterise the amenability of G . \square

Remark 2. We used in the above proof the surprising fact that Reiter's weak property (P_2^*) is equivalent to the stronger property (P_2) (saying that 1_G is weakly contained in λ_G , as representations of G) and, hence, to the amenability of G . Usually, the proof of this equivalence is a long and tedious one involving topological invariant means (compare [Pie], p. 56). It is worth mentioning that the argument used by Bédos in [Béd], Proof of Theorem 1, provides a quick and elegant proof for this equivalence. Indeed, assume 1_G is weakly contained in λ_G , as representations

of G_d . Then 1_G defines a state φ on $C_\delta^*(G)$ such that $\varphi(\lambda_G(x)) = 1$ for all $x \in G$. Extend φ to a state $\tilde{\varphi}$ on $\mathcal{L}(L^2(G))$. Cauchy-Schwarz inequality shows that

$$\tilde{\varphi}(\lambda_G(x)T) = \tilde{\varphi}(T\lambda_G(x))$$

for all $x \in G$, $T \in \mathcal{L}(L^2(G))$.

Then, denoting by M_f the multiplication operator on $L^2(G)$ by $f \in L^\infty(G)$, one defines a mean m on $L^\infty(G)$ as follows:

$$m(f) = \tilde{\varphi}(M_f), \quad \forall f \in L^\infty(G).$$

Since

$$M_{\lambda_G(x)f} = \lambda_G(x)M_f\lambda_G(x^{-1}), \quad \forall x \in G, \quad f \in L^\infty(G),$$

m is left invariant. This shows that G is amenable.

It should be observed that, using the notion of amenable representations as defined in [Bek], the above argument shows that λ_G is amenable.

The following corollary is also proved in [DuR], Proposition 3.2.

Corollary 2. *Let G be a locally compact group. Then*

$$\Phi \circ \Lambda : C_{\delta,\omega}^*(G) \rightarrow C_r^*(G_d)$$

is an isomorphism if and only if G_d is amenable.

Proof. If G_d is amenable, then, as shown earlier, Λ and Φ are isomorphisms.

Conversely, suppose $\Phi \circ \Lambda$ is an isomorphism. Since Λ is an isomorphism, by Theorem 4, G is amenable. As mentioned in the introduction, this implies that G_d is amenable, as Φ is an isomorphism. \square

REFERENCES

- [AkO] C. Akemann and P. Ostrand, *Computing norms in group C*-algebras*, Amer. J. Math. **98** (1976), 1015–1048. MR **56**:1079
- [Béd] E. Bédos, *On the C*-algebra generated by the left regular representation of a locally compact group*, Proc. Amer. Math. Soc. **120** (1994), 603–608. MR **94d**:22004
- [BCH] M.B. Bekka, M. Cowling and P. de la Harpe, *Some groups whose reduced C*-algebra is simple*, Publ. Math. IHES **80** (1994), 117–134.
- [Bek] M.B. Bekka, *Amenable unitary representations of locally compact groups*, Invent. Math. **100** (1990), 383–401. MR **91g**:22007
- [BeV] M.B. Bekka and A. Valette, *On duals of Lie groups made discrete*, J. reine angew. Math. **439** (1993), 1–10. MR **94k**:22003
- [Dix] J. Dixmier, *C*-algebras*, North-Holland, 1977. MR **56**:16388
- [DuR] C. Dunkl and D. Ramirez, *C*-algebras generated by Fourier-Stieltjes transforms*, Trans. Amer. Math. Soc. **164** (1972), 435–441. MR **46**:9646
- [Eym] P. Eymard, *L’algèbre de Fourier d’un groupe localement compact*, Bull. Soc. Math. France **92** (1964), 181–236. MR **37**:4208
- [HeR] E. Hewitt and K. Ross, *Abstract Harmonic Analysis I*, Springer-Verlag, 1994.
- [Kes] H. Kesten, *Symmetric random walks on groups*, Trans. Amer. Math. Soc. **92** (1959), 336–354. MR **22**:253
- [Lei] M. Leinert, *Faltungsoperatoren auf gewissen diskreten Gruppen*, Studia Math. **52** (1974), 149–158. MR **50**:7594

- [MoZ] D. Montgomery and L. Zippin, *Topological transformation groups*, Interscience (1955). MR **17**:383
- [Pat] A. Paterson, *Amenability*, Amer. Math. Soc. (1988). MR **90e**:43001
- [Pie] J.P. Pier, *Amenable Locally Compact Groups*, Wiley, 1984. MR **86a**:43001

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITY DE METZ, UFR M.I.M. ILE DU SAULCY, F-57045 METZ CEDEX 01, FRANCE

E-mail address: `bekka@poncelet.univ-metz.fr`

FACHBEREICH MATHEMATIK-INFORMATIK, UNIVERSITÄT-GH PADERBORN, WARBURGER STR. 100, D-33098 PADERBORN, GERMANY

E-mail address: `kaniuth@uni-paderborn.de`

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF ALBERTA, EDMONTON, ALBERTA, CANADA T6G 2G1

E-mail address: `tlau@vega.math.ualberta.ca`

MATHEMATISCHES INSTITUT, TECHNISCHE UNIVERSITÄT MÜNCHEN, ARCSSTRASSE 21, W-80333 MÜNCHEN 2, GERMANY

E-mail address: `gschlich@mathematik.tu-muenchen.de`