

AN EXTENSION OF THE VITALI-HAHN-SAKS THEOREM

ONÉSIMO HERNÁNDEZ-LERMA AND JEAN B. LASSERRE

(Communicated by Christopher D. Sogge)

ABSTRACT. The Vitali-Hahn-Saks theorem on the absolute continuity of the setwise limit of a sequence of bounded measures is extended to allow unbounded measures and convergence of integrals of continuous functions vanishing at infinity.

1. INTRODUCTION

A useful version of the Vitali-Hahn-Saks theorem gives conditions for a measure λ which is the setwise limit of a sequence of absolutely continuous (a.c.) measures $\{\lambda_n\}$ to be a.c. This result allows the underlying measure space (X, \mathcal{B}) to be arbitrary (cf. [2], p. 155), but it requires that $\{\lambda_n\}$ and λ , as well as the dominant measure, say μ , are all finite.

In this paper, we show that the same conclusion ($\lambda \ll \mu$) holds without requiring either *setwise* convergence or *finiteness* of the measures involved, but we impose in particular suitable topological conditions on X .

Moreover, we give conditions for the Radon-Nikodym derivative $d\lambda/d\mu$ to be in $L_p(X, \mathcal{B}, \mu)$ for $1 \leq p \leq \infty$.

In addition to being interesting in themselves, these results are also useful to study the existence of solutions to the Poisson Equation for a Markov kernel [3].

2. MAIN RESULTS

Let (X, \mathcal{B}, μ) be a σ -finite complete measure space, where X is a locally compact separable metric space, and \mathcal{B} is the completion (with respect to μ) of the σ -algebra of Borel subsets of X . Let $C_0(X)$ be the space of real-valued continuous functions on X vanishing at infinity. (Concerning the convergence in (2.1), see Remark 2.3(b) at the end of this section.)

Received by the editors January 24, 1995.

1991 *Mathematics Subject Classification*. Primary 28A33; Secondary 28C15.

Key words and phrases. Measures, setwise convergence, absolute continuity.

The research of the first author was partially supported by a visiting professorship at Paul Sabatier University, Toulouse, France, and by CONACYT Grant 1332-E9206.

This research was partially supported by the CNRS (France)- CONACYT (México) Scientific Cooperation Program.

Theorem 2.1. *Let $\{\lambda_n\}$ and λ be σ -finite (nonnegative) measures on (X, \mathcal{B}) such that, as $n \rightarrow \infty$,*

$$(2.1) \quad \int_X v d\lambda_n \rightarrow \int_X v d\lambda \quad \forall v \in C_0(X).$$

If, in addition, every λ_n is a.c. with respect to μ , then so is λ .

Suppose that the measures λ and λ_n in Theorem 2.1 are finite. Then, by the Radon-Nikodym theorem, there are nonnegative measurable functions u_n and u such that

$$\lambda_n(B) = \int_B u_n d\mu \quad \text{and} \quad \lambda(B) = \int_B u d\mu \quad \forall B \in \mathcal{B}.$$

The next theorem gives conditions for u to be in $L_p := L_p(X, \mathcal{B}, \mu)$, $1 \leq p \leq \infty$.

Theorem 2.2. *Fix $1 \leq p \leq \infty$. In addition to the hypotheses of Theorem 2.1, suppose that $u_n \in L_p \forall n$ and that, for some constant M , $\liminf_n \|u_n\|_p \leq M$. Then*

$$(2.2) \quad u \text{ is in } L_p.$$

Remark 2.3. (a) If the λ_n are all finite, then in Theorem 2.1, instead of (2.1), we only need to assume that $\lim_n \int_X v d\lambda_n$ exists and is finite for all $v \in C_0(X)$. Then, from [2], Theorem VIII.8, it follows that the sequence $\{\lambda_n\}$ converges to a measure λ , as in (2.1).

(b) Endowed with the sup-norm, $C_0(X)$ is a Banach space and its dual is the space $M(X)$ of finite signed measures on X with total variation norm [2],[5]. Thus, when the λ_n are finite measures, (2.1) can be stated equivalently as: λ_n converges to λ in the weak* topology $\sigma(M(X), C_0(X))$. If λ_n and λ are probability measures and (2.1) holds, it is sometimes said that λ_n converges vaguely to λ . If $C_0(X)$ is replaced by the (larger) space $C(X)$ of all continuous and bounded functions on X , it is said that λ_n converges weakly to λ (see e.g. [1]). It follows from Lemma 3.1(b) and the Portmanteau Theorem ([1], Theorem 2.1) that, in this case, vague convergence implies weak convergence (the converse being obviously true).

(c) To verify (2.1) it suffices to consider nonnegative $v \in C_0(X)$. Indeed, since any $v \in C_0(X)$ can be written as $v = v^+ + v^-$ with $v^+(x) := \max[0, v(x)]$ and $v^-(x) := \min[0, v(x)]$, it suffices to observe that both v^+ and v^- are in $C_0(X)$. Hence, convergence in (2.1) for all $0 \leq v \in C_0$ implies convergence for all $v \in C_0(X)$.

3. PROOFS

The proof of Theorem 2.1 is based on the following lemma, which is well known in the case of finite measures (and bounded function h); see e.g. [2], Theorem VIII.10.

Lemma 3.1. *In the context of Theorem 2.1:*

(a) *If h is a nonnegative and l.s.c. (lower semicontinuous) function on X , then*

$$\liminf_{n \rightarrow \infty} \int_X h d\lambda_n \geq \int_X h d\lambda;$$

(b) $\liminf_{n \rightarrow \infty} \lambda_n(B) \geq \lambda(B)$ *for any open set $B \in \mathcal{B}$.*

Proof. (a) As h is nonnegative and l.s.c., there exists an increasing sequence of nonnegative continuous bounded functions v_k on X such that $v_k(x) \uparrow h(x) \forall x \in X$. Similarly, as each v_k is a nonnegative continuous bounded function and X is σ -compact ([4], p. 203, Theorem 21), for every k there is an increasing sequence $\{v_{kl}, l = 1, 2, \dots\}$ of nonnegative functions v_{kl} in $C_0(X)$ with $v_{kl}(x) \uparrow v_k(x)$ for all $x \in X$ as $l \rightarrow \infty$. Hence,

$$\begin{aligned} \liminf_n \int h d\lambda_n &\geq \liminf_n \int v_k d\lambda_n \quad \forall k \\ &\geq \liminf_n \int v_{kl} d\lambda_n \quad \forall k, l \\ &= \int v_{kl} d\lambda \quad (\text{by (2.1)}). \end{aligned}$$

Thus, letting $l \rightarrow \infty$, and then $k \rightarrow \infty$, the Monotone Convergence Theorem yields (a).

(b) If $B \subset X$ is open, its characteristic (or indicator) function is l.s.c. and of course nonnegative. Hence, (b) follows from (a). \square

Proof of Theorem 2.1. For every measurable set $B \in \mathcal{B}$ and $\epsilon > 0$ there is an open set G_ϵ that contains B and $\mu(G_\epsilon - B) < \epsilon$ (see e.g. [2], p. 41). In particular, if B is a μ -null set, i.e., $\mu(B) = 0$ (and hence $\lambda_n(B) = 0 \forall n$), we have $\mu(G_\epsilon) = \mu(G_\epsilon - B) < \epsilon$. Moreover, by part (b) in Lemma 3.1

$$\liminf_n \lambda_n(G_\epsilon) \geq \lambda(G_\epsilon) \geq \lambda(B).$$

Therefore, letting $\epsilon \downarrow 0$ we obtain $\lambda(B) = 0$. \square

Proof of Theorem 2.2. Consider first the case $p = 1$. From $\liminf_n \|u_n\|_1 \leq M$ and Lemma 3.1(b) (with $B := X$), it immediately follows that $\lambda(X) < \infty$ so that $u \in L_1$.

Consider now the case $p > 1$ and let q be the exponent conjugate to p . Then the condition (2.2) holds if uv is in L_1 for every $v \in L_q$ (see e.g. [5], p. 133). In turn, for this to be true it suffices to show that (as $u \geq 0$) there is a constant M such that

$$(3.1) \quad \int uv d\mu \leq M \|v\|_q \quad \forall v \in L_q^+.$$

Moreover, since (for $1 \leq q < \infty$) the class of continuous functions in L_q is dense in L_q (see e.g. [2], p. 92) it suffices to prove (3.1) for all continuous functions v in L_q^+ . (Indeed, suppose the latter holds, and let $v_m \in L_q^+$ be a sequence of continuous functions converging in the L_q -norm to $v \in L_q^+$. Let v_{m_i} be a subsequence converging to v μ -a.e. Then

$$\begin{aligned} M \|v\|_q &= M \lim_i \|v_{m_i}\|_q \geq \lim_i \int uv_{m_i} d\mu \\ &\geq \int uv d\mu \quad (\text{by Fatou's Lemma}); \end{aligned}$$

i.e., $v \in L_q^+$ satisfies (3.1).) Now, let $v \in L_q^+$ be a continuous function. Hence, (2.1) and Lemma 3.1(a) yield:

$$\begin{aligned} \int uv d\mu = \int v d\lambda &\leq \liminf_n \int v d\lambda_n \\ &= \liminf_n \int v u_n d\mu \\ &\leq \liminf_n \|u_n\|_p \|v\|_q \quad (\text{Hölder's inequality}) \\ &\leq M \|v\|_q, \end{aligned}$$

with M as in Theorem 2.2. As v was an arbitrary continuous function in L_q^+ we obtain (3.1), hence (2.2). \square

REFERENCES

1. P. Billingsley, *Convergence of Probability Measures*, Wiley, New York, 1968. MR **38**:1718
2. J.L. Doob, *Measure Theory*, Springer-Verlag, New York, 1994. MR **95c**:28001
3. O. Hernández-Lerma and J.B. Lasserre, *Existence of solutions to the Poisson equation in L^p spaces*, LAAS Technical report 95045, Toulouse, 1995.
4. H.L. Royden, *Real Analysis*, 3rd Edition, Macmillan, New York, 1988. MR **90g**:00004
5. W. Rudin, *Real and Complex Analysis*, 3rd edition, McGraw-Hill, New York, 1986. MR **87f**:00009

DEPARTAMENTO DE MATEMÁTICAS, CINVESTAV-IPN, APDO. POSTAL 14-740, MÉXICO D.F. 07000, MEXICO

E-mail address: ohernand@math.cinvestav.mx.

LAAS-CNRS, 7 AVENUE DU COLONEL ROCHE, 31077 TOULOUSE CÉDEX, FRANCE

E-mail address: lasserre@laas.fr