

FACTORISATION IN NEST ALGEBRAS

M. ANOUSSIS AND E. G. KATSOULIS

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. We give a necessary and sufficient condition on an operator A for the existence of an operator B in the nest algebra $\text{Alg}N$ of a continuous nest N satisfying $AA^* = BB^*$ (resp. $A^*A = B^*B$). We also characterise the operators A in $B(H)$ which have the following property: For every continuous nest N there exists an operator B_N in $\text{Alg}N$ satisfying $AA^* = B_N B_N^*$ (resp. $A^*A = B_N^* B_N$).

1. INTRODUCTION–PRELIMINARIES

The problem of factorisation of operators with respect to a nest algebra has been studied by many authors [8], [1], [13], [9], [11], [12], [10]. In this work we give a necessary and sufficient condition on an operator A for the existence of an operator B in the nest algebra $\text{Alg}N$ of a continuous nest N satisfying $AA^* = BB^*$ (resp. $A^*A = B^*B$). This result improves Theorem 4.9 in [9] for continuous nests. We also characterise the operators A in $B(H)$ which have the following property: For every continuous nest N there exists an operator B_N in $\text{Alg}N$ satisfying $AA^* = B_N B_N^*$ (resp. $A^*A = B_N^* B_N$).

Throughout this work H denotes a separable Hilbert space and $B(H)$ the space of all bounded operators from H into itself. If V is a subset of H we denote by $[V]$ the linear span of V . By subspace of H we mean a subset of H which is closed under addition of vectors and scalar multiplication. If $\{V_n\}_{n=1}^\infty$ is a sequence of closed mutually orthogonal subspaces of H we denote by $\sum_{n=1}^\infty \oplus V_n$ the closure of their linear span. If A is in $B(H)$ we denote by $r(A)$ the range of A and by $\text{coker}A$ the orthogonal complement of the kernel of A . An operator range is the range of a bounded operator in H . A nest in H is a totally ordered set of closed subspaces of H containing $\{0\}$ and H which is closed under intersection and closed span. If N is a nest in H and P is in N we will denote by the same symbol the orthogonal projection on the subspace P . If N is a nest we denote by N^\perp the nest $\{P^\perp : P \in N\}$. A nest N is continuous if $P = \overline{[\bigcup_{Q < P} Q]}$ for every P in N . Given a nest N the associated nest algebra $\text{Alg}N$ is the set of operators A in $B(H)$ satisfying $PAP = AP$ for every P in N . For a general discussion of nest algebras the reader is referred to [3].

Received by the editors December 6, 1994 and, in revised form, April 5, 1995.
1991 *Mathematics Subject Classification*. Primary 47D25.

2. PROPER SUBSPACES

We introduce in this section the notion of N -proper subspace for a nest N . We show that a closed subspace of H of co-finite dimension is N -proper for every continuous nest N .

Definition 1. Let N be a nest on H . A vector x in H is called N -proper if $x = Px$ for some P in N , $P \neq I$.

Definition 2. Let N be a nest on H . A subspace V of H is called N -proper if $\{V \cap P : P \in N, P \neq I\}$ is dense in V .

Lemma 3. Let N be a continuous nest on H . Let $\{P_n\}_{n=1}^\infty$ be a sequence of elements of N such that: $P_n \neq I$, $P_{n+1} \geq P_n$, and P_n converges strongly to I . Let x_1, x_2, \dots, x_m be orthonormal vectors in H . Set $V = [x_1, x_2, \dots, x_m]^\perp$. Then:

(a) There exists n_0 such that $P_n x_1, P_n x_2, \dots, P_n x_m$ are linearly independent for $n \geq n_0$.

(b) We set $V_1 = P_1 H \ominus P_1 V^\perp$ and we define inductively

$$V_n = P_n H \ominus \left(\sum_{i=1}^{n-1} \oplus V_i \oplus P_n V^\perp \right).$$

Then $V = \sum_{i=1}^\infty \oplus V_i$.

Proof. (a) The Grammian of the vectors $P_n x_1, P_n x_2, \dots, P_n x_m$ converges to the Grammian of the vectors x_1, x_2, \dots, x_m which equals 1.

(b) It is easy to see that the V_n 's are mutually orthogonal and that V_n is contained in V for every n . We show that $(\sum_{i=1}^\infty \oplus V_i) \oplus V^\perp = H$. Let x be a vector in H which is orthogonal to $(\sum_{i=1}^\infty \oplus V_i) \oplus V^\perp$. For each n the vector $P_n x$ is orthogonal to $\sum_{i=1}^n \oplus V_i$ so $P_n x$ is in $P_n V^\perp$. For $n \geq n_0$ we have $P_n x = P_n (\sum_{i=1}^m a_i x_i)$, where the a_i 's are complex numbers not depending on n . So $x = \lim_{n \rightarrow \infty} P_n x = \sum_{i=1}^m a_i x_i$. But x is orthogonal to V^\perp , hence it is 0. \square

Proposition 4. Let N be a continuous nest and V a closed subspace of H of co-finite dimension. Then V is N -proper.

Proof. It follows immediately from Lemma 3. \square

Let N be a continuous nest on H and A an operator in $B(H)$. Consider the set $\bigcup_{P \in N, P \neq I} A^{-1}(P)$. This set is equal to $\bigcup_{P \in N, P \neq I} \text{Ker}(P^\perp A)$. If A is an Alg N the set $\bigcup_{P \in N, P \neq I} A^{-1}(P)$ contains $\bigcup_{P \in N, P \neq I} P$; hence it is dense in H . There exist operators A in $B(H)$ for which $\bigcup_{P \in N, P \neq I} A^{-1}(P)$ is not dense in H . We construct such an operator in Example 9. We will prove in the next section that $\bigcup_{P \in N, P \neq I} A^{-1}(P)$ is dense in H if and only if there exists an operator B in Alg N such that $AA^* = BB^*$. We first prove some preliminary results.

Lemma 5. Let N be a nest on H and A an operator in $B(H)$. The following are equivalent:

(a) The set $\bigcup_{P \in N, P \neq 0} (A^*)^{-1}(P^\perp)$ is dense in H .

(b) $\bigcap_{P \in N, P \neq 0} \overline{r(AP)} = \{0\}$.

Proof. We have that $(A^*)^{-1}(P^\perp) = \{x \in H : A^*x \in P^\perp\} = \{x \in H : P^\perp A^*x = A^*x\} = \text{Ker}(PA^*) = \overline{r(AP)}^\perp$ and $\bigcup_{P \in N, P \neq 0} \overline{r(AP)}^\perp$ is dense in $(\bigcap_{P \in N, P \neq 0} \overline{r(AP)})^\perp$. \square

Proposition 6. *Let N be a nest on H and A an operator in $B(H)$.*

(a) *Suppose that the set $\bigcup_{P \in N, P \neq I} A^{-1}(P)$ is dense in H . Then $r(A)$ is N -proper.*

(b) *Suppose that $r(A)$ is N -proper and closed. Then the set $\bigcup_{P \in N, P \neq I} A^{-1}(P)$ is dense in H .*

Proof. (a) The set $A(\bigcup_{P \in N, P \neq I} A^{-1}(P))$ is contained in $[r(A) \cap P : P \in N, P \neq I]$ and is dense in $r(A)$.

(b) The restriction of A to $\text{coker}A$ is an isomorphism from $\text{coker}A$ onto $r(A)$. Hence $(\bigcup_{P \in N, P \neq I} A^{-1}(P)) \cap \text{coker}A = A^{-1}(\bigcup_{P \in N, P \neq I} P) \cap \text{coker}A$ is dense in $\text{coker}A$. Therefore $\bigcup_{P \in N, P \neq I} A^{-1}(P)$ is dense in H . \square

Proposition 7. *Let N be a nest on H and A an operator in $B(H)$.*

(a) *Suppose that $\bigcap_{P \in N, P \neq 0} \overline{r(AP)} = \{0\}$. Then $\text{coker}A$ is N^\perp -proper.*

(b) *Suppose that $\text{coker}A$ is N^\perp -proper and $r(A)$ is closed. Then $\bigcap_{P \in N, P \neq 0} \overline{r(AP)} = \{0\}$.*

Proof. (a) It follows from Lemma 5 that $\bigcup_{P \in N, P \neq 0} (A^*)^{-1}(P^\perp)$ is dense in H . It follows from Proposition 6 that $r(A^*)$ is N^\perp -proper. Since the closure of an N^\perp -proper subspace is an N^\perp -proper subspace we conclude that $\text{coker}A$ is N^\perp -proper.

(b) It follows from [2, Ch. VI, Th. 1.10] that $r(A^*)$ is closed. Hence $r(A^*) = \text{coker}A$. It follows from Proposition 6 that $\bigcup_{P \in N, P \neq 0} (A^*)^{-1}(P^\perp)$ is dense in H . Therefore from Lemma 5 we conclude that $\bigcap_{P \in N, P \neq 0} \overline{r(AP)} = \{0\}$. \square

3. FACTORISATION

In this section we prove our main results and give some applications.

Theorem 8. *Let N be a continuous nest and A an operator in $B(H)$. The following are equivalent:*

(a) *There exists an operator B in $\text{Alg}N$ such that $AA^* = BB^*$.*

(b) *The set $\bigcup_{P \in N, P \neq I} A^{-1}(P)$ is dense in H .*

Proof. Assume (a) holds. In order to prove (b) it is enough to prove that the set $\bigcup_{P \in N, P \neq I} (A^{-1}(P) \cap \text{coker}A)$ is dense in $\text{coker}A$. Using polar decomposition one can see that there exists a partial isometry U with domain $\text{coker}A$ and range $\text{coker}B$ such that $A = BU$. We put $R = \overline{[\bigcup_{P \in N, P \neq I} (A^{-1}(P) \cap \text{coker}A)]}$ and $M = \text{coker}A \ominus R$. We will show that $M = \{0\}$. Take m in M and P in N , $P \neq I$. Since $r(A) = r(B)$ ([5, Th. 1]), we have $BPUm = Ax_P$ for some x_P in $\text{coker}A$. Since $BPUm$ is in P , x_P is in $A^{-1}(P) \cap \text{coker}A$ and hence in R . We have $BPUm = Ax_P = BUx_P$ and so $PUm - Ux_P$ is in $\ker B$. We have $PUm = PUm - Ux_P + Ux_P$ which belongs to $\ker B \oplus UR$. Note that the decomposition $H = \ker B \oplus UR \oplus UM$ is orthogonal. Therefore $Um = \lim_{P \in N, P \neq I, P \rightarrow I} PUm$ is in $(\ker B \oplus UR) \cap UM = \{0\}$. We conclude that $m = 0$.

Assume (b) holds. It is then clear that the set $\bigcup_{P \in N, P \neq I} (A^{-1}(P) \cap \text{coker}A)$ is dense in $\text{coker}A$. Take a sequence $\{P_n\}_{n=0}^\infty$ of elements of N such that: $P_0 = 0$, $P_{n+1} > P_n$, $P_n \neq I$ for every n and P_n converges strongly to I . We set: $R_1 = A^{-1}(P_1) \cap \text{coker}A$, $R_n = (A^{-1}(P_n) \cap \text{coker}A) \ominus R_{n-1}$ for $n > 1$.

It is clear that R_n is orthogonal to R_m for $n \neq m$ and that R_n is contained in $\text{coker}A$ for every n . We show that $\text{coker}A = \sum_{n=1}^\infty \oplus R_n$. Take y in $\text{coker}A$. If y is orthogonal to $\sum_{n=1}^\infty \oplus R_n$, then y is orthogonal to $A^{-1}(P_n) \cap \text{coker}A$ for every

n ; hence y is orthogonal to $(\bigcup_{n=1}^{\infty} (A^{-1}(P_n) \cap \text{coker}A))$. Since $(\bigcup_{n=1}^{\infty} (A^{-1}(P_n) \cap \text{coker}A))$ is dense in $\text{coker}A$, $y = 0$, and so $\text{coker}A = \sum_{n=1}^{\infty} \oplus R_n$.

Consider for $n \geq 1$ a partial isometry V_n with domain contained in $(P_{n+1} - P_n)H$ and range R_n . Put $V = \sum_{n=1}^{\infty} \oplus V_n$. Then V is a partial isometry with range $\text{coker}A$. Note that $A = AVV^*$. We show that AV belongs to $\text{Alg}N$. Let P be in N and x be a vector in P . We show that AVx is in P . If $P \leq P_1$ we have $AVx = 0$. If $P > P_1$ there exists $m \geq 1$ such that $P_m < P \leq P_{m+1}$. Then $AVx = A(\sum_{n=1}^m \oplus V_n)x$ and $(\sum_{n=1}^m \oplus V_n)x$ is contained in $(\sum_{n=1}^m \oplus R_n)$. Therefore AVx is in $A(\sum_{n=1}^m \oplus R_n)$ which is contained in P_m . Since $P_m < P$ we conclude that AVx is in P .

Put $B = AV$. Then $BB^* = AVV^*A^* = AA^*$ and B is in $\text{Alg}N$. \square

Remark. Theorem 8 remains true under the weaker assumption that N is a nest which satisfies $H = \overline{[\bigcup_{Q < H} Q]}$.

Let N be a continuous nest. We give an example of an operator with N -proper range which does not satisfy condition (b) of Theorem 8.

Example 9. Let N be a continuous nest. Take a sequence $\{P_n\}_{n=0}^{\infty}$ of elements of N such that:

$$P_0 = 0, \quad P_{n+1} > P_n, \quad P_n \neq I \quad \text{for every } n \text{ and } P_n \text{ converges strongly to } I.$$

For each n consider a vector e_n of norm 1 and such that $(P_{n+1} - P_n)e_n = e_n$. Put $y = \sum_{i=1}^{\infty} n^{-1}e_n$. Let A be the operator defined by: $Ae_n = n^{-1}e_n$ for $n \geq 1$, $Ae_0 = y$ and A is 0 on $[e_n : n = 0, 1, 2, \dots]^{\perp}$. Then $r(A)$ is N -proper and it is easy to see that A does not satisfy condition (b) of Theorem 8. In fact, e_0 is orthogonal to $\bigcup_{P \in N, P \neq I} A^{-1}(P)$. So A does not satisfy condition (a) of Theorem 8.

Corollary 10. *Let N be a continuous nest and A an operator in $B(H)$. The following are equivalent:*

- (a) *There exists an operator B in $\text{Alg}N$ such that $A^*A = B^*B$.*
- (b) $\bigcap_{P \in N, P \neq 0} \overline{r(AP)} = \{0\}$.

Proof. There exists an operator B in $\text{Alg}N$ such that $A^*A = B^*B$ if and only if there exists an operator C in $\text{Alg}N^{\perp}$ such that $A^*A = CC^*$. The corollary follows now from Theorem 8 and Lemma 5. \square

Corollary 11. *Let N be a continuous nest and A an operator in $B(H)$. Suppose A is onto (resp. one-to-one and $r(A)$ is closed). Then there exists an operator B in $\text{Alg}N$ such that $AA^* = BB^*$ (resp. $A^*A = B^*B$).*

Proof. It follows from Proposition 6 and Theorem 8 (resp. from Proposition 7 and Corollary 10). \square

Corollary 12. *Let N be a continuous nest and Q a projection in $B(H)$. Then there exists an operator B in $\text{Alg}N$ such that $Q = BB^*$ (resp. $Q = B^*B$) if and only if QH is N -proper (resp. N^{\perp} -proper).*

Proof. It follows from Proposition 6 and Theorem 8 (resp. from Proposition 7 and Corollary 10). \square

The following corollary answers a question posed by Shields in [13].

Corollary 13. *Let N be a continuous nest and A a positive operator in $B(H)$. Assume there exists an operator B in $\text{Alg}N$ such that $A^2 = B^*B$. Then there exists an operator C in $\text{Alg}N$ such that $A = C^*C$.*

Proof. We have to show that if $\bigcap_{P \in N, P \neq 0} \overline{r(AP)} = \{0\}$, then $\bigcap_{P \in N, P \neq 0} \overline{r(A^{1/2}P)} = \{0\}$. Let y be in $\bigcap_{P \in N, P \neq 0} \overline{r(A^{1/2}P)}$. Then $A^{1/2}y$ is in $\bigcap_{P \in N, P \neq 0} \overline{r(AP)} = \{0\}$; hence $A^{1/2}y = 0$. So y is in $\text{Ker } A^{1/2}$. Since y is also in $\overline{r(A^{1/2})}$ we see that $y = 0$. \square

We will characterise now the operators that satisfy condition (a) of Theorem 8 (resp. condition (a) of Corollary 10) for every continuous nest.

Proposition 14. *Let V be an operator range. Assume V is not of co-finite dimension in H . Then there exists a continuous nest N in H such that $V \cap P = \{0\}$ for every P in N , $P \neq I$.*

Proof. (i) We first show that there exists a non-closed operator range W which contains V . We will use the following fact: If V_1, V_2 are operator ranges, then $V_1 + V_2$ is an operator range [7, Ch. I, 1]. If V is closed we consider an operator range U which is non-closed and is contained in V^\perp . We set $W = V + U$. Then W is an operator range which is non-closed and contains V .

(ii) It follows from (i) above that we may assume that V is non-closed. An operator range R is called of type J_S (Dixmier's notation) if it is dense and there exists a sequence $\{H_n\}_{n=0}^\infty$ of closed mutually orthogonal infinite dimensional subspaces of H such that $R = \{\sum_{n=0}^\infty x_n : x_n \in H_n \text{ and } \sum_{n=0}^\infty (2^n \|x_n\|)^2 < \infty\}$. It is shown in the proof of Theorem 3.6 in [6] that any non-closed operator range is contained in an operator range of type J_S . It follows that there exists an operator range S of type J_S such that $V \subset S$. It follows from Theorem 3.6 in [6] that there exists a unitary operator U on H such that $US \cap S = \{0\}$. We conclude that there exists an operator range T of type J_S such that $V \cap T = \{0\}$. Now it is easy to see that there exists a continuous nest N in H such that $P \subset T$ for every P in N , $P \neq I$. It follows that $P \cap V = \{0\}$ for every P in N , $P \neq I$. \square

Theorem 15. *Let A be an operator in $B(H)$.*

(a) *There exists for every continuous nest N an operator B_N in $\text{Alg}N$ satisfying $AA^* = B_N B_N^*$ if and only if A is a right Fredholm operator.*

(b) *There exists for every continuous nest N an operator B_N in $\text{Alg}N$ satisfying $A^*A = B_N^* B_N$ if and only if A is a left Fredholm operator.*

Proof. (a) Assume that for every continuous nest N there exists an operator B_N in $\text{Alg}N$ satisfying $AA^* = B_N B_N^*$. It follows from Theorem 8 and Proposition 6 that $r(A)$ is N -proper for every continuous nest N . Proposition 14 implies that $r(A)$ is of co-finite dimension in H . If the range of an operator is of co-finite dimension, then it is closed [4, Prop. 3.7]. Therefore A is a right Fredholm operator. Assume now that A is a right Fredholm operator. Then $r(A)$ is closed and of co-finite dimension in H . By Proposition 4, $r(A)$ is N -proper for every continuous nest N . It follows then from Proposition 6 and Theorem 8 that for every continuous nest N there exists an operator B_N in $\text{Alg}N$ satisfying $AA^* = B_N B_N^*$.

(b) Consider the following properties of an operator A :

(i) *There exists for every continuous nest N an operator B_N in $\text{Alg}N$ satisfying $AA^* = B_N B_N^*$.*

- (ii) There exists for every continuous nest N an operator B_N in $\text{Alg}N$ satisfying $A^*A = B_N^*B_N$.

Since a nest N is continuous if and only if the nest N^\perp is continuous we see that an operator A has property (i) if and only if the operator A^* has property (ii). The assertion follows now from (a). \square

ADDED IN PROOF

After this work was submitted a paper of G. T. Adams, J. Froelich, P. J. McGuire, and V. I. Paulsen entitled *Analytic reproducing kernels and factorisation*, Indiana Univ. Math. J. **43** (1994), came to our attention. Condition (b) of our Theorem 8 is essentially the same with the density condition given in Theorem 3.1 of this paper in a different but related context.

REFERENCES

1. W. B. Arveson, *Interpolation problems in nest algebras*, J. Funct. Anal. **20** (1975), 208–233. MR **52**:3979
2. J. B. Conway, *A course in functional analysis*, Springer-Verlag, 1985. MR **86h**:46001
3. K. R. Davidson, *Nest algebras*, Pitman Research Notes in Mathematics Series, **191** (1988). MR **90f**:47062
4. J. Dixmier, *Etude sur les variétés et les opérateurs de Julia*, Bull. Soc. Math. France **77** (1949), 11–101. MR **11**:369f
5. R. G. Douglas, *On majorization, factorization and range inclusion of operators in Hilbert space*, Proc. Amer. Math. Soc. **17** (1966), 413–416. MR **34**:3315
6. P. A. Fillmore and J. P. Williams, *On operator ranges*, Advances in Math. **7** (1971), 254–281. MR **45**:2518
7. C. Foias, *Invariant para-closed subspaces*, Indiana Univ. Math. J. **21** (1972), 887–906. MR **53**:3734
8. I. C. Gohberg and M. G. Krein, *Theory and applications of Volterra operators in Hilbert space*, Transl. Math. Monographs, **24** (1970), AMS. MR **41**:9041
9. D. R. Larson, *Nest algebras and similarity transformations*, Ann. of Math. **121** (1985), 409–427. MR **86j**:47061
10. D. R. Pitts, *Factorization problems for nests: Factorization methods and characterizations of the universal factorization property*, J. Funct. Ana. **79** (1988), 57–90. MR **90a**:46160
11. S. C. Power, *Nuclear operators in nest algebras*, J. Operator Theory **10** (1983), 337–352. MR **85b**:47028
12. S. C. Power, *Factorisation in analytic operator algebras*, J. Funct. Anal. **67** (1986), 413–432. MR **87k**:47040
13. A. L. Shields, *An analogue of a Hardy-Littlewood-Fejer inequality for upper triangular trace class operators*, Math. Z. **182** (1983), 473–484. MR **85c**:47022

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF THE AEGEAN, KARLOVASSI 83200, GREECE

DEPARTMENT OF MATHEMATICS, EAST CAROLINA UNIVERSITY, GREENVILLE, NORTH CAROLINA 27858