

HILBERT C^* -MODULES IN WHICH ALL CLOSED SUBMODULES ARE COMPLEMENTED

BOJAN MAGAJNA

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. Let B be a C^* -algebra. If there exists a full Hilbert B -module X such that $X = Y \oplus Y^\perp$ for each closed submodule Y , then B is $*$ -isomorphic to a C^* -algebra of (not necessarily all) compact operators on a Hilbert space.

1. INTRODUCTION AND THE MAIN RESULT

It is well known that each closed subspace \mathcal{K} of a Hilbert space \mathcal{H} has an orthogonal complement \mathcal{K}^\perp such that $\mathcal{K} \oplus \mathcal{K}^\perp = \mathcal{H}$. (In fact, by [7] a similar property characterizes Hilbert spaces among all Banach spaces.) This property, however, does not hold in general for closed submodules of a Hilbert module over a C^* -algebra (see [5] or [9]; in [6] Hilbert modules that are complemented in any larger Hilbert module are characterized). In this note we shall characterize all C^* -algebras B such that all closed submodules in any Hilbert module over B are orthogonally complemented. Here a submodule Y of a Hilbert module X is called orthogonally complemented if there exists a self-adjoint projection p on X such that $Y = pX$ (see [5, p. 21]). More precisely, we shall prove the following result.

Theorem 1. *Let B be a C^* -algebra. If there exists a full Hilbert B -module in which every closed submodule is orthogonally complemented, then B is $*$ -isomorphic to a C^* -algebra of (not necessarily all) compact operators on some Hilbert space. Consequently, all closed submodules in all Hilbert B -modules are orthogonally complemented.*

2. PRELIMINARIES

A (right) Hilbert module over a C^* -algebra B is a right B -module X equipped with a B -valued inner product $\langle \cdot, \cdot \rangle$, which is linear over B in the second and conjugate linear in the first variable, such that X is a Banach space with the norm $\|x\| = \|\langle x, x \rangle\|^{1/2}$. Hilbert modules are important for operator K -theory, and the basic theory of such modules can be found in [4] and [5]. We denote by L (or by $L(X)$ if X is not clear from the context) the C^* -algebra of all operators (necessarily bounded and linear over B) on X that have an adjoint. For arbitrary $x, y \in X$ there is an operator $[y, x] \in L$ defined by $[y, x]z = y\langle x, z \rangle$ ($z \in X$). The closed linear span of such operators (which is the analogy for compact operators between

Received by the editors July 3, 1995 and, in revised form, September 26, 1995.

1991 *Mathematics Subject Classification.* Primary 46L05; Secondary 46C50.

Key words and phrases. Hilbert C^* -modules, complemented submodules.

Hilbert spaces) will be denoted by K (or $K(X)$). K is a closed two-sided ideal (not necessarily proper) in L .

For a general Hilbert B -module X the closed linear span of the set $\{\langle x, y \rangle : x, y \in X\}$ is a closed two-sided ideal in B and will be denoted by $\langle X, X \rangle$. If $\langle X, X \rangle = B$ then X is called a *full Hilbert B -module*.

The subset Λ of $L(X \oplus B)$ consisting of all matrices of the form

$$\begin{bmatrix} a & x \\ y^* & b \end{bmatrix},$$

where $a \in K$, $b \in B$ and $x, y \in X$, is a C^* -subalgebra. Here each $y \in X$ is regarded as an operator from B to X defined by $b \mapsto yb$, with the adjoint given by $y^*(x) = \langle y, x \rangle$. This algebra was introduced in [3] and called the *linking algebra of X* .

We shall regard B , X and $K = K$ as subsets of Λ in the obvious way:

$$B \cong \begin{bmatrix} 0 & 0 \\ 0 & B \end{bmatrix}, \quad X \cong \begin{bmatrix} 0 & X \\ 0 & 0 \end{bmatrix} \quad \text{and} \quad K \cong \begin{bmatrix} K & 0 \\ 0 & 0 \end{bmatrix}.$$

Note that the inner product $\langle x, y \rangle$ in X can be expressed as the product x^*y in Λ .

3. PROOF OF THE THEOREM

To prove the theorem we shall relate submodules of X to right ideals of the linking algebra Λ of X , but first we need a couple of simple lemmas.

Lemma 1. *If X is a Hilbert B -module such that every closed submodule of X is orthogonally complemented, then each closed right ideal J of $K = K(X)$ is of the form $J = pK$ for some projection $p \in L = L(X)$.*

Proof. Consider the closed submodule $[JX]$ of X . By hypothesis $[JX] = pX$ for some projection $p \in L$. Then $pax = ax$ for all $x \in X$ and $a \in J$, hence $pa = a$ and consequently $J \subseteq pK$. To prove the reverse inclusion, it suffices to show that $p[x, y] = [px, y]$ is in J for each 'rank one' operator $[x, y] \in K$ (since K is spanned by such operators). Since px can be approximated by finite sums of elements of the form az ($a \in J$, $z \in X$), $[px, y]$ can be approximated by finite sums of operators of the form $[az, y] = a[z, y]$, which are in J . This proves that $J = pK$. \square

Lemma 2. *If every closed right ideal J in a C^* -algebra A is of the form $J = pA$ for some projection p in the multiplier algebra $M(A)$ of A , then A is $*$ -isomorphic to a C^* -algebra of (not necessarily all) compact operators on some Hilbert space.*

Proof. First suppose that A is commutative, hence $A \cong C_0(T)$ and $M(A) \cong C_b(T)$ (= bounded continuous functions on a locally compact Hausdorff space T). By hypothesis for each $t \in T$ there exists a projection $p \in C_b(T)$ such that

$$\{f \in C_0(T) : f(t) = 0\} = pC_0(T).$$

Clearly such a p must be the characteristic function of the complement of $\{t\}$, hence each singleton in T is open and T is discrete. Consequently A is $*$ -isomorphic to a C^* -algebra of (diagonal) compact operators on some Hilbert space.

In general, we shall prove that for each maximal commutative C^* -subalgebra A_0 of A every maximal ideal J_0 of A_0 is of the form $J_0 = pA_0$ for some projection $p \in M(A_0)$. By previous paragraph this implies that the maximal ideal space of A_0 is discrete, hence by [2, p. 84] A is $*$ -isomorphic to a subalgebra of $K(\mathcal{H})$ for some

Hilbert space \mathcal{H} . Given a maximal ideal J_0 of A_0 , consider the closed right ideal $J = [J_0A]$ of A . By hypothesis $J = pA$ for some projection $p \in M(A)$. We may assume that $A \subseteq B(\mathcal{H})$ for some Hilbert space \mathcal{H} such that $[A\mathcal{H}] = \mathcal{H}$; then $M(A) = \{b \in A'' : bA + Ab \subseteq A\}$. Note that $[J_0\mathcal{H}] = [J_0A\mathcal{H}] = [J\mathcal{H}] = [pA\mathcal{H}] = p\mathcal{H}$; in particular the range of p is invariant under A'_0 , therefore $p \in A''_0$. From $p \in M(A)$ it then follows that $pA_0 \subseteq A''_0 \cap A$. Since $A''_0 \cap A$ is a commutative C*-subalgebra of A and contains A_0 , while A_0 is maximal commutative in A , $A''_0 \cap A = A_0$. Thus $pA_0 \subseteq A_0$, which implies that $p \in M(A_0)$.

Since the range of p is $[J_0\mathcal{H}]$, we have $J_0 \subseteq pA_0$; but, since J_0 is a maximal ideal in A_0 , the required equality $J_0 = pA_0$ follows if $pA_0 \neq A_0$. To see that $pA_0 \neq A_0$, let ρ_0 be a character on A_0 with $\ker \rho_0 = J_0$ and let ρ be a pure state on A extending ρ_0 . Then J is contained in the right kernel of ρ , hence A_0 is not contained in J . On the other hand, $pA_0 \subseteq J$ (since $J = pA$). □

Proof of Theorem 1. Let X be a full Hilbert module over B such that all closed submodules of X are orthogonally complemented. Then by Lemmas 1 and 2 there exists a *-monomorphism $\pi_K : K \rightarrow K(\mathcal{H}_K)$ for some Hilbert space \mathcal{H}_K (such that $[\pi_K(K)\mathcal{H}_K] = \mathcal{H}_K$). By a well known result concerning extensions of representations (see [8, p. 162]) there exist a Hilbert space $\mathcal{H} \supseteq \mathcal{H}_K$ and a representation π of the linking algebra Λ of X on \mathcal{H} such that $\pi(a)|_{\mathcal{H}_K} = \pi_K(a)$ for each $a \in K$. Since π is a *-representation we have

$$(1) \quad \pi(x)^*\pi(x) = \pi(\langle x, x \rangle) \quad \text{and} \quad \pi(x)\pi(x)^* = \pi([x, x])$$

for all $x \in X$. Put $\mathcal{H}_B = [\pi(X)^*\mathcal{H}_K]$ and note that \mathcal{H}_B is invariant under $\pi(B)$ since $BX^* = (XB)^* \subseteq X^*$. Since $BK = 0$, $KB = 0$, $XK = 0$ and $X^*X^* = 0$ (the products are computed in Λ), the subspaces $\mathcal{H}_K = [\pi(K)\mathcal{H}_K]$ and \mathcal{H}_B are orthogonal and we have

$$(2) \quad \pi(B)\mathcal{H}_K = 0, \quad \pi(K)\mathcal{H}_B = 0, \quad \pi(X)\mathcal{H}_K = 0 \quad \text{and} \quad \pi(X)^*\mathcal{H}_B = 0.$$

Moreover,

$$[\pi(X)\mathcal{H}_B] = [\pi(X)\pi(X)^*\mathcal{H}_K] = [\pi(K)\mathcal{H}_K] = \mathcal{H}_K.$$

These relations easily imply that $\mathcal{H}_K \oplus \mathcal{H}_B$ is invariant under π ; hence, replacing π by the corresponding restriction, we may assume that $\mathcal{H}_K \oplus \mathcal{H}_B = \mathcal{H}$. Then the second identity in (1) (together with the second identity in (2) and the fact that $\pi_K(K)$ consists of compact operators) implies that $\pi(x)$ is compact for all $x \in X$. Since $B = \langle X, X \rangle$, it follows then from the first identity in (1) that $\pi(B)$ consists of compact operators only. Moreover, since π_K is one to one, it is easy to see (using (1)) that the restriction of π to $(X$ and to B is one to one. (If $J \stackrel{\text{def}}{=} \ker \pi|_B \neq 0$, then $[XJ, XJ]$ would be a non-zero ideal of K contained in $\ker \pi_K$.) Thus B is *-isomorphic to a C*-subalgebra of $K(\mathcal{H}_B)$.

Conversely, suppose that B is isomorphic to a (nondegenerate) C*-algebra of compact operators on a Hilbert space \mathcal{H}_B by a *-isomorphism π_B . Let π be a representation of Λ on a Hilbert space $\mathcal{H} \supseteq \mathcal{H}_B$ such that $\pi(b)|_{\mathcal{H}_B} = \pi_B(b)$ for each $b \in B$, and put $\mathcal{H}_K = [\pi(X)\mathcal{H}_B]$. As in the previous paragraph, we may assume that $\mathcal{H}_K \oplus \mathcal{H}_B = \mathcal{H}$. Since π_B is one to one it follows (using (1)) that the restrictions of π to X and K are one to one, but then we see (using (2)) that π is also one to one. Thus, Λ is *-isomorphic to a C*-subalgebra of $K(\mathcal{H})$, hence by [1, 1.4.5] Λ is *-isomorphic to a (restricted) direct sum of elementary C*-algebras of the form $K(\mathcal{H}_j)$ for some Hilbert spaces \mathcal{H}_j . Since $K(\mathcal{H}_j)$ is strongly Morita

equivalent to \mathbb{C} , it would not be hard to see (using induced representations, see [10] or [5]) that $K(\mathcal{H}_j)$ and \mathbb{C} have equivalent categories of Hilbert modules, from which it is possible to conclude the proof, but we shall give here a more elementary approach. Since Λ is $*$ -isomorphic to a direct sum of algebras of the form $K(\mathcal{H}_j)$, each closed right ideal J of Λ is of the form $J = p_\Lambda \Lambda$ for some projection p_Λ in the multiplier algebra $M(\Lambda)$ of Λ . Using the above representation of Λ and the well known fact that the multiplier algebra of K is L (see [5]), it is easy to see that $M(\Lambda)$ is of the form

$$M(\Lambda) = \begin{bmatrix} L & Z \\ Z^* & M(B) \end{bmatrix},$$

where Z is a certain $L - M(B)$ -subbimodule of $B(\mathcal{H}_B, \mathcal{H}_K)$ containing X . Given a closed submodule Y of X , consider the closed right ideal J of Λ generated by Y , that is

$$J = \begin{bmatrix} [Y, X] & Y \\ 0 & 0 \end{bmatrix}.$$

If p_Λ is a projection in $M(\Lambda)$ such that $J = p_\Lambda \Lambda$, then a straightforward computation shows that p_Λ is necessarily of the form

$$p_\Lambda = \begin{bmatrix} p & 0 \\ 0 & 0 \end{bmatrix}$$

for some projection $p \in L$, and that $Y = pX$. □

REFERENCES

- [1] W. B. Arveson, *An invitation to C^* -algebras*, GTM **39**, Springer-Verlag, Berlin, 1976. MR **58**:23621
- [2] B. A. Barnes, G. J. Murphy, M. R. F. Smyth and T. T. West, *Riesz and Fredholm theory in Banach algebras*, Pitman Research Notes in Math. **67**, Pitman, 1982. MR **84a**:46108
- [3] L. G. Brown, P. Green and M. A. Rieffel, *Stable isomorphism and strong Morita equivalence of C^* -algebras*, Pacific J. Math. **71** (1977), 349–363. MR **57**:3866
- [4] K. Jensen and K. Thomsen, *Elements of KK -theory*, Birkhäuser, Basel, 1991. MR **94b**:19008
- [5] E. C. Lance, *Hilbert C^* -modules*, London Math. Soc Lecture Note Series **210**, Cambridge Univ. Press, Cambridge, 1995. CMP 95:10
- [6] H. Lin, *Injective Hilbert C^* -modules*, Pacific J. Math. **154** (1992), 131–164. MR **93c**:46110
- [7] J. Lindenstrauss and L. Tzafriri, On complemented subspaces problem, Israel J. Math. **9** (1971), 263–269. MR **43**:2474
- [8] G. J. Murphy, *C^* -algebras and operator theory*, Academic Press, London, 1990. MR **91m**:46084
- [9] W. Paschke, *Inner product modules over B^* -algebras*, Trans. Amer. Math. Soc. **182** (1973), 443–468. MR **50**:8087
- [10] M. Rieffel, *Induced representations of C^* -algebras*, Advances in Math. **13** (1974), 176–257. MR **50**:5489

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF LJUBLJANA, JADRANSKA 19, LJUBLJANA 1000, SLOVENIA

E-mail address: Bojan.Magajna@uni-lj.si