

## NOTE ON CLARK'S THEOREM FOR $p$ -ADIC CONVERGENCE

MINORU SETOYANAGI

(Communicated by Dennis A. Hejhal)

ABSTRACT. We must read Clark's statement under the hypothesis that the *negative* of each zero of the indicial polynomial is non-Liouville. In this note we shall give the example for which under the original hypothesis the statement does not hold.

### INTRODUCTION

Clark's article [1] is referred to by many authors because he first studied the  $p$ -adic convergence of power series solutions at a singular point and pointed out the importance of  $p$ -adic non-Liouvilleness of exponents. However, his result is not properly stated. We must read his statement under the hypothesis that the *negative* of each zero of the indicial polynomial is non-Liouville. In this note we shall give the example for which under the original hypothesis the statement does not hold. For another approach to  $p$ -adic convergence using the notion of  $p$ -adic Liouvilleness in the sense of Schikhof [2], see [3].

Throughout this note  $\mathbb{C}_p$  denotes the completion of the algebraic closure of  $\mathbb{Q}_p$ , and the  $p$ -adic valuation  $|\cdot|_p$  is simply written  $|\cdot|$ . By  $\mathbb{N}$  we denote the positive integers, and by  $\mathbb{N}^0$  the non-negative integers.  $\mathbb{Z}$  and  $\mathbb{Z}_p$  denote the rational integers and the ring of  $p$ -adic integers of  $\mathbb{Q}_p$ , respectively.

### 1. SOME DEFINITIONS

The statement in question is Clark's Theorem 3 [1]. We shall give a summary of this below. To understand it we refer to some definitions in his article.

**Definition 1.** An element  $\alpha \in \mathbb{C}_p$  is said to be ( $p$ -adically) *non-Liouville* if for  $s \in \mathbb{N}$ , we have

$$\text{ord}(\alpha + s) = O(\log s) \quad \text{as } s \rightarrow \infty.$$

**Definition 2.** For each non-Liouville number  $\alpha$  the *weight*  $w(\alpha)$  is defined by Clark. In this note we only use

$$w(\alpha) = \frac{1}{p-1}$$

for  $\alpha \in \mathbb{Z}_p$ .

---

Received by the editors October 8, 1995.

1991 *Mathematics Subject Classification.* Primary 12H25; Secondary 11S80, 34G05.

*Key words and phrases.*  $p$ -adic convergence,  $p$ -adically non-Liouville number.

**Definition 3.** The *weight*  $w(g)$  of a polynomial  $g \in \mathbb{C}_p[X]$  whose roots are non-Liouville is the sum of the weights of the roots.

The *ordinal* of a polynomial  $\Phi \in \mathbb{C}_p[X]$ , say  $\sum_{i=0}^n a_i X^i$ , is defined by

$$\text{ord } \Phi = \min_i \{\text{ord } a_i\}.$$

(Notice that “minimal” is carelessly mistaken for “maximal” in his article.)

We can summarize Clark’s statement as follows:

Consider a linear differential equation

$$(1) \quad A_n(x)y^{(n)} + A_{n-1}(x)y^{(n-1)} + \cdots + A_0(x)y = 0$$

where coefficients lie in  $\mathbb{C}_p[[x]]$  with nonzero radius of convergence. We may write

$$A_i = x^i \sum_{j=0}^{\infty} a_{ij} x^j \quad (0 \leq i \leq n) \text{ and } a_{i0} \neq 0 \text{ for some } i. \text{ Let } \Phi_j(s) = \sum_{i=0}^n a_{ij} \binom{s}{i} i! \text{ for}$$

$j \geq 0$ ; then  $\Phi_0$  is the indicial polynomial of (1) at  $x = 0$ .

**“Statement”.** *If every zero of the indicial polynomial of (1) is non-Liouville, then each power series solution converges for*

$$\text{ord } x > \sup_{j \geq 1} \left\{ \frac{\text{ord } \Phi_0 - \text{ord } \Phi_j}{j} \right\} + w(\Phi_0).$$

## 2. A NON-LIOUVILLE NUMBER

In order to give the example for which the Statement does not hold, we define a special non-Liouville number in the following lemma.

**Lemma.** *Let  $(e_k)_{k=1}^{\infty}$  be a sequence of positive integers given by  $e_1 = 1$  and  $e_{k+1} = kp^{e_k}$  ( $k \geq 1$ ), and let  $\lambda = \sum_{k=1}^{\infty} p^{e_k}$ . Then we have*

- (i)  $\varliminf_{n \rightarrow \infty} \sqrt[n]{|n - \lambda|} = 0$ ;
- (ii)  $\lambda \in \mathbb{Z}_p \setminus \mathbb{Z}$ ;
- (iii)  $\text{ord}(\lambda + n) = O(\log n)$  as  $n \rightarrow \infty$ .

*Proof.* (i) Let  $n_k = \sum_{j=1}^k p^{e_j}$ . Then  $(n_k)_{k=1}^{\infty}$  is a strictly increasing sequence of positive integers, and we have

$$\sqrt[n_k]{|\lambda - n_k|} = p^{-e_{k+1}/n_k}.$$

Since  $p^{e_k} \leq n_k < 2p^{e_k}$ , it follows that

$$-\frac{e_{k+1}}{n_k} < -\frac{e_{k+1}}{2p^{e_k}} = -\frac{k}{2},$$

and hence

$$\sqrt[n_k]{|\lambda - n_k|} < p^{-\frac{k}{2}}.$$

Thus we have

$$\lim_{k \rightarrow \infty} \sqrt[n_k]{|\lambda - n_k|} = 0,$$

and therefore

$$\varliminf_{n \rightarrow \infty} \sqrt[n]{|\lambda - n|} = 0.$$

(ii) Clearly  $\lambda \in \mathbb{Z}_p \setminus \mathbb{N}^0$ . If  $-\lambda \in \mathbb{N}$ , that is, there is a positive integer  $m$  such that  $-\lambda = m$ , then we have

$$\sqrt[n]{|n - \lambda|} = |n + m|^{\frac{1}{n}} \geq \left(\frac{1}{n + m}\right)^{\frac{1}{n+m} \frac{n+m}{n}},$$

and therefore

$$\lim_{n \rightarrow \infty} \sqrt[n]{|\lambda - n|} = 1,$$

which is contrary to (i).

(iii) Let  $n \geq p$ , and let

$$n = a_0 + a_1p + \dots + a_l p^l$$

be the  $p$ -adic expansion of  $n$ , where  $a_l \neq 0$ . Then, as  $p^l \leq n < p^{l+1}$ , we have

$$0 \leq \text{ord}(\lambda + n) \leq l + 2 \leq \frac{\log n}{\log p} + 2,$$

and therefore

$$0 \leq \frac{\text{ord}(\lambda + n)}{\log n} \leq \frac{3}{\log p}.$$

Hence

$$\text{ord}(\lambda + n) = O(\log n) \quad \text{as } n \rightarrow \infty.$$

□

### 3. EXAMPLE

Now, let  $\lambda$  be the non-Liouville number defined above. Consider the linear differential equation

$$(2) \quad (1 - x)x^2y'' + \{(1 - \lambda) - (2 - \lambda)x\}xy' + \lambda xy = 0.$$

In this equation, we have

$$\begin{aligned} A_2 &= x^2(1 - x), \\ A_1 &= x\{(1 - \lambda) - (2 - \lambda)x\}, \\ A_0 &= \lambda x, \end{aligned}$$

and so

$$\begin{aligned} a_{20} &= 1, & a_{21} &= -1, \\ a_{10} &= 1 - \lambda, & a_{11} &= -(2 - \lambda), \\ a_{00} &= 0, & a_{01} &= \lambda, \\ a_{ij} &= 0 \quad \text{if } j \geq 2. \end{aligned}$$

As  $\Phi_j(s) = a_{0j} + a_{1j}s + a_{2j}s(s - 1)$ , we have

$$\begin{aligned} \Phi_0(s) &= (1 - \lambda)s + s(s - 1) = s(s - \lambda), \\ \Phi_1(s) &= \lambda - (2 - \lambda)s - s(s - 1) = -s^2 - (1 - \lambda)s + \lambda, \\ \Phi_j(s) &= 0 \quad \text{if } j \geq 2. \end{aligned}$$

The roots of the indicial equation are 0 and  $\lambda$ , and so the requirement of the Statement is satisfied. Since  $\text{ord } \lambda \geq 0$  by Lemma (ii), we have

$$\begin{aligned}\text{ord } \Phi_0 &= \min\{\text{ord } 1, \text{ord } \lambda\} = 0, \\ \text{ord } \Phi_1 &= \min\{\text{ord } 1, \text{ord}(1 - \lambda), \text{ord } \lambda\} = 0, \\ \text{ord } \Phi_j &= \infty \quad \text{if } j \geq 2.\end{aligned}$$

This gives

$$\begin{aligned}\sup_{j \geq 1} \left\{ \frac{\text{ord } \Phi_0 - \text{ord } \Phi_j}{j} \right\} + w(\Phi_0) &= \sup_{j \geq 1} \left\{ \frac{-\text{ord } \Phi_j}{j} \right\} + w(0) + w(\lambda) \\ &= \frac{2}{p-1}.\end{aligned}$$

According to the Statement, the power series solution of (2) must converge for

$$\text{ord } x > \frac{2}{p-1},$$

but the equation (2) has the power series solution

$$y = \sum_{n=0}^{\infty} \frac{x^n}{n - \lambda},$$

whose radius of convergence is zero by virtue of Lemma (i).

#### 4. "COROLLARY"

Clark's original proof of Theorem 3 needs the Corollary on p.266 of his article:

**"Corollary"**. *Let  $h$  be a monic polynomial whose roots are non-Liouville and in  $\{\alpha \in \mathbb{C}_p : |\alpha| \leq 1\}$ , and let  $t$  be a fixed positive integer greater than all roots of  $h$  in  $\mathbb{Z}$ . If  $s, s' \in \mathbb{Z}, s > s' > t$ , then*

$$\sum_{j=s'}^s \text{ord } h(j) = w(h)(s - s') + O(\log s).$$

Professor Dwork suggested to the author that a similar error appears in this statement. Indeed, if it were true we should have the following claim, which is contrary to our Lemma.

*Claim.* Let  $\alpha \in \mathbb{Z}_p \setminus \mathbb{Z}$  such that  $\text{ord}(\alpha + s) = O(\log s)$ . Then

$$\underline{\lim}_{s \rightarrow \infty} \sqrt[s]{|s - \alpha|} \geq p^{-\frac{1}{p-1}}.$$

In fact, let  $h(X) = X - \alpha$ . This is the monic polynomial whose root is only  $\alpha$ . Therefore

$$w(h) = w(\alpha) = \frac{1}{p-1}.$$

By the Corollary we have

$$\sum_{j=s'}^s \text{ord}(j - \alpha) = \frac{s - s'}{p-1} + O(\log s),$$

where  $s > s' > t$ . Thus

$$0 \leq \text{ord}(s - \alpha) \leq \sum_{j=s'}^s \text{ord}(j - \alpha) \leq \frac{s-t}{p-1} + M \log s$$

for some  $M > 0$ , so that

$$|s - \alpha| \geq p^{-\frac{s-t}{p-1} - M \log s},$$

and therefore

$$\underline{\lim}_{s \rightarrow \infty} \sqrt[s]{|s - \alpha|} \geq p^{-\frac{1}{p-1}}.$$

However, this claim contradicts our Lemma when  $\alpha = \lambda$ .

#### ACKNOWLEDGEMENT

The author would like to thank Professor Dwork for his valuable comments and kind advice.

#### REFERENCES

1. D. N. Clark, *A note on the  $p$ -adic convergence of solutions of linear differential equations*, Proc. Amer. Math. Soc. **17** (1966), 262–269. MR **32**:4350
2. W. Schikhof, *Ultrametric calculus*, Cambridge Univ. Press, Cambridge, 1984. MR **86j**:11104
3. M. Setoyanagi, *On the convergence of solutions of  $p$ -adic linear differential equations*, preprint.

MAIZURU NATIONAL COLLEGE OF TECHNOLOGY, 234 SHIRAYA, MAIZURU, KYOTO 625, JAPAN  
*E-mail address*: set@maizuru-ct.ac.jp