

ON THE VON NEUMANN-JORDAN CONSTANT FOR BANACH SPACES

MIKIO KATO AND YASUJI TAKAHASHI

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. Let $C_{\text{NJ}}(E)$ be the von Neumann-Jordan constant for a Banach space E . It is known that $1 \leq C_{\text{NJ}}(E) \leq 2$ for any Banach space E ; and E is a Hilbert space if and only if $C_{\text{NJ}}(E) = 1$. We show that: (i) If E is uniformly convex, $C_{\text{NJ}}(E)$ is less than two; and conversely the condition $C_{\text{NJ}}(E) < 2$ implies that E admits an equivalent uniformly convex norm. Hence, denoting by $\tilde{C}_{\text{NJ}}(E)$ the infimum of all von Neumann-Jordan constants for equivalent norms of E , E is super-reflexive if and only if $\tilde{C}_{\text{NJ}}(E) < 2$. (ii) If $\tilde{C}_{\text{NJ}}(E) = 2^{2/p-1}$, $1 < p \leq 2$ (the same value as that of L_p -space), E is of Rademacher type r and cotype r' for any r with $1 \leq r < p$, where $1/r + 1/r' = 1$; the converse holds if E is a Banach lattice and l_p is finitely representable in E or E' .

INTRODUCTION AND PRELIMINARIES

In connection with the famous work [8] of Jordan and von Neumann concerning inner products, the von Neumann-Jordan (NJ-) constant $C_{\text{NJ}}(E)$ for a Banach space E was introduced by Clarkson [3] as the smallest constant C for which

$$(1) \quad \frac{1}{C} \leq \frac{\|x+y\|^2 + \|x-y\|^2}{2(\|x\|^2 + \|y\|^2)} \leq C$$

holds for all $x, y \in E$ with $(x, y) \neq (0, 0)$. As is easily seen, if C is best possible in the right-hand side inequality of (1), then so is $1/C$ in the left. Despite its fundamental nature very little is known about the NJ-constant; a summary is given in Theorem A below. In what follows, let $1 \leq p, q, r, t \leq \infty$ unless otherwise stated, and let p', q', r', t' be their conjugate numbers, respectively.

A. Theorem. (i) $1 \leq C_{\text{NJ}}(E) \leq 2$ for all Banach spaces E ; and $C_{\text{NJ}}(E) = 1$ if and only if E is a Hilbert space (Jordan and von Neumann [8]).

(ii) $C_{\text{NJ}}(L_p) = 2^{2/t-1}$, where $t = \min\{p, p'\}$ (Clarkson [3]; see also [5]).

(iii) For $L_p(L_q)$, L_q -valued L_p -space (on arbitrary measure spaces), $C_{\text{NJ}}(L_p(L_q)) = 2^{2/t-1}$, where $t = \min\{p, q, p', q'\}$; and for the Sobolev space $W_p^k(\Omega)$, $C_{\text{NJ}}(W_p^k(\Omega)) = 2^{2/t-1}$, where $t = \min\{p, p'\}$ (Kato and Miyazaki [10]).

Received by the editors September 8, 1995.

1991 Mathematics Subject Classification. Primary 46B20, 46B03, 46B42.

Key words and phrases. von Neumann-Jordan constant, uniform convexity, super-reflexivity, type and cotype, finite representability, p -convexity and p -concavity for a Banach lattice.

The authors were supported in part by Grants-in-Aid for Scientific Research from the Ministry of Education, Science and Culture (07640225 (first author), 07640240 (second author)).

(iv) For $E = C_c(K)$, resp. $C_b(K)$ (the spaces of continuous functions on a locally compact Hausdorff space K which have compact support, resp. are bounded), $C_{NJ}(E) = 2$ (Kato and Miyazaki [9]).

The following facts are readily seen:

B. Proposition. (i) $C_{NJ}(E) = 2^{2/t-1}$, $1 \leq t \leq 2$, if and only if

$$\|A: l_2^2(E) \rightarrow l_2^2(E)\| = 2^{1/t},$$

where $A = \begin{pmatrix} 1 & -1 \\ 1 & -1 \end{pmatrix}$ and $l_2^2(E)$ denotes the E -valued l_2^2 -space; and hence

(ii) $C_{NJ}(E') = C_{NJ}(E)$, where E' is the dual space of E . (This was observed for L_p in Clarkson [3].)

In this paper we first show that a uniformly convex Banach space E is nearly characterized by the condition $C_{NJ}(E) < 2$; more precisely the uniform convexity for E implies $C_{NJ}(E) < 2$, while conversely $C_{NJ}(E) < 2$ assures the existence of an equivalent uniformly convex norm in E . Hence, letting $\tilde{C}_{NJ}(E)$ be the infimum of all NJ-constants for equivalent norms of E , E is super-reflexive if and only if $\tilde{C}_{NJ}(E) < 2$. Secondly we show that if $\tilde{C}_{NJ}(E) = 2^{2/p-1}$, $1 < p \leq 2$ (the same value as that of L_p and of $L_{p'}$ as well), then E is of type r and cotype r' for any r with $1 \leq r < p$; and the converse holds if E is a Banach lattice and l_p is finitely representable in E or E' . Thus, for a Banach lattice E the number p satisfying $\tilde{C}_{NJ}(E) = 2^{2/p-1}$ is coincident with $\min\{p(E), q(E)'\}$, where $p(E) = \sup\{p; E \text{ is of type } p\}$, $q(E) = \inf\{q; E \text{ is of cotype } q\}$ and $1/q(E) + 1/q(E)' = 1$.

Let us recall some definitions. A Banach space E is called *strictly convex* if $\|(x+y)/2\| < 1$ whenever $\|x\| = \|y\| = 1$, $x \neq y$. E is said to be *uniformly convex* provided for each ε ($0 < \varepsilon < 2$) there exists a $\delta > 0$ such that $\|(x+y)/2\| < 1 - \delta$ whenever $\|x - y\| \geq \varepsilon$, $\|x\| = \|y\| = 1$. E is called $(2, \varepsilon)$ -convex, $\varepsilon > 0$, if $\min\{\|x+y\|, \|x-y\|\} \leq 2(1 - \varepsilon)$ whenever $\|x\| = \|y\| = 1$.

A Banach space F is said to be *finitely representable in E* provided for any $\lambda > 1$ and each finite-dimensional subspace F_1 of F , there is an isomorphism T of F_1 into E for which

$$\lambda^{-1}\|x\| \leq \|Tx\| \leq \lambda\|x\| \quad \text{for all } x \in F_1.$$

E is said to be *super-reflexive* ([7]; cf. [1], [12]) if no non-reflexive Banach space is finitely representable in E .

Super-reflexive spaces are characterized as those uniformly convexifiable:

C. Theorem (Enflo [4]; cf. [1], [15]). *A Banach space E is super-reflexive if and only if E admits an equivalent uniformly convex norm.*

A Banach space E is said to be of (Rademacher) *type p* ($1 \leq p \leq 2$), resp. of *cotype q* ($2 \leq q \leq \infty$), if there exists some $M > 0$ such that

$$\int_0^1 \left\| \sum_{j=1}^n r_j(t)x_j \right\| dt \leq M \left\{ \sum_{j=1}^n \|x_j\|^p \right\}^{1/p},$$

resp.

$$\left\{ \sum_{j=1}^n \|x_j\|^q \right\}^{1/q} \leq M \int_0^1 \left\| \sum_{j=1}^n r_j(t)x_j \right\| dt,$$

for all finite systems $\{x_j\}$ in E , where $r_j(t)$ are the Rademacher functions, i.e., $r_j(t) = \text{sgn}(\sin 2^j \pi t)$.

D. Theorem (Maurey and Pisier [14]; also [16], esp. Theorem 3.11, or [12],v. 3).
 Let E be a Banach space of infinite dimension. Let

$$p(E) := \sup\{p; E \text{ is of type } p\},$$

resp.

$$q(E) := \inf\{q; E \text{ is of cotype } q\}.$$

Then

$$p(E) = \min\{p; l_p \text{ is finitely representable in } E\},$$

resp.

$$q(E) = \max\{q; l_q \text{ is finitely representable in } E\}.$$

(Note that $p(E) \leq 2 \leq q(E)$.)

A Banach lattice E is called p -convex, resp. p -concave, $1 \leq p < \infty$, if there exists a constant M such that

$$(2) \quad \left\| \left\{ \sum_{j=1}^n |x_j|^p \right\}^{1/p} \right\| \leq M \left\{ \sum_{j=1}^n \|x_j\|^p \right\}^{1/p},$$

resp.

$$(3) \quad \left\{ \sum_{j=1}^n \|x_j\|^p \right\}^{1/p} \leq M \left\| \left\{ \sum_{j=1}^n |x_j|^p \right\}^{1/p} \right\|,$$

for all finite systems $\{x_j\}$ in E . The smallest possible value of M in (2), resp. (3), is called the p -convexity, resp. p -concavity, constant for E .

Now, we begin with the following proposition which will give effective examples later.

1. Proposition. Let $1 < p \leq 2$ and $\lambda > 1$. Let $X_{p,\lambda}$ be the space $l_{p'}$ equipped with the norm $\|x\|_{p,\lambda} := \max\{\|x\|_{p'}, \lambda\|x\|_\infty\}$, where $1/p + 1/p' = 1$. Then:

$$(i) \quad C_{\text{NJ}}(X_{p,\lambda}) = \begin{cases} \lambda^2 2^{2/p-1} & \text{if } 1 < \lambda \leq 2^{1/p'}, \\ 2 & \text{if } \lambda \geq 2^{1/p'}, \end{cases}$$

(ii) $X_{2,\lambda}$ is isomorphic to a Hilbert space and

$$C_{\text{NJ}}(X_{2,\lambda}) = \min\{\lambda^2, 2\},$$

(iii) $X_{p,\lambda}$ is not strictly convex for any $\lambda > 1$.

Proof. (i) Since

$$(4) \quad \|x\|_{p'} \leq \|x\|_{p,\lambda} \leq \lambda \|x\|_{p'} \quad \text{for all } x \in X_{p,\lambda},$$

we have

$$\begin{aligned} \|x + y\|_{p,\lambda}^2 + \|x - y\|_{p,\lambda}^2 &\leq \lambda^2 (\|x + y\|_{p'}^2 + \|x - y\|_{p'}^2) \\ &\leq \lambda^2 2^{2/p} (\|x\|_{p'}^2 + \|y\|_{p'}^2) \\ &\leq \lambda^2 2^{2/p} (\|x\|_{p,\lambda}^2 + \|y\|_{p,\lambda}^2), \end{aligned}$$

where the second inequality follows from the fact $C_{NJ}(l_{p'}) = 2^{2/p-1}$ (Theorem A (ii)) and Proposition B (i). This implies $C_{NJ}(X_{p,\lambda}) \leq \lambda^2 2^{2/p-1}$. Let $1 < \lambda \leq 2^{1/p'}$. Put $x = (1/\lambda, 1/\lambda, 0, \dots)$, $y = (1/\lambda, -1/\lambda, 0, \dots) \in X_{p,\lambda}$. Then, since $\|x\|_{p'} = \|y\|_{p'} = \lambda^{-1} 2^{1/p'} \geq 1$ and $\|x\|_\infty = \|y\|_\infty = \lambda^{-1}$, we have

$$\|x\|_{p,\lambda} = \|y\|_{p,\lambda} = \lambda^{-1} 2^{1/p'},$$

and clearly

$$(5) \quad \|x + y\|_{p,\lambda} = \|x - y\|_{p,\lambda} = 2.$$

Hence,

$$(6) \quad \frac{\|x + y\|_{p,\lambda}^2 + \|x - y\|_{p,\lambda}^2}{2(\|x\|_{p,\lambda}^2 + \|y\|_{p,\lambda}^2)} = \lambda^2 2^{2/p-1},$$

which implies $C_{NJ}(X_{p,\lambda}) = \lambda^2 2^{2/p-1}$. Let $\lambda \geq 2^{1/p'}$, and let x and y be as above. Then we have $\|x\|_{p,\lambda} = \|y\|_{p,\lambda} = 1$ and (5). Therefore the left side of (6), in this case, takes the value two, or $C_{NJ}(X_{p,\lambda}) = 2$.

(ii) is clear by (4) and (i).

(iii) Take $\alpha = (1/\lambda)^{p'} + \alpha^{p'} \leq 1$ and $0 < \alpha \leq 1/\lambda$. Put $x = (1/\lambda, 0, 0, \dots)$ and $y = (1/\lambda, \alpha, 0, \dots)$. Then $\|x\|_{p,\lambda} = \|y\|_{p,\lambda} = 1$, whereas $\|x + y\|_{p,\lambda} = 2$, so X is not strictly convex. This completes the proof. \square

2. Theorem. *Let E be uniformly convex. Then $C_{NJ}(E) < 2$.*

Proof. Let E be uniformly convex. Let ε be any positive number with $0 < \varepsilon < 2^{1/2}$. Then there exists a $\delta > 0$ such that $\|x\| \leq 1$, $\|y\| \leq 1$, and $\|x - y\| \geq \varepsilon$ imply $\|(x + y)/2\|^2 \leq (1 - \delta)[(\|x\|^2 + \|y\|^2)/2]$ (see [1], p. 190, or [11], p. 360). Let x and y be any elements in E with $\|x\|^2 + \|y\|^2 = 1$. We first assume that $\|x - y\| \geq \varepsilon$. Then

$$\left\| \frac{x + y}{2} \right\|^2 + \left\| \frac{x - y}{2} \right\|^2 \leq (1 - \delta) \frac{\|x\|^2 + \|y\|^2}{2} + \frac{\|x\|^2 + \|y\|^2}{2} \leq 1 - \frac{\delta}{2},$$

and hence

$$(7) \quad \|x + y\|^2 + \|x - y\|^2 \leq 2(2 - \delta).$$

Next, let $\|x - y\| \leq \varepsilon$. Then

$$(8) \quad \|x + y\|^2 + \|x - y\|^2 \leq 2(\|x\|^2 + \|y\|^2) + \varepsilon^2 \leq 2(1 + \varepsilon^2/2).$$

Consequently, by (7) and (8) we have

$$\frac{\|x + y\|^2 + \|x - y\|^2}{2(\|x\|^2 + \|y\|^2)} \leq 1 + \max\{1 - \delta, \varepsilon^2/2\},$$

or $C_{NJ}(E) < 2$. \square

The converse of Theorem 2 is not true:

3. Theorem. *For any $\varepsilon > 0$ there exists a Banach space E , isomorphic to a Hilbert space, with $C_{NJ}(E) < 1 + \varepsilon$ which is not strictly convex.*

In fact, for the spaces $X_{2,\lambda}$ ($\lambda > 1$), we have $C_{NJ}(X_{2,\lambda}) \rightarrow 1$ as $\lambda \rightarrow 1$, whereas $X_{2,\lambda}$ is not strictly convex.

Although the condition $C_{NJ}(E) < 2$ does not even imply strict convexity for E , it assures the existence of an equivalent norm on E for which E becomes uniformly convex (cf. Theorem C).

4. Theorem. *Let $C_{\text{NJ}}(E) < 2$. Then E is super-reflexive; the converse is not true.*

Proof. Assume $C := C_{\text{NJ}}(E) < 2$. We see that E is $(2, \varepsilon)$ -convex with some ε , which implies the super-reflexivity for E (James [6]; see also [1], [12]). Let x and y be any elements in E with $\|x\| = \|y\| = 1$. Then

$$\begin{aligned} \min_{\varepsilon_i = \pm 1} \|\varepsilon_1 x + \varepsilon_2 y\| &\leq \left\{ \frac{1}{2} (\|x + y\|^2 + \|x - y\|^2) \right\}^{1/2} \\ &\leq C^{1/2} (\|x\|^2 + \|y\|^2)^{1/2} \\ &= (2C)^{1/2} = 2(1 - \varepsilon), \end{aligned}$$

where $\varepsilon = 1 - (C/2)^{1/2}$. For the latter assertion consider the space $X_{2, \sqrt{2}}$. Indeed, $X_{2, \sqrt{2}}$ is isomorphic to a Hilbert space and hence super-reflexive, whereas $C_{\text{NJ}}(X_{2, \sqrt{2}}) = 2$ by Proposition 1. (l_1^n and l_∞^n are also such examples.) \square

5. Definition. Let $\tilde{C}_{\text{NJ}}(E)$ denote the infimum of all von Neumann-Jordan constants for equivalent norms of a Banach space E .

Owing to Theorems 2 and 4, we obtain the following theorem.

6. Theorem. *A Banach space E is super-reflexive if and only if $\tilde{C}_{\text{NJ}}(E) < 2$.*

So far we have treated Banach spaces whose NJ-constant is less than two or can be so taken by renorming. Let us now consider the spaces with NJ-constant $2^{2/t-1}$, $1 < t \leq 2$, which is the same as that of L_p -space if $t = \min\{p, p'\}$. We first note the following facts; the proof is easy and is omitted.

7. Proposition. (i) *Let F be a Banach space which is finitely representable in E . Then $C_{\text{NJ}}(F) \leq C_{\text{NJ}}(E)$.*

(ii) *If l_p is finitely representable in E , $C_{\text{NJ}}(E) \geq 2^{2/t-1}$, where $t = \min\{p, p'\}$.*

8. Proposition. *Let $C_{\text{NJ}}(E) \leq 2^{2/p-1}$, $1 < p \leq 2$. Then E is of type r and cotype r' for any r with $1 \leq r < p$. The converse is not true.*

Proof. Assume $C_{\text{NJ}}(E) \leq 2^{2/p-1}$. Suppose that E is not of type r for some $r < p$. Then l_r is finitely representable in E by Theorem D. Hence, $C_{\text{NJ}}(E) \geq 2^{2/r-1} > 2^{2/p-1}$ by the preceding proposition, which is a contradiction. By Proposition B, E' is also of type r for any $r < p$, which implies that E is of cotype r' for any such r (cf. [13], Proposition 1.e.17; or [1], p. 309).

The latter part is a consequence of a result by Pisier and Xu [17] which asserts that for any $q > 2$ there exists a non-reflexive Banach space E ($C_{\text{NJ}}(E) = 2$) such that E is of type 2 and cotype q . \square

9. Remark. Propositions 7 and 8 remain valid if $C_{\text{NJ}}(E)$ is replaced by $\tilde{C}_{\text{NJ}}(E)$ (note that “finite representability” and “type, cotype” are isomorphic properties).

For a Banach lattice the situation in Proposition 8 or Remark 9 is preferable, and we obtain the following result.

10. Theorem. *Let E be a Banach lattice and let $1 < p \leq 2$. Then the following assertions are equivalent:*

- (i) $\tilde{C}_{\text{NJ}}(E) \leq 2^{2/p-1}$,
- (ii) E is of type r and cotype r' for any $r < p$,

- (iii) E and E' are of type r for any $r < p$,
- (iv) E and E' are of cotype r' for any $r < p$,
- (v) E is r -convex and r' -concave for any $r < p$.

For the proof of Theorem 10 we need the following two lemmas.

11. Lemma ([13], Corollary 1.f.9). *Let $1 < r < \infty$. Let E be a Banach lattice of type r , resp. cotype r . Then E is $(r - \varepsilon)$ -convex, resp. $(r + \varepsilon)$ -concave, for any $\varepsilon > 0$.*

The following lemma is an easy consequence of Theorem 1.d.1 of [13] (see also the proof of Theorem 1.f.1 of [13]).

12. Lemma (cf. [13], Theorem 1.d.1). *Let $1 \leq p \leq 2$. Let E be a Banach lattice. Then, for any x and y in E ,*

$$(|x + y|^{p'} + |x - y|^{p'})^{1/p'} \leq 2^{1/p'} (|x|^p + |y|^p)^{1/p}.$$

Proof of Theorem 10. We have already seen that the implications (i) \Rightarrow (iii) \Rightarrow (ii) are valid in the proof of Proposition 8 and Remark 9. The implication (ii) \Rightarrow (iv) is clear, since if E is of type r , then E' is of cotype r' (cf. [13], Proposition 1.e.17; or [1], p. 309).

(iv) \Rightarrow (v): Assume that E and E' are of cotype r' for any $r < p$. Then by Lemma 11, E and E' are r' -concave for any $r < p$, and the latter implies r -convexity of E (see [13], Proposition 1.d.4).

(v) \Rightarrow (i): Let E be r -convex and r' -concave for any $r < p$. Here we can take both r -convex and r' -concave constants to be one by a suitable renorming ([13], Proposition 1.d.8). We denote such an equivalent norm by $\|\cdot\|_1$. Then by Lemma 12 we have

$$\begin{aligned} (\|x + y\|_1^2 + \|x - y\|_1^2)^{1/2} &\leq 2^{1/2-1/r'} (\|x + y\|_1^{r'} + \|x - y\|_1^{r'})^{1/r'} \\ &\leq 2^{1/2-1/r'} \|(|x + y|^{r'} + |x - y|^{r'})^{1/r'}\|_1 \\ &\leq 2^{1/2} \|(|x|^r + |y|^r)^{1/r}\|_1 \\ &\leq 2^{1/2} (\|x\|_1^r + \|y\|_1^r)^{1/r} \\ &\leq 2^{1/r} (\|x\|_1^2 + \|y\|_1^2)^{1/2}, \end{aligned}$$

which implies $C_{\text{NJ}}((E, \|\cdot\|_1)) \leq 2^{2/r-1}$. Consequently $\tilde{C}_{\text{NJ}}(E) \leq 2^{2/p-1}$. This completes the proof. \square

13. Remark. In Theorem 10 the assertion (i) may be replaced by

$$(i') \quad \tilde{C}_{\text{NJ}}(E) = 2^{2/p-1}$$

if l_p is finitely representable in E or E' (cf. Remark 9).

By Theorem 10 we obtain the following result.

14. Corollary. *Let E be a Banach lattice, and let $\tilde{C}_{\text{NJ}}(E) = 2^{2/p_0-1}$, $1 \leq p_0 \leq 2$. Then*

$$\begin{aligned} p_0 &= \sup\{p : E \text{ is of type } p \text{ and cotype } p'\} \\ &= \sup\{p : E \text{ and } E' \text{ are of type } p\} \\ &= \sup\{p : E \text{ and } E' \text{ are of cotype } p'\} \\ &= \sup\{p : E \text{ is } p\text{-convex and } p'\text{-concave}\} \\ &= \sup\{p : E \text{ and } E' \text{ are } p\text{-convex}\} \\ &= \sup\{p : E \text{ and } E' \text{ are } p'\text{-concave}\}. \end{aligned}$$

The proof for the case $p_0 = 1$ goes in the same way as in the proof of Theorem 10. (Note Lemma 1.f.3 of [13] for the last three identities.)

15. Remark. Let E be a Banach lattice, and let $p(E)$, $q(E)$ be as in Theorem D. Then the number p_0 in Corollary 14 is also represented as

$$p_0 = \min\{p(E), q(E)'\} = 2\{1 + \log_2 \tilde{C}_{\text{NJ}}(E)\}^{-1},$$

where $1/q(E) + 1/q(E)' = 1$.

ACKNOWLEDGEMENT

The authors thank the referee for some helpful comments.

REFERENCES

1. B. Beauzamy, *Introduction to Banach spaces and their geometry*, 2nd Ed., North Holland, Amsterdam-New York-Oxford, 1985. MR **88f**:46021
2. J. A. Clarkson, *Uniformly convex spaces*, Trans. Amer. Math. Soc. **40** (1936), 396–414.
3. J. A. Clarkson, *The von Neumann-Jordan constant for the Lebesgue space*, Ann. of Math. **38** (1937), 114–115.
4. P. Enflo, *Banach spaces which can be given an equivalent uniformly convex norm*, Israel J. Math. **13** (1972), 281–288. MR **49**:1073
5. E. Hewitt and K. Stromberg, *Real and abstract analysis*, Springer, New York-Heidelberg-Berlin, 1965. MR **32**:5826
6. R. C. James, *Uniformly non-square Banach spaces*, Ann. of Math. **80** (1964), 542–550. MR **30**:4139
7. R. C. James, *Super-reflexive Banach spaces*, Canad. J. Math. **24** (1972), 896–904. MR **47**:9248
8. P. Jordan and J. von Neumann, *On inner products in linear metric spaces*, Ann. of Math. **36** (1935), 719–723.
9. M. Kato and K. Miyazaki, *Remark on generalized Clarkson's inequalities for extreme cases*, Bull. Kyushu Inst. Tech., Math. Natur. Sci. **41** (1994), 27–31. MR **95h**:46044
10. M. Kato and K. Miyazaki, *On generalized Clarkson's inequalities for $L_p(L_q)$ and Sobolev spaces*, Math. Japon. **43** (1996), 505–515. CMP 96:13
11. G. Köthe, *Topologische lineare Räume I*, Springer, Berlin-Heidelberg-New York, 1966. MR **33**:3069
12. J. Kuelbs, *Probability on Banach spaces*, Marcel Dekker, New York-Basel, 1978. MR **80c**:60007
13. J. Lindenstrauss and L. Tzafriri, *Classical Banach spaces II*, Springer, Berlin-Heidelberg-New York, 1979. MR **81c**:46001
14. B. Maurey and G. Pisier, *Séries de variables aleatoires vectorielles independantes et proprietes geometriques des espaces de Banach*, Studia Math. **58** (1976), 45–90. MR **56**:1388
15. G. Pisier, *Martingales with values in uniformly convex spaces*, Israel J. Math. **20** (1975), 326–350. MR **52**:14940

16. G. Pisier, *Factorization of linear operators and geometry of Banach spaces*, Amer. Math. Soc., Providence, RI, 1986. MR **88a**:47020
17. G. Pisier and Q. Xu, *Random series in the real interpolation spaces between the spaces v_p* , Lecture Notes in Math., vol. 1267, Springer-Verlag, Berlin-Heidelberg-New York, 1987, pp. 185–209. MR **89d**:46011

DEPARTMENT OF MATHEMATICS, KYUSHU INSTITUTE OF TECHNOLOGY, TOBATA, KITAKYUSHU 804, JAPAN

DEPARTMENT OF SYSTEM ENGINEERING, OKAYAMA PREFECTURAL UNIVERSITY, SOJA 719-11, JAPAN