

## ITERATION OF COMPACT HOLOMORPHIC MAPS ON A HILBERT BALL

CHO-HO CHU AND PAULINE MELLON

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. Given a compact holomorphic fixed-point-free self-map,  $f$ , of the open unit ball of a Hilbert space, we show that the sequence of iterates,  $(f^n)$ , converges locally uniformly to a constant map  $\xi$  with  $\|\xi\| = 1$ . This extends results of Denjoy (1926), Wolff (1926), Hervé (1963) and MacCluer (1983). The result is false without the compactness assumption, nor is it true for all open balls of  $J^*$ -algebras.

### 1. INTRODUCTION

There has been extensive literature on the subject of iterating holomorphic functions since the early works of Julia [14], Fatou [6], [7], Denjoy [3] and Wolff [23], [24]. We refer to [2], [20] for some interesting surveys and references.

Given a fixed-point-free holomorphic map  $f: \Delta \rightarrow \Delta$  where  $\Delta = \{z \in \mathbb{C} : |z| < 1\}$ , Wolff's theorem [24] states that there is a boundary point  $u \in \partial\Delta$  such that every closed disc internally tangent to  $\Delta$  at  $u$  is invariant under the iterates of  $f$ . From this follows the result of Denjoy [3] and Wolff [23] that the iterates,  $f^n = \underbrace{f \circ \cdots \circ f}_n$ ,

of  $f$  converge to  $u$  uniformly on compact subsets of  $\Delta$ . Wolff's theorem has been extended to Hilbert balls [8], and the convergence result of Denjoy and Wolff also extends to the open unit ball of  $\mathbb{C}^n$  [13], [15], as well as some other domains in  $\mathbb{C}^n$  [1]. Nevertheless, the convergence result fails for infinite dimensional Hilbert balls and Stachura [18] has given an example to show that it fails even for *biholomorphic* self-maps.

Recently, Wolff-type theorems have been established for *compact* holomorphic self-maps of the open unit balls of  $J^*$ -algebras (which include  $C^*$ -algebras and Hilbert spaces) [6], [25]. A natural question is whether a Denjoy-Wolff-type convergence result for *compact* holomorphic maps on  $J^*$ -algebras might also follow from these Wolff-type theorems. We show that this is the case for Hilbert spaces, but not the case even for finite-dimensional  $C^*$ -algebras. We prove the following result.

**Theorem.** *Let  $H$  be a Hilbert space with open unit ball  $B$ . Let  $f: B \rightarrow B$  be a compact holomorphic map with no fixed point in  $B$ . Then there exists  $\xi \in \partial B$  such that the sequence  $(f^n)$  of iterates of  $f$  converges locally uniformly on  $B$  to the constant map taking value  $\xi$ .*

---

Received by the editors December 27, 1995.

1991 *Mathematics Subject Classification.* Primary 46G20, 32A10, 32A17; Secondary 32M15.

©1997 American Mathematical Society

We will give a simple example to show that the above result is false if  $H$  is replaced by a  $C^*$ -algebra. We also note that it has been shown in [10] that if  $f: B \rightarrow B$  is fixed-point-free and so-called *firmly holomorphic*, then the iterates  $(f^n)$  converge *pointwise* to a boundary point  $\xi \in \partial B$ .

It may be useful to recall the Earle-Hamilton Theorem [5] which states that every holomorphic map  $f: B \rightarrow B$ , where  $B$  is a bounded domain in a Banach space, has a fixed-point if  $f(B)$  is *strictly* contained in  $B$ .

## 2. PRELIMINARIES

All Banach spaces will be complex. Given *bounded* domains  $D$  and  $D'$  in any Banach spaces, we denote by  $H(D, D')$  the space of all holomorphic maps  $f: D \rightarrow D'$ . We write  $H(D)$  for  $H(D, D)$ . Every nonempty open ball  $B$  in  $D$  induces a norm  $\|\cdot\|_B$  on  $H(D, D')$  where  $\|f\|_B = \sup_{x \in B} \|f(x)\|$  for  $f \in H(D, D')$ . The *topology of local uniform convergence* on  $H(D, D')$  is the topology induced by the norms  $\|\cdot\|_B$  where  $B$  is an open ball in  $D$  satisfying  $\text{dist}(B, \partial D) > 0$ ,  $\partial D$  being the boundary of  $D$ . Using Hadamard's three circles theorem, it has been shown in [21], [22] (see also [19, Lemma 13.1]) that  $\|\cdot\|_{B_1}$  and  $\|\cdot\|_{B_2}$  induce the same topology for any open balls  $B_1, B_2$  in  $D$  satisfying  $\text{dist}(B_1, \partial D) > 0$  and  $\text{dist}(B_2, \partial D) > 0$ . It follows that a sequence  $(f_n)$  in  $H(D, \overline{D})$  converges to  $f \in H(D, \overline{D})$  locally uniformly if, and only if, for every  $x \in D$ ,  $(f_n)$  converges uniformly to  $f$  on *some* open ball  $B$  containing  $x$  and satisfying  $\text{dist}(B, \partial D) > 0$ . Given any  $x$  in a Banach space  $X$ , and  $r > 0$ , we let  $B(x, r) = \{y \in X : \|y - x\| < r\}$ . A map  $f: D \rightarrow D' \subset X$  is called *compact* if the closure  $\overline{f(D)}$  is compact in  $X$ .

**Lemma 1.** *Let  $B$  be the open unit ball of a Banach space  $X$  and let  $f: B \rightarrow B$  be a compact holomorphic map. Then the sequence  $(f^n)$  of iterates of  $f$  has a subsequence converging locally uniformly to a function in  $H(B, \overline{B})$ .*

*Proof.* Choose a sequence  $(r_n)$  in  $(0, 1)$  such that  $r_n \uparrow 1$  and  $f(B) \cap B(0, r_1) \neq \emptyset$ . We have  $f(B) = \bigcup_{n=1}^{\infty} (f(B) \cap B(0, r_n))$ . We first find a subsequence of  $(f^n)$  converging uniformly on  $f(B) \cap B(0, r_1)$ . By compactness of  $\overline{f(B) \cap B(0, r_1)} \subset \overline{f(B)}$ , there is a countable set  $\{z_n\}$  in  $f(B) \cap B(0, r_1)$ , which is dense in  $\overline{f(B) \cap B(0, r_1)}$ .

Since  $f$  is compact,  $(f^n)$  has a subsequence,  $(f^{(n,1)})$ , such that  $(f^{(n,1)}(z_1))$  converges. Likewise,  $(f^{(n,1)})$  has a subsequence  $(f^{(n,2)})$  such that  $(f^{(n,2)}(z_2))$  converges. Proceed to find subsequences  $(f^{(n,k)})_n$  which converge at  $z_1, \dots, z_k$ . We show that the diagonal sequence  $(f^{(k,k)})$  converges uniformly on  $f(B) \cap B(0, r_1)$ . It suffices to show that it is uniformly Cauchy on  $f(B) \cap B(0, r_1)$ . Let  $\varepsilon > 0$ . Since  $\text{dist}(B(0, r_1), \partial B) = 1 - r_1 > 0$ , we have

$$(1) \quad \|h(z) - h(w)\| \leq \frac{\|z - w\|}{1 - r_1}$$

for  $h \in H(B)$  and  $z, w \in B(0, r_1)$  (cf. [19, 1.17]). By compactness, there exist  $z_{n_1}, \dots, z_{n_l}$  in  $\{z_n\}$  such that

$$\overline{f(B) \cap B(0, r_1)} \subset \bigcup_{i=1}^l B(z_{n_i}, \frac{\varepsilon}{3}(1 - r_1)).$$

There exists  $N$  such that  $j, k > N$  implies

$$\|f^{(j,j)}(z_{n_i}) - f^{(k,k)}(z_{n_i})\| < \frac{\varepsilon}{3}$$

for  $i = 1, \dots, l$ . Hence, for any  $z \in f(B) \cap B(0, r_1)$ , we have  $z \in B(z_{n_i}, \frac{\varepsilon}{3}(1 - r_1))$  for some  $i$ , and

$$\begin{aligned} \|f^{(j,j)}(z) - f^{(k,k)}(z)\| &\leq \|f^{(j,j)}(z) - f^{(j,j)}(z_{n_i})\| + \|f^{(j,j)}(z_{n_i}) - f^{(k,k)}(z_{n_i})\| \\ &\quad + \|f^{(k,k)}(z_{n_i}) - f^{(k,k)}(z)\| \\ &< \frac{\varepsilon(1 - r_1)}{3(1 - r_1)} + \frac{\varepsilon}{3} + \frac{\varepsilon(1 - r_1)}{3(1 - r_1)} = \varepsilon \end{aligned}$$

whenever  $j, k > N$ . This shows that  $(f^{(k,k)})$  is uniformly convergent on  $f(B) \cap B(0, r_1)$ .

We repeat the diagonal process as follows. Choose a subsequence  $(f^{n_1})$  of  $(f^n)$  converging uniformly on  $f(B) \cap B(0, r_1)$ . Then choose a subsequence  $(f^{n_2})$  of  $(f^{n_1})$  converging uniformly on  $f(B) \cap B(0, r_2)$ , and so on. The diagonal sequence  $(f^{n_n})$  then converges uniformly on  $f(B) \cap B(0, r_k)$  for  $k = 1, 2, \dots$

Finally, we show that  $(f^{n_n+1})$  converges locally uniformly on  $B$ . Pick  $x \in B$  and choose  $r, R > 0$  such that  $r + R = 1 - \|x\|$  and  $\frac{r}{R} < 1 - \|f(x)\|$ . Then  $B(x, r)$  and  $B(f(x), \frac{r}{R})$  are contained in  $B$ . As in (1),  $\text{dist}(B(x, r), \partial B) \geq R > 0$  implies

$$f(B(x, r)) \subset B(f(x), \frac{r}{R}) \cap f(B) \subset B(0, r_k) \cap f(B)$$

for some  $k$ . It follows that  $(f^{n_n})$  converges uniformly on  $f(B(x, r))$  and hence  $(f^{n_n+1})$  converges uniformly on  $B(x, r)$ .  $\square$

*Remark 1.* The above proof implies that every subsequence  $(f^{n_k})$  of the iterates  $(f^n)$  has a locally uniformly convergent subsequence.

We need the following version of the maximum modulus principle and we include a proof for completeness (cf. [4, p.95]).

**Lemma 2.** *Let  $D$  be a domain in a Banach space  $X$  and let  $B$  be the open unit ball of a Hilbert space  $H$ . Given any holomorphic function  $f: D \rightarrow \overline{B}$ , we have either  $f(D) \subset B$  or  $f(z) = \xi \in \partial B$  for all  $z \in D$ .*

*Proof.* Suppose  $f(z_0) = \xi \in \partial B$  for some  $z_0 \in D$  where  $D$  contains some open ball  $B(z_0, r)$  with  $r > 0$ . We show that  $f(z) = \xi$  for all  $z \in D$ . It suffices to show  $f(v) = \xi$  for all  $v \in B(z_0, r)$ . Fix  $v$  arbitrary in  $B(z_0, r)$ . Define  $\varphi_v: \Delta \rightarrow \mathbb{C}$  by

$$\varphi_v(\lambda) = \langle f(z_0 + \lambda(v - z_0)), \xi \rangle \quad (\lambda \in \Delta)$$

where  $\langle \cdot, \cdot \rangle$  denotes the inner product on  $H$ . Then  $\varphi_v: \Delta \rightarrow \overline{\Delta}$  and  $\varphi_v(0) = 1$  imply  $\varphi_v \equiv 1$  by the maximum modulus principle. It follows that  $f(z_0 + \lambda(v - z_0)) = \xi$  for all  $\lambda \in \Delta$  which gives  $f(v) = \xi$  by continuity.  $\square$

### 3. DENJOY-WOLFF-TYPE RESULT

In this section, we prove the **Theorem** and give some simple examples. Wolff's theorem has been extended to fixed-point-free holomorphic self-maps  $f$  of a Hilbert ball  $B$ , in which case there exists  $\xi \in \partial B$  such that the "ellipsoids"

$$E(\xi, \lambda) = \left\{ x \in B : \frac{|1 - \langle x, \xi \rangle|^2}{1 - \|x\|^2} < \lambda \right\} \quad (\lambda > 0)$$

are invariant under  $f$ , and further,  $\overline{E(\xi, \lambda)} \cap \partial B = \{\xi\}$  (cf. [8]). If  $\dim B < \infty$ , then the iterates  $f^n$  must converge locally uniformly to the "Wolff point"  $\xi$  (cf. [13, 15]).

As remarked before, the latter result is false in infinite dimensions by Stachura’s example [18].

Let  $\mathcal{L}(H, K)$  be the Banach space of bounded linear operators between Hilbert spaces  $H$  and  $K$ . A closed linear subspace  $Z \subset \mathcal{L}(H, K)$  is called a  $J^*$ -algebra if  $TT^*T \in Z$  whenever  $T \in Z$ , where  $T^*$  denotes the adjoint of  $T$  (cf. [11, 19]). Every Hilbert space  $H = \mathcal{L}(\mathbb{C}, H)$  is a  $J^*$ -algebra, and so is every  $C^*$ -algebra.

Let  $B$  be the open unit ball of a  $J^*$ -algebra  $Z$  and let  $f: B \rightarrow B$  be a fixed-point-free compact holomorphic map. A Wolff-type result has been obtained in [16, Theorem 5] which states that under certain conditions on  $f$ , there exist a “Wolff point”  $\xi \in \partial B$  and circular domains  $D_{z,\xi}$  ( $z \in B$ ) invariant under  $f$  (see also [25]). The question of whether the iterates  $f^n$  would converge to  $\xi$  was unanswered in [16]. The following example gives a negative answer.

**Example 1.** Let  $Z = \mathbb{C} \times \mathbb{C}$  be equipped with the coordinatewise product and norm  $\|(z, w)\| = \max(|z|, |w|)$ . Then  $Z$  is a  $C^*$ -algebra with open unit ball  $\Delta \times \Delta$ . Pick any fixed-point-free  $h \in H(\Delta)$ . Define  $f: \Delta \times \Delta \rightarrow \Delta \times \Delta$  by

$$f(z, w) = (iz, h(w)) \quad (z, w \in \Delta).$$

Then  $f$  is fixed-point-free and we have

$$f^n(z, w) = (i^n z, h^n(w)),$$

where  $(h^n)$  converges locally uniformly on  $\Delta$  to some  $\xi \in \partial\Delta$ . The iterates  $(f^n)$  clearly do not converge to any boundary point in  $\partial(\Delta \times \Delta)$ .

Nevertheless, we can still derive a Denjoy-Wolff-type convergence result for compact holomorphic maps on Hilbert spaces, by adapting MacCluer’s arguments for  $\mathbb{C}^n$  in [15]. A crucial step in the proof depends on the fact that the automorphisms of a Hilbert ball map affine sets to affine sets, and consequently that the fixed-point set of a nonconstant holomorphic map is affine. In contrast, the automorphisms of the open unit ball of an arbitrary  $J^*$ -algebra may distort the affine sets and the fixed-point set of a nonconstant holomorphic map need *not* be affine, even in the simple case of the bidisc as shown by the example below. This is one reason why a Denjoy-Wolff-type result fails for arbitrary  $J^*$ -algebras.

Let  $B$  be the open unit ball of a Banach space  $X$ . By an *affine subset* of  $B$  we mean a nonempty set of the form  $(c + L) \cap B$  where  $c \in X$  and  $L$  is a closed linear subspace of  $X$ . If  $X$  is a Hilbert space,  $c$  can be chosen to be orthogonal to  $L$ , and also, for nonconstant  $h \in H(B)$ , its fixed-point set  $\text{Fix}(h) = \{x \in B : h(x) = x\}$  is affine by [12] (see also [9, Theorem 23.2]). This was proved in [17] in finite dimensions.

**Example 2.** Let  $Z = \mathbb{C} \times \mathbb{C}$  be as in Example 1, with open unit ball  $\Delta \times \Delta$ . Fix  $a \in \Delta \setminus \{0\}$ . Define  $h: \Delta \times \Delta \rightarrow \Delta \times \Delta$  by

$$h(z, w) = (g_a(w), g_{-a}(z)) \quad (z, w \in \Delta)$$

where  $g_a(w) = \frac{a + w}{1 + \bar{a}w}$ . Then  $\text{Fix}(h) = \{(z, g_{-a}(z)) : z \in \Delta\}$  which is *not* affine since  $(0, -a)$  and  $(a, 0)$  are in  $\text{Fix}(h)$  while  $\frac{1}{2}(0, -a) + \frac{1}{2}(a, 0) \notin \text{Fix}(h)$ . We also note that  $h^{2n}(z, w) = (z, w)$  and  $h^{2n+1}(z, w) = h(z, w)$ . So  $(h^n(z, w))_n$  does not converge if  $(z, w) \notin \text{Fix}(h)$ .

We are now ready to prove the Theorem.

*Proof of the Theorem.* We will use the same symbol throughout for both a constant function and its value.

Let  $\xi \in \partial B$  be the “Wolff point” of  $f$  as mentioned in the beginning of this section. Let  $\Gamma(f)$  be the set of all subsequential limits of  $\{f^n : n = 1, 2, \dots\}$  in  $H(B, \overline{B})$  with respect to the topology of local uniform convergence. By Lemma 1,  $\Gamma(f) \neq \emptyset$ .

We first show that  $\Gamma(f)$  consists of constant maps only. Suppose, otherwise, that  $\Gamma(f)$  contains a nonconstant map  $g \in H(B, \overline{B})$ . We deduce a contradiction. By Lemma 2,  $g(B) \subset B$ . Let  $(f^{n_k})$  be a subsequence of  $(f^n)$  converging to  $g$ . Let  $m_k = n_{k+1} - n_k$ . By Remark 1,  $(f^{m_k})$  has a convergent subsequence, and we may assume, without loss of generality, that  $f^{m_k} \rightarrow h_0 \in H(B, \overline{B})$ . Since  $f^{n_{k+1}} = f^{m_k} \circ f^{n_k} \rightarrow h_0 \circ g$ , we have  $h_0 \circ g = g$  and  $h_0$  is the identity on  $g(B)$ . So  $h_0$  is nonconstant,  $h_0(B) \subset B$  and  $A_0 = \text{Fix}(h_0)$  is an affine subset of  $B$ . Since  $A_0 \subset \overline{f(B)}$  which is compact, it follows that  $\dim A_0 < \infty$ . Clearly,  $A_0 \subset h_0(B)$ . If  $A_0 \neq h_0(B)$ , then we repeat the above process. Letting  $p_k = m_{k+1} - m_k$ , we may assume  $f^{p_k} \rightarrow h_1 \in H(B)$  satisfying  $h_1 \circ h_0 = h_0$ . So  $h_1$  is the identity on  $h_0(B)$  and  $h_0(B) \subset A_1 = \text{Fix}(h_1)$ . We have  $A_0 \subsetneq A_1$  and  $A_1$  is a finite dimensional affine subset of  $B$ , with  $\dim A_1 > \dim A_0$ . If  $A_1 \neq h_1(B)$ , we repeat the process again. Continuing in this manner, we must eventually find some  $h_i \in H(B)$  such that  $h_i(B) = A_i = \text{Fix}(h_i)$ . For otherwise, we can construct a sequence  $(v_j)_j \subset \cup_{i=1}^{\infty} A_i$  with the property that  $\|v_i - v_j\| > \delta$  for all  $i \neq j$  and some  $\delta > 0$ . Since  $\cup_{i=1}^{\infty} A_i \subset \overline{f(B)}$  which is compact, this is clearly impossible. It follows that  $h_i^2 = h_i$ . Let  $(f^{l_k})$  converge to  $h_i$  locally uniformly. Note that  $f(A_i) \subset A_i$ . Since  $A_i$  is a finite dimensional affine subset of  $B$  and the automorphisms act transitively on  $B$ , a similar argument to that in [15, p. 98] shows that  $A_i$  is biholomorphically equivalent to the open unit ball of  $\mathbb{C}^n$  where  $n = \dim A_i$ . Now  $f|_{A_i} : A_i \rightarrow A_i$  is fixed-point-free and by [13, 15], there exists  $\mu \in \partial A_i$  such that  $(f|_{A_i})^n$  converges locally uniformly to  $\mu$  on  $A_i$ . So  $h_i|_{A_i} = \lim_{k \rightarrow \infty} (f|_{A_i})^{l_k} = \mu$  which is impossible as  $h_i$  is nonconstant and  $h_i(B) = A_i$ .

Therefore  $\Gamma(f)$  must consist of constant maps only. Now take any  $\eta \in \Gamma(f)$ . Then  $\eta \in \partial B$ , for otherwise  $\eta$  would be a fixed point of  $f$  in  $B$ . There is a subsequence  $(f^{n_k})$  converging to  $\eta$ . Let  $\lambda > 0$  and let  $z \in E(\xi, \lambda)$ . We have

$$\eta = \lim_{k \rightarrow \infty} f^{n_k}(z) \in \overline{E(\xi, \lambda)} \cap \partial B = \{\xi\}$$

since  $E(\xi, \lambda)$  is  $f$ -invariant. Therefore every convergent subsequence of  $(f^n)$  converges to the constant map  $\xi$ . It follows from Remark 1 that  $(f^n)$  must converge locally uniformly to  $\xi$  and the proof is complete.  $\square$

We end with the following example of a fixed-point-free compact holomorphic map on a Hilbert ball.

**Example 3.** Let  $B$  be the open unit ball of the (complex) Hilbert space  $l_2$ . Define  $f : B \rightarrow B$  by

$$\begin{aligned} f(x_1, x_2, \dots) &= \left( \frac{1+x_1}{2}, \left( \frac{1-x_1}{2} \right) \frac{x_1}{2}, \left( \frac{1-x_1}{2} \right) \frac{x_2}{3}, \dots \right) \\ &= \left( \frac{1+x_1}{2}, 0, 0, \dots \right) + \frac{1-x_1}{2} \left( 0, \frac{x_1}{2}, \frac{x_2}{3}, \dots \right) \end{aligned}$$

for  $(x_1, x_2, \dots) \in B$ . Then  $f$  is fixed-point-free and holomorphic. Moreover,  $f$  is compact since it is the sum of two compact maps. Also the *Wolff point* is  $(1, 0, 0, \dots)$ .

#### ACKNOWLEDGEMENTS

This research was carried out during the authors' visits to each other's institution. We gratefully acknowledge financial support from the British Council/Forbairt Joint Research Scheme and the Arts Faculty of University College Dublin.

#### REFERENCES

- [1] M. Abate, *Iteration theory, compactly divergent sequences and commuting holomorphic maps*, Ann. della Scuola Normale Superiore di Pisa **18** (1991) 67-191. MR **93e**:32034a
- [2] R. B. Burckel, *Iterating analytic self-maps of discs*, Amer. Math. Monthly **88** (1981) 396-407. MR **82g**:30046
- [3] A. Denjoy, *Sur l'itération des fonctions analytiques*, C. R. Acad. Sc. Paris **182** (1926) 255-257.
- [4] S. Dineen, "The Schwarz lemma", Oxford Univ. Press, Oxford, 1989. MR **91f**:46064
- [5] C. J. Earle and R. S. Hamilton, *A fixed point theorem for holomorphic mappings*, Proc. Symposia Pure Math **16** (1969) 61-65. MR **42**:918
- [6] M. P. Fatou, *Sur l'équations fonctionnelles*, Bull. Soc. Math. France **47** (1919) 161-271.
- [7] M. P. Fatou, *Sur l'équations fonctionnelles*, Bull. Soc. Math. France **48** (1920) 22-94 and 208-314.
- [8] K. Goebel, *Fixed points and invariant domains of holomorphic mappings of the Hilbert ball*, Nonlinear Analysis **6** (1982) 1327-1334. MR **84j**:58027
- [9] K. Goebel and S. Reich, "Uniform convexity, hyperbolic geometry and nonexpansive mappings", Marcel Dekker, 1984. MR **86d**:58012
- [10] K. Goebel and S. Reich, *Iterating holomorphic self-mappings of the Hilbert ball*, Proc. Japan. Acad. **58** (1982) 349-352. MR **84b**:47065
- [11] L. A. Harris, *A generalization of  $C^*$ -algebras*, Proc. London Math. Soc. **42** (1981) 331-361. MR **82e**:46089
- [12] T. L. Hayden and T. J. Suffridge, *Biholomorphic maps in Hilbert space have a fixed point*, Pacific J. Math. **38** (1971) 419-422. MR **46**:4288
- [13] M. Hervé, *Quelques propriétés des applications analytiques d'une boule à  $m$  dimensions dans elle-meme*, J. Math. Pures et Appl. **42** (1963) 117-147. MR **28**:3177
- [14] G. Julia, *Mémoire sur l'itération des fonctions rationnelles*, J. Math. Pures et Appl. **8** (1918) 47-245.
- [15] B. D. MacCluer, *Iterates of holomorphic self-maps of the open unit ball in  $\mathbb{C}^n$* , Michigan Math J. **30** (1983) 97-106. MR **85c**:32047a
- [16] P. Mellon, *Another look at results of Wolff and Julia type for  $J^*$ -algebras*, J. Math. Anal. Appl. **198** (1996) 444-457. CMP 96:08
- [17] W. Rudin, "Function theory in the unit ball of  $\mathbb{C}^n$ ", Springer, New York, 1980. MR **82i**:32002
- [18] A. Stachura, *Iterates of holomorphic self-maps of the unit ball in Hilbert space*, Proc. Amer. Math. Soc. **93** (1985) 88-90. MR **86b**:47117
- [19] H. Upmeyer, "Symmetric Banach manifolds and Jordan  $C^*$ -algebras", North-Holland, 1985. MR **87a**:58022
- [20] E. Vesentini, *Iteration of holomorphic maps*, Russ. Math. Survey **40** (1985) 7-11.
- [21] J. P. Vigué, *Sur le groupe des automorphismes analytiques d'un ouvert borné d'un espace de Banach complexe*, C. R. Acad. Sc. Paris **278** (1974) 617-620. MR **49**:3230
- [22] J. P. Vigué, *Le groupe des automorphismes analytiques d'un domaine borné d'un espace de Banach complexe, application aux domaines bornés symétriques*, Ann. Sc. Ec. Norm. Sup. **9** (1976) 203-282. MR **55**:3340
- [23] J. Wolff, *Sur l'itération des fonctions bornées*, C. R. Acad. Sc. Paris **182** (1926) 42-43 and 200-201.

- [24] J. Wolff, *Sur une généralisation d'un théorème de Schwarz*, C. R. Acad. Sc. Paris **182** (1926) 918-920 and 183 (1926) 500-502.
- [25] K. Włodarczyk, *Julia's lemma and Wolff's theorem for  $J^*$ -algebras*, Proc. Amer. Math. Soc. **99** (1987) 472-476. MR **88a**:46049

GOLDSMITHS COLLEGE, UNIVERSITY OF LONDON, LONDON SE14 6NW, ENGLAND  
*E-mail address*: `maa01chc@gold.ac.uk`

DEPARTMENT OF MATHEMATICS, UNIVERSITY COLLEGE DUBLIN, BELFIELD, DUBLIN 4, IRELAND  
*E-mail address*: `pmellon@irlearn.ucd.ie`