

TORI IN CERTAIN ASPHERICAL FOUR-MANIFOLDS

ALLAN L. EDMONDS

(Communicated by Ronald A. Fintushel)

ABSTRACT. The homology classes represented by embedded or mapped tori in the product of two surfaces are completely characterized.

If X^4 is a 4-manifold, then any element $\xi \in H_2(X^4)$ can be represented by a map from a closed, oriented surface, and also by an embedded, closed, oriented surface (perhaps of bigger genus). If X^4 is simply connected then $H_2(X^4) \approx \pi_2(X^4)$ by the Hurewicz Theorem, so that every class ξ is represented by a map $S^2 \rightarrow X^4$. It has been an intensively studied fundamental problem to determine which classes are represented by smooth or topologically embedded 2-spheres, and, more generally, to determine the minimum genus of an embedded surface representing ξ . In general there are striking differences between the smooth and topological categories. See, for example, [2] and [3] for some of the latest results in this area.

In this note we begin to address similar questions in the aspherical case, in which there are no interesting 2-spheres at all, apart from “local knot theory.”

We determine which 2-dimensional homology classes can be represented by embedded tori in $X^4 = F^2 \times G^2$, where F^2 and G^2 are closed orientable surfaces of genus greater than 1. It turns out that there is no difference between the topological and differentiable categories in this case.

Theorem 1. *Let F^2 and G^2 be closed orientable surfaces of genus greater than 1. An element $\xi \in H_2(F^2 \times G^2)$ is represented by a topological or differentiable embedded torus $T^2 \subset F^2 \times G^2$ if and only if $\xi = \alpha \times \beta$ for some $\alpha \in H_1(F^2)$ and $\beta \in H_1(G^2)$.*

This follows immediately from the following two more precise propositions.

Proposition 1. *Any map $f : T^2 \rightarrow F^2 \times G^2$ represents a homology class of the form $\alpha \times \beta$ for some $\alpha \in H_1(F^2)$ and $\beta \in H_1(G^2)$.*

Proposition 2. *If $\alpha \in H_1(F^2)$ and $\beta \in H_1(G^2)$ are given, then the homology class $\alpha \times \beta$ is represented by a smooth embedding $f : T^2 \rightarrow F^2 \times G^2$.*

Lemma 1. *If F^2 is a surface of negative euler characteristic, then any two commuting elements in $\pi_1(F^2)$ generate a cyclic subgroup of $\pi_1(F^2)$.*

Proof. Let $\alpha, \beta \in \pi_1(F^2)$ be commuting elements and let $S \rightarrow F^2$ be the covering map, where $\pi_1(S)$ corresponds to the subgroup of $\pi_1(F^2)$ generated by α and β . If it is a finite-sheeted covering, then $S \cong T^2$ by the classification of surfaces, and

Received by the editors October 7, 1996.

1991 *Mathematics Subject Classification.* Primary 57N13, 57N35.

hence $\chi(S) = 0$. But multiplicativity of the euler characteristic then implies that $\chi(F^2) = 0$ too, a contradiction. Therefore the covering must be an infinite-sheeted covering. In particular, S is non-compact, and hence has a free fundamental group. Since $\pi_1(S)$ is abelian, this implies that it is cyclic, as required. \square

Corollary 1. *If F^2 is a closed orientable surface of negative euler characteristic, and $f : T^2 \rightarrow F^2$ is a map, then there is a primitive homology class $\alpha \in H_1(F^2)$ such that the image of $f_* : H_1(T^2) \rightarrow H_1(F^2)$ is contained in the cyclic subgroup generated by α .*

Proof. Since $\pi_1(T^2) \approx H_1(T^2)$, the image of $H_1(T^2)$ in $H_1(F^2)$ is the image under the Hurewicz homomorphism of the image of $\pi_1(T^2)$ in $\pi_1(F^2)$, which is cyclic by Lemma 1, and hence contained in a cyclic subgroup generated by a primitive element. \square

We will need the following well-known result. For a proof, see [4].

Lemma 2. *If $\alpha \in H_1(F^2)$ is a nonzero homology class, then α is represented by a simple closed curve embedded in F^2 if and only if α is primitive.* \square

Proof of Proposition 1. Let $f : T^2 \rightarrow F^2 \times G^2$ represent a nonzero homology class and let $p_F : F^2 \times G^2 \rightarrow F^2$ and $p_G : F^2 \times G^2 \rightarrow G^2$ denote the projections onto the two factors. By Corollary 1 the image of $(p_F \circ f)_* : H_1(T^2) \rightarrow H_1(F^2)$ is contained in a cyclic subgroup $\langle \alpha \rangle$ generated by a primitive element $\alpha \in H_1(F^2)$. Similarly, the image of $(p_G \circ f)_* : H_1(T^2) \rightarrow H_1(G^2)$ is contained in a cyclic subgroup $\langle \beta \rangle$ generated by a primitive element $\beta \in H_1(G^2)$. Now by Lemma 2 both α and β are represented by simple closed curves A and B in F^2 and G^2 , respectively. It follows that $f_*[T^2]$ is a multiple of $[A \times B]$ in $H_2(F^2 \times G^2)$. But $n(\alpha \times \beta) = (n\alpha) \times \beta$, which is what is needed. \square

Proof of Proposition 2. If we write $\alpha = m\alpha^*$ and $\beta = n\beta^*$, where α^* and β^* are primitive homology classes, then $\alpha \times \beta = mn\alpha^* \times \beta^* = (mn\alpha^*) \times \beta^*$. Now, by Lemma 2, α^* and β^* can be represented by simple closed curves A^* and B^* , respectively. Since A^* and B^* have trivial normal bundles, so does the torus $A^* \times B^* \subset F^2 \times G^2$. The proof then is completed by Lemma 3 below. \square

Lemma 3. *The map $T^2 \rightarrow T^2 \times D^2$ given by $(z, w) \mapsto (z^k, w, 0)$ is homotopic to an embedding.*

Proof. Just define an explicit embedding by $(z, w) \mapsto (z^k, w, z)$ \square

Remark 1. Lemma 3 generalizes to state that if F^2 is a closed oriented surface of genus g and n is a positive integer, then the homology class in $H_2(F^2 \times D^2)$ corresponding to $n[F^2 \times \{0\}]$ is represented by an embedding of the surface of genus $ng - (n - 1)$. This is the least possible genus that can even represent $n[F^2 \times \{0\}]$ by a map, by Kneser's inequality. For a modern exposition of Kneser's theorem, see [5].

Remark 2. Any map of nonzero degree $T^2 \rightarrow T^2$ is homotopic to a covering, and any covering $T^2 \rightarrow T^2 \times \{0\} \subset T^2 \times D^2$ is homotopic to an embedding. The proof is somewhat more difficult than that of Lemma 3 and depends on lifting a pair of commuting permutations $\sigma, \tau \in S_n$, that generate a transitive subgroup of S_n , to a pair of commuting braids b_σ and b_τ over them in the braid group B_n .

Remark 3. No nontrivial connected covering F^2 of the 0-section of a *nontrivial* oriented 2-disk bundle E over a closed orientable surface G^2 is homotopic to an embedding of F^2 in E . See [1], for example.

It follows from the results of this note that a map $T^2 \rightarrow F^2 \times G^2$ is homologous to an embedding $T^2 \rightarrow F^2 \times G^2$. Is it possible that a map $T^2 \rightarrow F^2 \times G^2$ is actually *homotopic* to an embedding? It seems unlikely.

REFERENCES

1. A. L. Edmonds, *Embedding coverings in bundles*, Indiana University preprint, 1996.
2. P. B. Kronheimer and T. S. Mrowka, *The genus of embedded surfaces in the projective plane*, Math. Research Letters **1** (1994), 797–808. MR **96a**:57073
3. Ronnie Lee and Dariusz Wilczyński, Representing homology classes by locally flat surfaces of minimum genus, preprint (1995)
4. M. D. Meyerson, *Representing homology classes of closed orientable surfaces*, Proc. Amer. Math. Soc. **61** (1976), 181–182. MR **54**:13916
5. R. Skora, *The degree of a map between surfaces*, Math. Ann. **276** (1987), 415–423. MR **88f**:57018

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON, INDIANA 47405
E-mail address: edmonds@indiana.edu