

A CHARACTERIZATION OF GORENSTEIN RINGS IN CHARACTERISTIC $p (> 0)$

S. P. DUTTA

(Communicated by Wolmer V. Vasconcelos)

ABSTRACT. A new characterization of Gorenstein rings in characteristic $p (> 0)$ is proved. It involves asymptotic behaviour of lengths of modules under the Frobenius map.

In this note we are going to study how the Gorenstein property for a local ring $(R, m, k = R/m)$ of characteristic $p (> 0)$, dimension d , is connected with the behaviour of $\lim_{n \rightarrow \infty} \ell(F^n(M))/p^{nd}$ and $\lim_{n \rightarrow \infty} \ell(F^n(M^v))/p^{nd}$, when M is a module of finite length and finite projective dimension over R , $M^v = \text{Hom}_R(M, E)$, E is the injective hull of k over R and $F^n(M)$ is obtained from M by applying the Frobenius map n times ($(M \otimes_R f_R^n$ [P-S]); Notation). It is well known that for a module M of finite length over any local ring R of $\dim d$ and $\text{ch. } p (> 0)$ $\lim_{n \rightarrow \infty} \ell(F^n(M))/p^{nd}$ is positive, and if R is Cohen-Macaulay and $M = R/(x_1, \dots, x_d)$, where x_1, \dots, x_d is a system of parameters in R , the above limit is $\ell(R/(x_1, \dots, x_d))$. Recall that when R is Gorenstein $M^v \simeq \text{Ext}_R^d(M, R)$ for any module M of finite length. Moreover when $pd_R M < \infty$, we have $(F^n(M))^v \simeq \text{Ext}^d(F^n(M), R) \simeq F^n(\text{Ext}^d(M, R)) \simeq F^n(M^v)$. Thus $\ell(F^n(M)) = \ell(F^n(M^v))$, and obviously the above limits are same. This leads us to raise the same question for Cohen-Macaulay rings which are not Gorenstein. Our main theorem shows that the answer, in this case, is in the negative even for cyclic modules of the form $R/(x_1, \dots, x_d)$, where x_1, \dots, x_d is a system of parameters of R . In fact our main theorem is the following:

Theorem. *A Cohen-Macaulay local ring R of dimension d and $\text{ch. } p (> 0)$ is Gorenstein if and only if there exists a system of parameters x_1, \dots, x_d such that*

$$\lim_{n \rightarrow \infty} \ell(F^n((R/\underline{x})^v))/p^{nd} = \ell(R/\underline{x}).$$

Motivation for the study of the above result came from the study of the Strong Intersection Conjecture due to Peskine and Szpiro [P-S]. In this conjecture they assert that, given a pair of finitely generated modules M and N over a local ring R such that $\ell(M \otimes_R N) < \infty$ and $pd_R M < \infty$, $\dim N \leq \text{grade } M$. While studying the conjecture over a Cohen-Macaulay ring of dimension d and positive characteristic, with a canonical module $\Omega (\neq R)$, this author was led to inquire whether $\lim_{n \rightarrow \infty} \ell(H_n^0(F^n(\Omega/(x_2, \dots, x_d)\Omega)))/p^{nd}$ is 0. Recall that, though $\Omega/(x_2, \dots, x_d)\Omega$ is a Cohen-Macaulay module of dimension 1, $F^n(\Omega/(x_2, \dots, x_d)\Omega)$

Received by the editors March 15, 1996 and, in revised form, November 26, 1996.

1991 *Mathematics Subject Classification.* Primary 13D02, 13H10.

This research was partially supported by an NSF grant and an NSA grant.

is not necessarily Cohen-Macaulay. In the course of our proof of the main theorem, we show that the above limit is never zero. If the limit were zero, one could venture for the construction of a counterexample to the above conjecture.

I am thankful to the referee for comments and suggestions, some of which have been incorporated into the body of this paper.

Notations. Throughout this work (R, m, k) denotes a local ring of characteristic $p (> 0)$, dimension d , m its maximal ideal; $k = R/m$ and k is perfect. The Frobenius map $f : R \rightarrow R$, given by $f(x) = x^p$ for all $x \in R$, is a ring homomorphism. We denote by f_R^n the bi-algebra R , having the structure of an R -algebra from the left by f^n and from the right by the identity map, i.e. if $\alpha \in R$, $x \in f_R^n$, $\alpha \cdot x = \alpha^{p^n} x$ and $x \cdot \alpha = x\alpha$. For any module M , we write $F_R^n(M)$ for $M \otimes_R f_R^n$ (we usually drop R from the notation when there is no chance of confusion) and when $\ell(M)$ (= length of M) $< \infty$, M^v for $\text{Hom}_R(M, E)$, where E is the injective hull of k over R .

If $pd_R M < \infty$, $\chi_i(M, N)$ will denote $\sum_{j=0}^{pdM} (-1)^j \ell(\text{Tor}_{i+j}^R(M, N))$. For any finitely generated module M , $\mu(M)$ denotes the minimal number of generators of M over R . We write \lim to denote $\lim_{n \rightarrow \infty}$ and Ω to denote a canonical module of R . For any system of parameters x_1, \dots, x_d of R and for any module M , $\underline{x}_i M$ denotes the submodule $(x_i, \dots, x_d)M$ and \overline{M} denotes $M/x_1 M$. If $\theta = (\theta_{ij})$ denotes an $s \times r$ matrix, then $\theta^{[p^n]}$ stands for the $s \times r$ matrix $(\theta_{ij}^{p^n})$. Also for any module M , and for any element $x \in R$, $(0 : x)M = \{\alpha \in M \mid x\alpha = 0\}$. If I denotes the ideal (y_1, \dots, y_t) in R , then $I^{[p^n]}$ denotes the ideal $(y_1^{p^n}, \dots, y_t^{p^n})$ in R .

Now we state our main theorem.

Theorem. *Let R be a Cohen-Macaulay local ring of dimension d and characteristic $p (> 0)$. Then R is Gorenstein if and only if there exists a system of parameters x_1, \dots, x_d such that*

$$\lim_{n \rightarrow \infty} \ell(F^n((R/\underline{x})^v))/p^{nd} = \ell(R/\underline{x}).$$

Proof. We note that without any loss of generality one can assume R is complete. Recall that for any system of parameters x_1, \dots, x_d of R

$$\lim \ell(F^n(R/\underline{x}))/p^{nd} = \ell(R/\underline{x}).$$

Now let R be Gorenstein. Then we have

$$(F^n(R/\underline{x}))^v \simeq \text{Ext}_R^d(F^n(R/\underline{x}), R) \simeq F^n(\text{Ext}^d(R/\underline{x}, R)) \simeq F^n((R/\underline{x})^v).$$

Hence

$$\lim \ell(F^n((R/\underline{x})^v))/p^{nd} = \lim \ell((F^n(R/\underline{x}))^v)/p^{nd} = \lim \ell(F^n(R/\underline{x}))/p^{nd} = \ell(R/\underline{x}).$$

Next, let us assume that R is not Gorenstein. Our proof is completed in two parts.

In Part 1, we will show that it is possible to construct a system of parameters x_1, \dots, x_d of R such that

$$\lim \ell(F^n((R/\underline{x})^v))/p^{nd} \neq \ell(R/\underline{x});$$

and in Part 2 we will prove the above result for any system of parameters.

Part 1. We first note a rather trivial fact in dimension 0, which will be used in our proof:

Fact. Let R be a local ring of dimension zero. Let M be a module of finite length. Let $\mu(M)$ denote the number of minimal generators of M . Then $\ell(F^n(M)) = \mu(M)\ell(R)$ for $n \gg 0$.

Proof of the Fact. Consider a minimal representation of M :

$$R^r \xrightarrow{\theta} R^s \rightarrow M \rightarrow 0$$

where the entries of θ are in m . Applying $\otimes f_R^n$, we get an exact sequence

$$R^r \xrightarrow{\theta^{[p^n]}} R^s \rightarrow F^n(M) \rightarrow 0.$$

Since $\dim R = 0$, for $n \gg 0$, entries of $\theta^{[p^n]}$ are all 0. Hence the result.

We note that $(R/\underline{x})^v \simeq \text{Ext}^d(R/\underline{x}, \Omega) = \Omega/\underline{x}\Omega$. So we need to establish that for a certain system of parameters x_1, \dots, x_d

$$\ell(R/\underline{x}) (= \lim \ell(F^n(R/\underline{x}))/p^{nd} = \lim \ell(R/\underline{x}^{p^n})/p^{nd}) \neq \lim \ell(F^n(\Omega/\underline{x}\Omega))/p^{nd}.$$

(\underline{x}^{p^n} stands for $(x_1^{p^n}, \dots, x_d^{p^n})$ and $\underline{x}_i^{p^n}$ stands for $(x_i^{p^n}, \dots, x_d^{p^n})$). Note that $\ell(\Omega/\underline{x}\Omega) = \ell(R/\underline{x})$ (for any module M of finite length, $\ell(M) = \ell(M^v)$).

Step 0. $\dim R = 0$. In this case $\ell(F^n(\Omega)) = \mu(\Omega)\ell(R)$ for $n \gg 0$ (Fact).

Thus the above limits are equal if and only if $\mu(\Omega) = 1$ i.e., R is Gorenstein.

Now suppose x_1, \dots, x_d ($d > 0$) is a system of parameters of R such that

$$(1) \quad \lim \ell(R/\underline{x}^{p^n})/p^{nd} = \lim \ell(F^n(\Omega/\underline{x}\Omega))/p^{nd}.$$

Note that the left-hand side is $\ell(R/\underline{x})$. We also recall that $F^n(M \otimes_R N) \simeq F^n(M) \otimes_R F^n(N)$. So $F^n(\Omega/\underline{x}\Omega) \simeq F^n(\Omega) \otimes F^n(R/\underline{x}) = F^n(\Omega) \otimes R/\underline{x}^{p^n}$. Since R is Cohen-Macaulay, $pd_R R/\underline{x} < \infty$.

Step 1. (1) holds if and only if $\Omega \subset R$ and $\chi_1(R/\underline{x}^{p^n}, F^n(\Omega))/p^{nd} \rightarrow 0$ as $n \rightarrow \infty$.

Since $\text{support } \Omega = \text{support } R$ and $F^n(\Omega)_p = F_{R_p}^n(\Omega_p)$ [P-S], we have

$$\begin{aligned} \chi(R/\underline{x}^{p^n}, F^n(\Omega)) &= p^{nd} \chi(R/\underline{x}, F^n(\Omega)) (\chi \text{ is additive}) \\ &= p^{nd} \sum_{P \in \text{Ass}(R)} \ell(F_{R_p}^n(\Omega_p)) \chi(R/\underline{x}, R/P). \end{aligned}$$

We have

$$\chi(R/\underline{x}^{p^n}, F^n(\Omega)) = p^{nd} \sum_{P \in \text{Ass}(R)} \mu_{R_p}(\Omega_p) \ell(R_P) \chi(R/\underline{x}, R/P)$$

for $n \gg 0$ (Fact).

Recall ([L]) that for any finitely generated module T such that $\ell(T/\underline{x}T) < \infty$, $\chi(R/\underline{x}, T) \geq 0$ and equality holds if and only if $\dim T < d$. Moreover $\chi_i(R/\underline{x}, T) \geq 0$. Thus, if $\mu_{R_p}(\Omega_p) > 1$ for any prime $P \in \text{Ass}(R)$, it follows from the above that

$$\lim \chi(R/\underline{x}^{p^n}, F^n(\Omega))/p^{nd} > \sum_{P \in \text{Ass}(R)} \ell(R_P) \chi(R/\underline{x}, R/P).$$

But the right-hand side of the above inequality is nothing but $\chi(R/\underline{x}, R) = \ell(R/\underline{x})$. Since $\chi_1(R/\underline{x}^{p^n}, F^n(\Omega)) \geq 0$, when $\lim \ell(F^n(\Omega/\underline{x}\Omega))/p^{nd} = \ell(R/\underline{x})$, the above inequality forces us to have

$$(2) \quad \mu_{R_p}(\Omega_p) = 1, \forall P \in \text{Ass}(R) \quad \text{and} \quad \lim \chi_1(R/\underline{x}^{p^n}, F^n(\Omega))/p^{nd} = 0.$$

This can easily be checked to be both necessary and sufficient in order that (1) may hold. Note that Ω_p is a canonical module of R_p , and thus (2) implies that R is generically Gorenstein. So, we can assume that Ω is an ideal of height 1 in R . Recall that in such a case R/Ω is Gorenstein.

Now we need to study what it means to have $\lim \chi_1(R/\underline{x}^{p^n}, F^n(\Omega))/p^{nd} = 0$.

Step 2. We can choose x_1 so that $\lim \chi_1(R/\underline{x}^{p^n}, F^n(\Omega))/p^{nd} = 0$ if and only if

$$\lim \ell(H_m^0(F^n(\Omega/\underline{x}_2\Omega)))/p^{nd} = 0.$$

From Step 1, $\Omega \subset R$ is a height one ideal of R . Take $x_1 \in \Omega$.

Consider the short exact sequence

$$0 \rightarrow \Omega \rightarrow R \rightarrow R/\Omega \rightarrow 0.$$

Applying $\otimes f_R^n$, we obtain the following short exact sequences:

$$0 \rightarrow \text{Tor}_1(R/\Omega, f_R^n) \rightarrow F^n(\Omega) \rightarrow \Omega^{[p^n]} \rightarrow 0, \quad 0 \rightarrow \Omega^{[p^n]} \rightarrow R \rightarrow R/\Omega^{[p^n]} \rightarrow 0.$$

Note that $\ell(R/(\Omega + \underline{x}_2)) < \infty$ and hence $\ell(\text{Tor}_i(R/\underline{x}_2, F^n(\Omega))) < \infty$. Now consider the exact sequence

$$0 \rightarrow R/\underline{x}_2^{p^n} \xrightarrow{x_1^{p^n}} R/\underline{x}_2^{p^n} \rightarrow R/\underline{x}^{p^n} \rightarrow 0.$$

Applying $\otimes F^n(\Omega)$ to the above sequence, from the long exact sequence of Tor's we obtain

$$\chi_1(F^n(\Omega), R/\underline{x}^{p^n}) = \ell((0 : x_1^{p^n})F^n(\Omega/\underline{x}_2\Omega)).$$

Again from the exact sequence

$$0 \rightarrow \Omega/\underline{x}_2\Omega \rightarrow R/\underline{x}_2 \rightarrow R/(\Omega + \underline{x}_2) \rightarrow 0,$$

applying $\otimes f_R^n$, we get the following exact sequence:

$$0 \rightarrow \text{Tor}_1(R/(\Omega + \underline{x}_2), f_R^n) \rightarrow F^n(\Omega/\underline{x}_2\Omega) \rightarrow R/\underline{x}_2^{p^n} \rightarrow R/(\Omega + \underline{x}_2)^{[p^n]} \rightarrow 0.$$

Since $x_1 \in \Omega$, it follows from the above sequence that

$$(0 : x_1^{p^n})F^n(\Omega/\underline{x}_2\Omega) = \text{Tor}_1(R/(\Omega + \underline{x}_2), f_R^n) = H_m^0(F^n(\Omega/\underline{x}_2\Omega)).$$

Thus $\lim \chi_1(R/\underline{x}^{p^n}, F^n(\Omega))/p^{nd} = 0$ if and only if

$$(3) \quad \lim \ell(H_m^0(F^n(\Omega/\underline{x}_2\Omega)))/p^{nd} = 0.$$

We have now two cases to consider:

- (a) $\ell(H_m^0(F^n(\Omega/\underline{x}_2\Omega))) \leq Kp^{n(d-2)}$, where K is a constant.
- (b) $\ell(H_m^0(F^n(\Omega/\underline{x}_2\Omega))) = \mathbf{O}(p^{n(d-1)})$.

Step 3. Case (a). We consider the case when $\ell(H_m^0(F^n(\Omega/\underline{x}_2\Omega))) \leq Kp^{n(d-2)}$. We use induction on d (≥ 0). For $d = 0$, see Step 0. Consider the exact sequence

$$(4) \quad 0 \rightarrow H_m^0(F^n(\Omega/\underline{x}_2\Omega)) \rightarrow F^n(\Omega/\underline{x}_2\Omega) \rightarrow N_n \rightarrow 0.$$

Because of (1), it follows that $\lim \ell(N_n/x_1^{p^n}N_n)/p^{nd} = \ell(R/\underline{x})$. But x_1 is not a zero-divisor on N_n , so we have $\lim \ell(N_n/x_1N_n)/p^{n(d-1)} = \ell(R/\underline{x})$. Tensoring (4) by R/x_1R , we get the following exact sequence:

$$(5) \quad 0 \rightarrow \overline{H_m^0(F^n(\Omega/\underline{x}_2\Omega))} \rightarrow F_R^n(\overline{\Omega}/\underline{x}_2\overline{\Omega}) \rightarrow \overline{N}_n \rightarrow 0.$$

(For any module M , we write \overline{M} to denote M/x_1M .) Since $\ell(H_m^0(F^n(\Omega/\underline{x}_2\Omega))) \leq Kp^{n(d-2)}$, $\lim \ell(\overline{H_m^0(F^n(\Omega/\underline{x}_2\Omega))})/p^{n(d-1)} = 0$. This shows that

$$\lim \ell(F_{\overline{R}}^n(\overline{\Omega}/\underline{x}_2\overline{\Omega}))/p^{n(d-1)} = \lim \ell(\overline{N}_n)/p^{n(d-1)} = \ell(R/\underline{x}) = \ell(\overline{R}/\underline{x}_2\overline{R}).$$

Since the dimension of \overline{R} is $d - 1$, we are done by induction on $d = \dim R$ ($\overline{\Omega} = \Omega/x_1\Omega$ is a canonical module for $\overline{R} = R/x_1R$).

Step 4. Case (b). Now we consider the case when $\ell(H_m^0(F^n(\Omega/\underline{x}_2\Omega)))$ is strictly $\mathbf{O}(p^{n(d-1)})$. If $\ell(\overline{H_m^0(F^n(\Omega/\underline{x}_2\Omega))})$ is $\mathbf{O}(p^{n(d-2)})$, we are done by Step 3. So assume that $\ell(\overline{H_m^0(F^n(\Omega/\underline{x}_2\Omega))}) = \mathbf{O}(p^{n(d-1)})$. Recall from Step 2, that $H_m^0(F^n(\Omega/\underline{x}_2\Omega)) = \text{Tor}_1(R/(\Omega + \underline{x}_2), f_R^n)$. We have

$$(6) \quad \ell(H_m^0(F^n(\Omega/\underline{x}_2\Omega))) \leq c p^{n(d-1)},$$

where c is a constant. Suppose that

$$(7) \quad \ell(\overline{H_m^0(F^n(\Omega/\underline{x}_2\Omega))}) = c_1 p^{n(d-1)} + \mathbf{O}(p^{n(d-2)})$$

(the existence of c_1 follows from (5)) and

$$\ell(H_m^0(F^n(\Omega/\underline{x}_2\Omega)) \otimes R/x_1^i R) = c_i p^{n(d-1)} + \mathbf{O}(p^{n(d-2)}).$$

Let t be an integer such that $t > [c/c_1]$. Applying $\otimes R/x_1^i R$ to (4), we obtain the following exact sequence:

$$(8) \quad 0 \rightarrow H_m^0(F^n(\Omega/\underline{x}_2\Omega)) \otimes R/x_1^i R \rightarrow F^n(\Omega/\underline{x}_2\Omega) \otimes R/x_1^i R \rightarrow N_n/x_1^i N_n \rightarrow 0.$$

Since x_1 is a non-zero-divisor on N_n , $\ell(N_n/x_1^i N_n) = i\ell(\overline{N}_n)$. Again, from the exact sequence

$$0 \rightarrow R/x_1 R \rightarrow R/x_1^2 R \rightarrow R/x_1 R \rightarrow 0$$

we get

$$\ell(F^n(\Omega/x_1\Omega) \otimes R/x_1^2 R) = 2\ell(F^n(\Omega/\underline{x}_2\Omega) \otimes R/x_1 R) + \mathbf{O}(p^{n(d-2)})$$

(by applying Proposition 1, [S]). Hence for any fixed i ,

$$\ell(F^n(\Omega/x_1\Omega) \otimes R/x_1^i R) = i\ell(F_{\overline{R}}^n(\overline{\Omega}/\underline{x}_2\overline{\Omega})) + \mathbf{O}(p^{n(d-2)}).$$

Now choose an $i > t$. It then follows from (8) and the above equality that, for $n \gg 0$,

$$\begin{aligned} \ell(H_m^0(F^n(\Omega/\underline{x}_2\Omega)) \otimes R/x_1^i R) &= i\ell(F_{\overline{R}}^n(\overline{\Omega}/\underline{x}_2\overline{\Omega})) - i\ell(\overline{N}_n) + \mathbf{O}(p^{n(d-2)}) \\ &= i\ell(\overline{H_m^0(F^n(\Omega/\underline{x}_2\Omega))}) + \mathbf{O}(p^{n(d-2)}) \quad \text{from (5)} \\ &> t\ell(\overline{H_m^0(F^n(\Omega/\underline{x}_2\Omega))}) + \mathbf{O}(p^{n(d-2)}). \end{aligned}$$

Thus, (6), (7), (8) and the above inequality imply that

$$\ell(H_m^0(F^n(\Omega/\underline{x}_2\Omega)) \otimes R/x_1^i R) > \ell(H_m^0(F^n(\Omega/\underline{x}_2\Omega)))$$

for $n \gg 0$, which is impossible. (The above arguments also show that c cannot be equal to c_1 .) Hence, by Step 3 and Step 4, $\lim \ell(F^n(H_m^0(\Omega/\underline{x}_2\Omega)))/p^{nd}$ cannot be 0. Thus for any system of parameters x_1, \dots, x_d of R , with $x_1 \in \Omega$,

$$\ell(R/\underline{x}) = \lim \ell(F^n(R/\underline{x}))/p^{nd} \neq \lim \ell(F^n((R/\underline{x})^v))/p^{nd}.$$

Part 2. If possible, let there be a system of parameters x_1, \dots, x_d such that

$$\ell(R/\underline{x}) = \lim \ell(F^n(R/\underline{x}))/p^{nd} = \lim \ell(F^n((R/\underline{x})^v))/p^{nd}.$$

Then as in Part 1, we conclude that Ω is a height 1 ideal of R and

$$\lim \chi_1(R/\underline{x}^{p^n}, F^n(\Omega))/p^{nd} = 0.$$

If any $x_i \in \Omega$, we arrive at a contradiction by Theorem 1. So we can assume that not a single x_i is in Ω . Now consider $x_1^{t_1}, x_2^{t_2}, \dots, x_d^{t_d}$, where t_1, \dots, t_d are positive integers. Then

$$\chi_1(R/(x_1^{t_1 p^n}, \dots, x_d^{t_d p^n}), F^n(\Omega)) \leq t_1 t_2 \dots t_d \chi_1(R/\underline{x}^{p^n}, F^n(\Omega)).$$

Hence

$$\lim \chi_1(R/(x_1^{t_1 p^n}, \dots, x_d^{t_d p^n}), F^n(\Omega))/p^{nd} = 0.$$

Thus x_1^t, \dots, x_d^t also satisfy the above equality, i.e.

$$\ell(R/(x_1^t, \dots, x_d^t)) = \lim \ell(F^n(R/(x_1^t, \dots, x_d^t)^v))/p^{nd} \quad (\text{Step 1, Theorem 1}).$$

Now, since $\ell(R/(\Omega + \underline{x})) < \infty$, we can reorganize the ideal (x_1, \dots, x_d) in such a way that x_1, \dots, x_{d-1} is a system of parameters on R/Ω (Theorem 124, [K]). Choose t so that $x_d^t \in \Omega + (x_1, \dots, x_{d-1})$. Then, by our observation above, the system of parameters $x_1, \dots, x_{d-1}, x_d^t$ also satisfies the following:

$$\ell(R/(x_1, \dots, x_{d-1}, x_d^t)) = \lim \ell(F^n(\Omega/(x_1, \dots, x_{d-1}, x_d^t)\Omega))/p^{nd}.$$

But this contradicts the result proved in Part 1, since

$$(x_1, \dots, x_{d-1}, x_d^t) = (x_1, \dots, x_{d-1}, x'_d),$$

where $x'_d \in \Omega$.

Remark. On a complete Cohen-Macaulay local ring R of dimension d , if M is a module of finite length and finite projective dimension, $\ell(M)$ may be different from $\ell(\tilde{M})$, where $\tilde{M} = \text{Ext}^d(M, R)$. (See [R], for such an example.) Still, one can show, by using Theorem 1.5 of [D2], that

$$\lim \ell(F^n(M))/p^{nd} = \lim \ell(F^n(\tilde{M}))/p^{nd}.$$

Actually, a more general statement is valid: for any module M of finite length

$$\lim \ell(\text{Ext}^d(F^n(M), R))/p^{nd} = \lim \ell(F^n(M))/p^{nd}.$$

REFERENCES

- [D1] S. P. Dutta, *Frobenius and Multiplicities*, J. Algebra No. 2 **85** (1983), 424–448. MR **85f**:13022
- [D2] S. P. Dutta, *Ext and Frobenius*, J. Algebra No. 1 **127** (1989), 163–177. MR **91g**:13018
- [K] I. Kaplansky, *Commutative Rings, Revised Edition*, Univ. of Chicago Press (1974). MR **49**:10674
- [L] S. Lichtenbaum, *On the vanishing of Tor in regular local rings*, Illinois J. Math. **10** (1966), 220–226. MR **32**:5688
- [P-S] C. Peskine and L. Szpiro, *Dimension projective finie et cohomologie local*, Publ. I.H.E.S. Paris **42** (1973), 47–119. MR **51**:10330

- [R] P. Roberts, *Intersection theorems*, Commutative algebra, Hochster, Huneke, Sally Editors, MSRI Publications, 15, Springer-Verlag (1989), 417–437. MR **90j**:13024
- [S] G. Seibert, *Complexes with homology of finite length and Frobenius functions*, J. Algebra **125** (1989), 278–287. MR **90j**:13012

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS, 1409 WEST GREEN STREET, URBANA, ILLINOIS 61801

E-mail address: dutta@math.uiuc.edu