

## ENUMERATIONS, COUNTABLE STRUCTURES AND TURING DEGREES

STEPHAN WEHNER

(Communicated by Andreas R. Blass)

ABSTRACT. It is proven that there is a family of sets of natural numbers which has enumerations in every Turing degree except for the recursive degree. This implies that there is a countable structure which has representations in all but the recursive degree. Moreover, it is shown that there is such a structure which has a recursively represented elementary extension.

### 1. INTRODUCTION

In the following we are concerned with countable structures in a recursive language. Researchers have investigated how one could measure the intuitive idea of *information content* of such structures and tried to relate each one of them to a Turing degree [2], [3], [1]. The natural starting point is to look at the collection of *representations*. Let  $\mathfrak{A}$  be a structure. If  $\mathfrak{B}$  is an isomorphic structure with universe  $\omega$ , then  $\mathfrak{B}$  is called a representation of  $\mathfrak{A}$  (written  $\mathfrak{B} \simeq \mathfrak{A}$ ).  $D(\mathfrak{B})$ , its open diagram, can be regarded as a subset of  $\omega$  so that it has a Turing degree, and one can look at the collection of degrees  $\{deg(D(\mathfrak{B})) : \mathfrak{B} \simeq \mathfrak{A}\}$ . A first guess for capturing the complexity of  $\mathfrak{A}$  would be to let its degree be the least element of this collection, especially in the light of the following theorem [2, Theorem 4.1]:

**Theorem 1.1** (Knight). *Let  $\mathfrak{A}$  be a structure in a relational language. Then exactly one of the following holds: (1) For any  $\mathfrak{d} > deg(D(\mathfrak{A}))$ , there is a representation  $\mathfrak{B}$  of  $\mathfrak{A}$  such that  $deg(D(\mathfrak{B})) = \mathfrak{d}$ . (2) There is a finite subset  $S$  of the universe of  $\mathfrak{A}$  such that all permutations of the universe which fix  $S$  are automorphisms of  $\mathfrak{A}$ .*

But this idea fails. For example, Richter [3, Theorem 3.3] shows that for any countable order  $\mathfrak{C}$  which has no recursive representation the collection  $\{deg(D(\mathfrak{B})) : \mathfrak{B} \simeq \mathfrak{C}\}$  has no least element. Therefore more involved concepts have been tried to assign degrees to structures [1].

Now for the particular problems addressed in this paper. Steffen Lempp asked (unpublished): Does a structure with representations in all non-recursive degrees have a recursive representation? Julia Knight asked some related questions: With a binary relation  $R \subseteq \omega^2$  associate a family of subsets of  $\omega$  given by  $\mathcal{F}_R := \{R_n : n \in \omega\}$ , where  $R_n := \{x : (n, x) \in R\}$ ; say that  $R$  is an enumeration of  $\mathcal{F}_R$ . She

---

Received by the editors September 17, 1996 and, in revised form, January 6, 1997.

1991 *Mathematics Subject Classification*. Primary 03D45.

Many thanks go to Julia Knight and Carl Jockusch!

asked (also unpublished): If a family  $\mathcal{F}$  has the feature that for every non-recursive set  $X$ ,  $\mathcal{F}$  has an enumeration recursive in  $X$ , does  $\mathcal{F}$  have a recursive enumeration? Similarly, if for every non-recursive set  $X$ ,  $\mathcal{F}$  has an enumeration r.e. in  $X$ , does  $\mathcal{F}$  have an r.e. enumeration?

In the next section we give some positive results on Knight's questions under extra hypotheses. In Section 3 we prove that the answer to Knight's questions is negative, by constructing a single suitable family. This implies that Lempp's question also has a negative answer, as is shown in the last section. The same finding is obtained in [4], but by another approach. We close by discussing the difference.

The notation is quite standard and follows [5]. All sets considered are subsets of  $\omega$ , the set of natural numbers. We call countable collections of subsets of  $\omega$  *families*. Let  $\varphi$  be a Gödel-numbering of the partial recursive functions (of varying arity) and  $W_i := \text{Rng}(\varphi_i)$  be an enumeration of the recursively enumerable sets as usual.  $W_{i,s}$  is the set of numbers enumerated in  $W_i$  by stage  $s$ . We use recursive bijections  $\langle \cdot, \cdot \rangle$ ,  $\langle \cdot, \cdot, \cdot \rangle$  between  $\omega$  and  $\omega^2$ ,  $\omega$  and  $\omega^3$ , respectively. We also use projections  $(\cdot)_1$  and  $(\cdot)_2$  so that, for example,  $(\langle a, b \rangle)_1 = a$ . Let  $\langle x, A \rangle$  be the set  $\{\langle x, a \rangle : a \in A\}$ . Similarly,  $A_2 := \{(a)_2 : a \in A\}$ . We let  $A + x = \{a + x : a \in A\}$  and  $A - x = \{b : b + x \in A\}$ .

Fix an effective listing  $\Omega$  of recursively enumerable enumerations of all families with r.e. enumerations:

$$\Omega^{(e)} := \{(i, x) : \langle i, x \rangle \in W_e\},$$

and write  $\Omega_i^{(e)}$  for  $\{x : \langle i, x \rangle \in W_e\}$ , the  $i$ -th set of the enumeration  $\Omega^{(e)}$ . Define  $\mathcal{C}^{(e)} := \{\Omega_i^{(e)} : i \in \omega\}$  to be the family enumerated by  $\Omega^{(e)}$ .

$D : \omega \rightarrow 2^\omega$  denotes the canonical enumeration of the family of finite sets; write  $D_n$  for the  $n$ -th finite set. Then the binary predicates  $x \in D_n$  and  $x = |D_n|$  are recursive.

## 2. POSITIVE RESULTS

In this section we give conditions on a family which ensure that the implications of Knight's two questions hold. These are given in Theorems 2.3 and 2.4 and are due to Julia Knight. Jockusch gave a proof of Theorem 2.4 which also showed the following: There is a property which families with a recursive (r.e.) enumeration share with families that have, for all non-recursive degrees  $\mathbf{d}$ , an enumeration recursive (r.e.) in  $\mathbf{d}$ . These seem to be the only positive statements possible about such families.

What is this property? By a rather straightforward forcing construction it follows that if a set of natural numbers is recursive (r.e.) in all non-recursive degrees, then it is recursive (r.e.). Therefore, the members of a family are recursive (recursively enumerable) if the family has, for all non-recursive degrees  $\mathbf{d}$ , an enumeration recursive (r.e.) in  $\mathbf{d}$ . Hence such families  $\mathcal{F}$  are fully described by the index set

$$I_r(\mathcal{F}) := \{i : (\exists A \in \mathcal{F})(\varphi_i = \chi_A)\}$$

or, respectively,

$$I_{re}(\mathcal{F}) := \{i : (\exists A \in \mathcal{F})(W_i = A)\}.$$

Both the index set  $I_r$  of a family with a recursive enumeration and the index set  $I_{re}$  of a family with a recursively enumerable enumeration are  $\Sigma_3^0$  in the arithmetical hierarchy. Here is the coincidence:

**Theorem 2.1.** *Let  $\mathcal{F}$  be a family. If, for every non-recursive degree  $\mathbf{d}$ ,  $\mathcal{F}$  has an enumeration recursive in  $\mathbf{d}$ , then its index set  $I_r(\mathcal{F})$  is  $\Sigma_3^0$ .*

**Theorem 2.2** (Jockusch). *Let  $\mathcal{F}$  be a family. If, for every non-recursive degree  $\mathbf{d}$ ,  $\mathcal{F}$  has an enumeration recursively enumerable in  $\mathbf{d}$ , then its index set  $I_{re}(\mathcal{F})$  is  $\Sigma_3^0$ .*

Towards giving a sufficient condition under which the implication of her first question holds, Julia Knight defines an *extension function* for a family  $\mathcal{F}$  to be a (possibly partial) function  $f: 2^{<\omega} \rightarrow \omega$  such that if  $\sigma \in 2^{<\omega}$  and there exists a set  $A \in \mathcal{F}$  such that  $\chi_A \supseteq \sigma$ , then  $\varphi_{f(\sigma)} = \chi_A$  for some such  $A$ . We mention two facts: Any family with a recursive enumeration has a partial recursive extension function, and so does a family containing all finite sets.

We prove Theorems 2.1 and 2.2 simultaneously with the following two.

**Theorem 2.3** (Knight). *Let  $\mathcal{F}$  be a family which has, for all non-recursive  $\mathbf{d}$ , an enumeration recursive in  $\mathbf{d}$ . If  $\mathcal{F}$  has a partial recursive extension function, then  $\mathcal{F}$  has a recursive enumeration.*

*Proof (of Theorems 2.3 and 2.1).* Let  $\mathcal{F}$  be a family which has, for any non-recursive set  $X$ , an enumeration recursive in  $X$ . We construct a generic set  $D$ , attempting to meet the following requirements and expecting to fail. Below, this failure will be exploited to prove the statements of the two theorems for  $\mathcal{F}$  separately.

$R_e$ :  $\varphi_e^D$  is not the characteristic function of an enumeration of  $\mathcal{F}$ .

The set  $D$  will be Cohen-generic. The set of forcing conditions is  $2^{<\omega}$  and the partial order is given by  $\subseteq$ . We use the old-fashioned notion of a *complete forcing sequence* (c.f.s.), where  $p_{n+1} \supseteq p_n$ , with  $p_{n+1}$  entering the  $n$ th dense set in some countable collection.  $D$  is the set with characteristic function  $\bigcup_{n \in \omega} p_n$ .

Fix a condition  $p \in 2^{<\omega}$  and  $e \in \omega$ . We consider the following four possibilities for  $p$  and  $e$ , showing how extensions of  $p$  may force satisfaction of  $R_e$  in each case.

- P1. For some  $q \supseteq p, n, x \in \omega, q \vdash \varphi_e^D(n, x) \downarrow \neq 0, 1$ , or  $q \vdash \varphi_e^D(n, x) \uparrow$ .  
We include  $q$  in the c.f.s., thereby satisfying  $R_e$ .
- P2. For some  $q \supseteq p$  and some  $n$ , for all  $q' \supseteq q$  there exist  $x$  and  $r_0, r_1 \supseteq q'$  such that  $r_i \vdash \varphi_e^D(n, x) = i$ .  
For each  $A \in \mathcal{F}$  the set

$$D_A^1 := \{r : r \supseteq q \Rightarrow (\exists x)(r \vdash \varphi_e^D(n, x) \neq \chi_A(x))\}$$

is dense. We add  $q$  to the c.f.s. and enter the sets  $D_A^1$ . Then the requirement  $R_e$  is satisfied.

We write  $q \vdash E_n = A$  if for all  $x$  and all  $q' \supseteq q$  there is an  $r \supseteq q'$  such that  $r \vdash \varphi_e^D(n, x) \downarrow = \chi_A(x)$ . Note that for  $q$  and  $n$  there is at most one set  $A$  such that  $q \vdash E_n = A$ .

- P3. For some  $q \supseteq p, n \in \omega$  and  $B \notin \mathcal{F}$  we have  $q \vdash E_n = B$ .  
By putting  $q$  into the c.f.s., we meet  $R_e$ .
- P4. Not P2 but there exist  $A \in \mathcal{F}$  and  $q \supseteq p$  such that for all  $q' \supseteq q$  and all  $n$ , if  $q' \vdash E_n = B$  then  $A \neq B$ .

Since P2 does not hold, the set

$$D_n^2 := \{q' : q' \supseteq q \Rightarrow (\exists B)(q' \vdash E_n = B)\}$$

is dense for each  $n$ . By including  $q$  in the c.f.s. and entering the sets  $D_n^2$ , we meet  $R_e$ .

If for all  $p \in 2^{<\omega}$  and  $e \in \omega$  one of the cases P1, ..., P4 holds, then the forcing construction yields a generic (and so non-recursive) set  $D$ , in which  $\mathcal{F}$  has no recursive enumeration, contrary to the assumption on  $\mathcal{F}$ .

So let  $p \in 2^{<\omega}$  and  $e \in \omega$  be such that none of the cases P1, ..., P4 hold. It follows that if  $q \vdash E_n = A$ , then  $A \in \mathcal{F}$  for all  $q \supseteq p$ ,  $n$  and  $A \subseteq \omega$ . Moreover, all elements of  $\mathcal{F}$  occur in this way.

We first complete the proof of Theorem 2.3. Let  $f$  be a partial recursive extension function for  $\mathcal{F}$ . Fix  $n \in \omega$  and  $p \subseteq q \in 2^{<\omega}$ . Say that  $\sigma \in 2^{<\omega}$  has a  $q$ -computation if there is an  $r \supseteq q$  such that  $r \vdash \varphi_e^D(n, x) = \sigma(x)$  for all  $x \in \text{dom}(\sigma)$ . We make the following observations:

- $q \vdash E_n = A$  and  $q' \supseteq q$  implies  $q' \vdash E_n = A$ .
- The set  $\{\sigma \in 2^{<\omega} : \sigma \text{ has a } q\text{-computation}\}$  is r.e. (uniformly in  $q$ ).
- By definition,  $q \vdash E_n = A$  if and only if any  $\sigma \in 2^{<\omega}$  has a  $q$ -computation if and only if  $\sigma \subseteq \chi_A$ .
- Since  $p, e$  do not satisfy P1, for any  $q' \supseteq q$  there is a  $\sigma \in 2^{<\omega}$  which has a  $q'$ -computation.
- The previous two items imply that the predicate  $q \supseteq p \Rightarrow (\forall A)(q \not\vdash E_n = A)$  in  $q$  and  $n$  is r.e.
- Since  $p, e$  do not satisfy P3, if  $q \supseteq p$  and  $q \not\vdash E_n = A$  for any  $A \in \mathcal{F}$ , then for any  $\sigma \in 2^{<\omega}$  with a  $q$ -computation there is a set  $A \in \mathcal{F}$  such that  $\sigma \subseteq \chi_A$ .

We form a recursive enumeration  $R$  of  $\mathcal{F}$ , using pairs  $(q, n)$  as indices, where  $q \in 2^{<\omega}$  with  $q \supseteq p$  and  $n \in \omega$ . Since there is a recursive bijection between  $P \times \omega$  and  $\omega$ , where  $P = \{q \in 2^{<\omega} : q \supseteq p\}$ , this suffices. Let  $(\sigma_q^{(n)})_{n \in \omega}$  be an effective enumeration of  $\{\sigma \in 2^{<\omega} : \sigma \text{ has a } q\text{-computation}\}$  by the second observation. We choose this enumeration so that, additionally, for every  $\sigma$  with a  $q$ -computation there are infinitely many  $i$  such that  $\sigma_q^{(i)} = \sigma$ . Define a partial recursive function  $a_q$  by  $a_q(0) := 0$  and  $a_q(n+1) := \mu m > n. \sigma_q^{(m)} \supseteq \sigma_q^{(a_q(n))}$ . By the fourth observation, let  $h : \omega \rightarrow 2^{<\omega} \times \omega$  be a recursive function with range  $\{(q, n) : q \supseteq p \text{ and } (\forall A)(q \not\vdash E_n = A)\}$ .

$R_{(q,n)}$  is defined by

$$R_{(q,n)} := \begin{cases} \bigcup_{0 \leq i \leq m} \sigma_q^{(a_q(i))} \cup \varphi_{f(\sigma_q^{(a_q(m))})} & \text{if } m \text{ is least such that } h(m) = (q, n), \\ \bigcup_{i \in \omega} \sigma_q^{(a_q(i))} & \text{otherwise.} \end{cases}$$

Fix  $q \supseteq p$  and  $n \in \omega$ . By definition of the functions  $h$ , there is  $m$  satisfying the first case if and only if  $(\forall A)(q \not\vdash E_n = A)$ . If  $q \vdash E_n = A$ , then the choice of the enumeration  $(\sigma_q^{(n)})_{q \in 2^{<\omega}, n \in \omega}$  and the function  $a$  guarantees  $R_{(q,n)} = A$ . If  $q \not\vdash E_n = A$  for any  $A$ , then by the fact that  $f$  is an extension function for  $\mathcal{F}$ , it follows that  $R_{(q,n)} \in \mathcal{F}$ .

Since for every  $A \in \mathcal{F}$  there are  $q$  and  $n$  such that  $q \vdash E_n = A$ , it follows that  $R$  is an enumeration of  $\mathcal{F}$ , and Theorem 2.3 is proved.

We turn to the proof of Theorem 2.1. As mentioned above, if none of the cases holds for  $p$  and  $e$ , then we have

$$\mathcal{F} = \{X : (\exists q \supseteq p)(\exists n)(q \vdash E_n = X)\}.$$

The ternary relation  $(q \vdash E_n = X) \wedge (\chi_X = \varphi_i)$  (in  $q, n$  and  $i$ ), when restricted to  $i$  such that  $\varphi_i$  is the characteristic function of a set, is  $\Pi_2^0$  so that  $I_r(\mathcal{F})$  is  $\Sigma_3^0$ .  $\square$

**Theorem 2.4** (Knight). *Let  $\mathcal{F}$  be a family such that for all non-recursive  $X$ ,  $\mathcal{F}$  has an enumeration r.e. in  $X$ . If  $\mathcal{F}$  contains all finite sets, then  $\mathcal{F}$  has an r.e. enumeration.*

*Proof of Theorems 2.4 and 2.2.* As above, we construct a generic set  $D$ , attempting to meet the following requirements and expecting to fail.

$R_e$ .  $W_e^D$  is not an enumeration of  $\mathcal{F}$ .

We use the same forcing notion as was used for the previous proof. Fix  $p \in 2^{<\omega}$  and  $e \in \omega$ . We consider the following three cases.

- C1. For some  $q \supseteq p$  and some  $n \in \omega$ , for all  $q' \supseteq q$ , there exist  $x \in \omega$  and  $r_0, r_1 \supseteq q'$  such that  $r_0 \vdash (n, x) \in W_e^D$  and  $r_1 \vdash (n, x) \notin W_e^D$ .

For each  $A \in \mathcal{F}$  the set

$$D_A^1 := \{r : r \supseteq q \Rightarrow (\exists x)[(r \vdash (n, x) \in W_e^D \wedge x \notin A) \text{ or } (r \vdash (n, x) \notin W_e^D \wedge x \in A)]\}$$

is dense. We put  $q$  into the c.f.s. and enter the sets  $D_A^1$ . Requirement  $R_e$  is satisfied.

We write  $q \vdash E_n = A$  if for all  $x$ , if  $x \in A$ , then for all  $q' \supseteq q$  there is  $r \supseteq q'$  such that  $r \vdash (n, x) \in W_e^D$ , and if  $x \notin A$  then  $q \vdash (n, x) \notin W_e^D$ .

- C2. Case C1 does not hold, but  $q \vdash E_n = B$  for some  $q \supseteq p, n \in \omega$  and  $B \notin \mathcal{F}$ .

Requirement  $R_e$  is met by putting  $q$  in the c.f.s.

- C3. Case C1 does not hold, but there exists  $A \in \mathcal{F}$  such that for all  $q' \supseteq q$  and all  $n$ , if  $q' \vdash E_n = B$ , then  $A \neq B$ .

Since Case C1 does not hold, the set

$$D_n^2 := \{q' : q' \supseteq q \Rightarrow (\exists B)(q' \vdash E_n = B)\}$$

is dense for all  $n$ . We meet the requirement  $R_n$  by including  $q$  in the c.f.s. and entering all sets  $D_n^2$ .

If for all  $p \in 2^{<\omega}$  and  $e \in \omega$  one of the cases C1, C2, or C3 holds, then the forcing construction yields a generic (and hence non-recursive) set  $D$ , in which  $\mathcal{F}$  has no r.e. enumeration. This contradicts the assumption of both Theorem 2.2 and Theorem 2.4. So let  $p \in 2^{<\omega}$  and  $e \in \omega$  be such that none of the cases C1, C2, C3 holds. We show that the index set  $I_{re}$  of  $\mathcal{F}$  is  $\Sigma_3^0$ . As in the proof of Theorem 2.3, for all  $q \supseteq p, n$ , and  $A \subseteq \omega$ , if  $q \vdash E_n = A$ , then  $A \in \mathcal{F}$ , and all members of  $\mathcal{F}$  occur in this way. Therefore we have

$$\{i : W_i \in \mathcal{F}\} = \{i : (\exists q \supseteq p)(\exists n)(q \vdash E_n = W_i)\}.$$

The relation “ $q \vdash E_n = W_i$ ” is  $\Pi_2^0$ , and so the index set of  $\mathcal{F}$  is  $\Sigma_3^0$ . This completes the proof of Theorem 2.2. To complete the proof of Theorem 2.4, note that since  $\mathcal{F}$  includes all finite sets and only contains r.e. sets, by a theorem of Yates [7, Theorem 8],  $\mathcal{F}$  has an r.e. enumeration.  $\square$

3. A FAMILY OF FINITE SETS

In this section we first define a family  $\mathcal{C}$  which has no r.e. enumeration. Then we show that for every non-recursive set  $X$  there is an enumeration of  $\mathcal{C}$  which is recursive in  $X$ , and finally that every non-recursive degree contains an enumeration of  $\mathcal{C}$ . This corrects an earlier statement in [6, p 187]. In particular, we apologize for connecting the error with Martin Kummer.

Let  $r$  be the partial recursive function defined by

$$r(e) := (\mu \langle i, x, s \rangle . \langle e, x \rangle \in \Omega_{i,s}^{(e)}) . 1.$$

Informally, the value of  $r(e)$  is the first index  $i$  to be found such that there is a number of the form  $\langle e, x \rangle \in \Omega_i^{(e)}$ ; if there is no such  $i$  then  $r(e)$  is not defined. Let the family  $\mathcal{C}$  be defined by

$$\mathcal{C} := \{ \langle e, A \rangle : A \text{ is finite, } e \in \omega \} - \{ \langle e, \omega \rangle \cap \Omega_{r(e)}^{(e)} : r(e) \downarrow \}.$$

$\mathcal{C}$  does not have an r.e. enumeration; for suppose  $\Omega^{(e_0)}$  is an enumeration of  $\mathcal{C}$ . Then  $r(e_0)$  is defined, and the set  $\Omega_{r(e_0)}^{(e_0)} = \langle e, \omega \rangle \cap \Omega_{r(e_0)}^{(e_0)}$  is not a member of  $\mathcal{C}$ .

Let  $X$  be an arbitrary non-recursive set. To see how to construct an enumeration  $S^X$  of  $\mathcal{C}$  such that  $S^X \leq_T X$  we use the following lemma.

**Lemma 3.1.** *Uniformly in  $i$  and recursively in  $X$  there is a finite set  $A_i^X \leq_T X$  such that  $W_i \neq A_i^X$ .*

Let  $g$  be a partial recursive function such that  $g(e) \downarrow$  if and only if  $r(e) \downarrow$  and if  $r(e) \downarrow$  then  $W_{g(e)} = (\langle e, \omega \rangle \cap \Omega_{r(e)}^{(e)}) . 2$ . Let  $h$  be a partial recursive function such that  $h(e, a) \downarrow$  if and only if  $g(e) \downarrow$  and if  $g(e) \downarrow$  then  $W_{h(e,a)} = W_{g(e)} - a$ .

Define

$$S_{\langle n,t,e \rangle}^X := \langle e, B^X(n, t, e) \rangle,$$

where

$$B^X(n, t, e) := \begin{cases} D_n \cup (s_0 + A_{h(e,s_0)}^X) & \text{if there is } s > t, \max(D_n) + 1 \\ & \text{such that } g_s(e) \downarrow \text{ and } W_{g(e),s} = D_n, \\ & \text{and } s_0 \text{ is the least such,} \\ D_n & \text{otherwise.} \end{cases}$$

By the lemma,  $S^X$  is recursive in  $X$ . We claim that  $S^X$  is an enumeration of  $\mathcal{C}$ .

“ $\subseteq$ ”. First of all, it follows from the lemma that all sets  $B^X(n, t, e)$  are finite and so the family enumerated by  $S^X$  is contained in  $\{ \langle e, A \rangle : A \text{ is finite} \}$ .

Suppose  $r(e) \downarrow$  and  $S_{\langle n,t,e \rangle}^X = \langle e, \omega \rangle \cap \Omega_{r(e)}^{(e)}$ . It follows that  $B^X(n, t, e) = W_{g(e)}$ . There has to be a stage  $s > t, \max(D_n) + 1$  such that  $W_{g(e),s} = D_n$ , because otherwise  $B^X(n, t, e) = D_n \neq W_{g(e)}$ , a contradiction. Let  $s_0$  be the least such  $s$ . Then  $B^X(n, t, e) = D_n \cup (s_0 + A_{h(e,s_0)}^X) = W_{g(e)}$ , so that  $A_{h(e,s_0)}^X = W_{h(e,s_0)}$ , a contradiction.

“ $\supseteq$ ”. Let  $e, n \in \omega$  such that  $C = \langle e, D_n \rangle \in \mathcal{C}$ . We want to find a number  $z$  such that  $S_z^X = C$ . Let  $t$  be such that  $W_{g(e),s} \neq D_n$  for all  $s > t$ . (If there is no such  $t$  then  $C \notin \mathcal{C}$ .) By definition,  $B^X(n, t, e) = D_n$ , so that  $S_{\langle n,t,e \rangle}^X = C$ .

We have constructed an enumeration of  $\mathcal{C}$  which is recursive in  $X$ . Simple coding is sufficient to obtain an enumeration T-equivalent to  $X$ : Choose  $A, B \in \mathcal{C}$  so that

$A - B \neq \emptyset$ . Define another enumeration of  $\mathcal{C}$ ,

$$P_{2n}^X := S_n^X, \quad P_{2n+1}^X := \begin{cases} A & \text{if } n \in X, \\ B & \text{if } n \notin X. \end{cases}$$

Clearly,  $P^X$  is recursive in  $X$  and enumerates  $\mathcal{C}$ . Let  $x \in A - B$ . Then  $z \in X$  if and only  $x \in P_{2z+1}^X$ , so that  $X$  is recursive in  $P^X$ .

*Proof of Lemma 3.1.* Let  $y, n : \omega^2 \rightarrow \omega$  be two recursive one-one functions such that their ranges are disjoint and cover  $\omega$ . Let

$$\alpha(s, x) := \begin{cases} y(s, x) & \text{if } x \in X, \\ n(s, x) & \text{otherwise,} \end{cases}$$

so that  $\alpha \leq_T X$ . We construct the set  $A_i^X$  in stages uniformly in  $i$  and recursively in  $X$ . In the course of the construction, numbers  $a_x$  may become defined.

Stage 0. Set  $x := 0$ .  $A_{i,0}^X$  is empty.

Stage  $s + 1$ . If  $W_{i,s} \neq A_{i,s}^X$ , pass to the next stage. Otherwise enumerate

$\alpha(s, x)$  in  $A_i^X$ , let  $a_x := s$  and increase  $x$  by one.

End of construction.

The set  $A_i^X$  is (by construction) r.e. in  $X$ . It is also recursive in  $X$ : By inspection,  $A_i^X$  only contains numbers  $\alpha(a, b)$ . If  $y(s, x) \in A_i^X$ , then  $y(s, x) \in A_{i,s+1}^X$  and the same holds for numbers  $n(s, x)$ . This together with the choice of  $y$  and  $n$  is sufficient.

*Claim 1.* The set  $A_i^X$  is finite.

Suppose  $A_i^X$  is infinite. During the construction of  $A_i^X$  infinitely many numbers  $a_x$  are defined. By induction on  $x$ , it follows that  $A_{i,a_x}^X = \{\alpha(a_j, j) : j \leq x\}$ , and therefore  $W_{i,a_x} = \{\alpha(a_j, j) : j < x\}$  for all  $x \in \omega$ . Thus,  $W_i = \{\alpha(a_x, x) : x \in \omega\}$ . Now,  $x \in X$  if and only if  $(\exists t)(y(t, x) \in W_i)$ , and  $x \notin X$  if and only if  $(\exists t)(n(t, x) \in W_i)$ . This means that  $X$  is recursive, a contradiction.

*Claim 2.*  $A_i^X$  is different from  $W_i$ .

By the previous claim it is sufficient to consider the case when  $W_i$  is finite. Let  $s_0$  be the least number such that  $W_{i,s_0} = W_i$ . Then either  $W_{i,s_0} \neq A_{i,s_0}^X$ , in which case  $W_{i,s_0} \neq A_{i,s_0}^X = A_i^X$ ; or  $W_{i,s_0} = A_{i,s_0}^X$ , so that at stage  $s_1 = s_0 + 1$  a new number is enumerated in  $A_i^X$ , whence  $W_i = W_{i,s_1} = A_{i,s_0}^X \neq A_i^X$ .

Lemma 3.1 is proved. □

#### 4. APPLICATION TO LEMPP'S QUESTION

Let  $\mathcal{F}$  be a family. With  $\mathcal{F}$  we associate the following countable structure  $\mathfrak{A}_{\mathcal{F}}$  in the language  $L = (S, Z, I)$ , where  $S$  is a binary predicate symbol and  $Z$  and  $I$  are unary predicate symbols. The universe of  $\mathfrak{A}_{\mathcal{F}}$  is  $\mathcal{F} \times \omega \times \omega$ . For every  $A \in \mathcal{F}$ , set  $Z((A, x, 0))$  and  $S((A, x, n), (A, x, n + 1))$ . Set  $I((A, x, n))$  if and only if  $n \in A$ . Thus, countably many  $S$ -chains  $(A, x, 0), (A, x, 1), (A, x, 2), \dots$  are associated with every  $A \in \mathcal{F}$ , and in every chain  $I$  holds of the  $n$ -th member if  $n \in A$ .

**Theorem 4.1.** *Let  $\mathbf{d}$  be a Turing degree and  $\mathcal{F}$  be a family. Then  $\mathcal{F}$  has an enumeration recursive in  $\mathbf{d}$  if and only if the structure  $\mathfrak{A}_{\mathcal{F}}$  has a representation recursive in  $\mathbf{d}$ .*

*Proof.* “ $\Rightarrow$ ”. Let  $Q^X$  be an enumeration of  $\mathcal{F}$  which is recursive in  $X \in \mathbf{d}$ . Define  $R^X$  to be another  $X$ -recursive enumeration of  $\mathcal{F}$  by  $R_{\langle n, i \rangle}^X := Q_i^X$ . A representation of  $\mathfrak{A}_{\mathcal{F}}$  is given by  $\mathfrak{B}$ , where  $Z_{\mathfrak{B}}(\langle x, 0 \rangle)$ ,  $S_{\mathfrak{B}}(\langle x, n \rangle, \langle x, n + 1 \rangle)$ , and  $I_{\mathfrak{B}}(\langle x, n \rangle)$  if and only if  $x \in R_n^X$ . These predicates are recursive in  $R^X$ , and therefore  $D(\mathfrak{B})$  is recursive in  $X$ .

“ $\Leftarrow$ ”. Let  $\mathfrak{B}$  be a representation of  $\mathfrak{A}_{\mathcal{F}}$  whose open diagram is recursive in  $X \in \mathbf{d}$ . The set  $Z := \{x : Z_{\mathfrak{B}}(x)\}$  is recursive in  $X$ . With each  $x \in Z$  we associate the set  $S_x := \{n : I_{\mathfrak{B}}(f^{(n)}(x))\}$ , where  $f(x)$  is the unique  $y$  such that  $S_{\mathfrak{B}}(x, y)$ , and  $f^{(n)}(x)$  denotes the  $n$ -fold application of  $f$  to  $x$ . The sets  $S_x$  are uniformly in  $x \in Z$  recursive in  $X$ , and, by definition of  $\mathfrak{A}_{\mathcal{F}}$ ,  $\mathcal{F} = \{S_x : x \in Z\}$ . This suffices.  $\square$

**Corollary 4.2** (Slaman [4]). *There is a structure which has representations only in the non-recursive degrees.*

*Proof.* Apply the theorem to the family  $\mathcal{C}$  defined in the previous section to obtain representations of  $\mathfrak{A}_{\mathcal{C}}$  below any non-recursive T-degree. Apply Theorem 1.1 to obtain representations in all non-recursive T-degrees.  $\square$

Slaman remarked (private communication) that the construction given in [4] yields a structure which is not elementarily equivalent to any recursively represented structure. Informally, the reason for this is as follows. Essentially, the construction proceeds in such a way that the final outcome of the actions taken to diagonalize against the recursive representations can be read off the theory of the structure (which is called  $\mathfrak{M}$ ):

He writes ([4, Section 2.1]): “We will ensure that either  $R^{-1}(T_i)^{\mathfrak{R}_i} = \emptyset$ , or  $\langle T_i, <_L \upharpoonright T_i \rangle^{\mathfrak{R}_i}$  is not isomorphic to  $\langle T_i, <_L \upharpoonright T_i \rangle^{\mathfrak{M}}$ , or there is a  $p$  in  $R^{-1}(T_i)^{\mathfrak{R}_i}$  such that  $\zeta(p)^{\mathfrak{R}_i}$  is not maximal, or there is a  $p$  in  $R^{-1}(T_i)^{\mathfrak{R}_i}$  such that  $\zeta(p)^{\mathfrak{R}_i}$  is infinite. Since none of these disjuncts apply to  $\mathfrak{M}$ , we will thus ensure that  $\mathfrak{M}$  has no recursive presentation.”

Fix  $i \in \omega$ . The first and third disjunct can be directly formalized in the language  $\mathcal{L}$  of the structure. If the second disjunct holds, then the construction [4, Section 2.2.1] yields a tree  $\langle T_i, <_L \upharpoonright T_i \rangle^{\mathfrak{M}}$  which is finite. Therefore this tree is described by a sentence in the theory of  $\mathfrak{M}$ . The fourth disjunct also cannot be formalized in  $\mathcal{L}$ , but, provided the first three disjuncts are not true, the strategy used in [4, Section 2.2.2] results in

$$(\exists p)(R(p, s^i(0)) \wedge (\forall x)(R(p, x) \rightarrow (\exists y)(x <_T y \wedge R(p, y))))$$

being true in  $\mathfrak{R}_i$ . This is a formula in  $\mathcal{L}$ , and (by the same strategy) not true in  $\mathfrak{M}$ . Hence there is no recursively represented structure which is elementarily equivalent to  $\mathfrak{M}$ .

The structure obtained from the family  $\mathcal{C}$  by Theorem 4.1 is of a different kind:

**Theorem 4.3.** *There is a structure which has representations only in the non-recursive degrees and has a recursively represented elementary extension.*

*Proof.* Let us look at the following family:

$$\mathcal{D} := \{\langle e, A \rangle : A \text{ is finite, } e \in \omega\}.$$

Obviously,  $\mathcal{D}$  has a recursive enumeration, and  $\mathfrak{A}_{\mathcal{D}}$  has a recursive representation. At the same time, the structure  $\mathfrak{A}_{\mathcal{C}}$  from the proof of Corollary 4.2 is contained in  $\mathfrak{A}_{\mathcal{D}}$ , and they are elementarily equivalent:

It suffices to show that for any formula  $(\exists x)(\phi)$  in variables  $x_1, \dots, x_n$ , if

$$\mathfrak{A}_{\mathcal{D}} \models (\exists x)(\phi)[x_1 := c_1, \dots, x_n := c_n]$$

and  $c_1, \dots, c_n \in \mathfrak{A}_{\mathcal{C}}$ , then there is  $c \in \mathfrak{A}_{\mathcal{C}}$  such that

$$\mathfrak{A}_{\mathcal{D}} \models \phi[x := c, x_1 := c_1, \dots, x_n := c_n].$$

Note that if  $\mathfrak{A}_{\mathcal{D}} \models (\exists x)(\phi)[x_1 := c_1, \dots, x_n := c_n]$ , then there is a  $d$  in  $\mathfrak{A}_{\mathcal{D}}$  such that either

$$\begin{aligned} \mathfrak{A}_{\mathcal{D}} \models & \phi[x := d, x_1 := c_1, \dots, x_n := c_n] \\ & \wedge \neg S(d, c_1) \wedge \neg S(c_1, d) \wedge \dots \wedge \neg S(d, c_n) \wedge \neg S(c_n, d), \end{aligned}$$

or

$$\begin{aligned} \mathfrak{A}_{\mathcal{D}} \models & \phi[x := d, x_1 := c_1, \dots, x_n := c_n] \\ & \wedge (S(d, c_1) \vee S(c_1, d) \vee \dots \vee S(d, c_n) \vee S(c_n, d)). \end{aligned}$$

In the former case, choose  $c$  from  $\mathfrak{A}_{\mathcal{C}}$  outside of the  $S$ -chains of  $\mathfrak{A}_{\mathcal{C}}$  which  $c_1, \dots, c_n$  belong to such that it satisfies  $Z$  and  $I$  in the same way  $d$  does. In the latter case,  $d$  is already part of an  $S$ -chain which is contained in  $\mathfrak{A}_{\mathcal{C}}$ , and so an element of  $\mathfrak{A}_{\mathcal{C}}$ .  $\square$

#### REFERENCES

- [1] Christopher J. Ash, Carl G. Jockusch, jr. and Julia F. Knight; *Jumps of orderings*, Trans. Amer. Math. Soc., vol. 319, (1990), p. 573 – 599. MR **90j**:03081
- [2] Julia F. Knight; *Degrees Coded in Jumps of Orderings*, J. Symbolic Logic, vol. 51, (1986), p. 1034 – 1042. MR **88j**:03030
- [3] Linda J. Richter; *Degrees of Structures*, J. Symbolic Logic, vol. 46 (1981), p. 723 – 731. MR **83d**:03048
- [4] Theodore A. Slaman; *Relative to any Nonrecursive Set*, Proc. Amer. Math. Soc., vol. 126 (1998), 2117–2122. CMP 97:11
- [5] Robert I. Soare; *Recursively Enumerable Sets and Degrees*, Springer Verlag, Berlin, Heidelberg, New York, Tokyo, 1987. MR **88m**:03003
- [6] Stephan Wehner; *On Injective Enumerability of Recursively Enumerable Classes of Cofinite Sets*, Arch. Math. Logic, vol. 34, (1995), p. 183 – 196. MR **96d**:03062
- [7] C.E.M. Yates; *On the Degrees of Index Sets II*, Trans. Amer. Math. Soc., vol. 135 (1969), p. 249 – 266. MR **39**:2637

DEPARTMENT OF CHEMISTRY, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, BRITISH COLUMBIA, CANADA V6T 1Z1

*E-mail address:* `stephan@pepe.chem.ubc.ca`