

THE EQUIVALENCE OF SOME BERNOULLI CONVOLUTIONS TO LEBESGUE MEASURE

R. DANIEL MAULDIN AND KÁROLY SIMON

(Communicated by Frederick W. Gehring)

ABSTRACT. Since the 1930's many authors have studied the distribution ν_λ of the random series $Y_\lambda = \sum \pm \lambda^n$ where the signs are chosen independently with probability $(1/2, 1/2)$ and $0 < \lambda < 1$. Solomyak recently proved that for almost every $\lambda \in [\frac{1}{2}, 1]$, the distribution ν_λ is absolutely continuous with respect to Lebesgue measure. In this paper we prove that ν_λ is even equivalent to Lebesgue measure for almost all $\lambda \in [\frac{1}{2}, 1]$.

1. INTRODUCTION

For each $\lambda \in (0, 1)$ we define the random variable

$$Y_\lambda = \sum_{n=0}^{\infty} \theta_n \cdot \lambda^n,$$

where θ_n are independent random variables with $Prob(\theta_n = -1) = Prob(\theta_n = 1) = \frac{1}{2}$. The distribution ν_λ of Y_λ is sometimes called a symmetric infinite Bernoulli convolution. One can easily see that for $0 < \lambda < \frac{1}{2}$ the distribution ν_λ is supported on a Cantor set of zero Lebesgue measure. Since the 1930's a lot of work has been done to characterize ν_λ for $\frac{1}{2} < \lambda$ (for a good survey see e.g. Peres, Solomyak (1996a) [4]). Among these results the most interesting ones are as follows: P. Erdős (1939) [1] proved that ν_λ is singular with respect to Lebesgue measure, if λ is the reciprocal of a PV number. (An algebraic integer is a PV number provided all of its conjugates are less than one in modulus.) On the other hand, Wintner (1935) [7] proved that ν_λ is absolutely continuous for $\lambda = 2^{-\frac{1}{k}}$, for each $k \geq 1$, and Garsia (1962) [3] found some other algebraic integers for which ν_λ is absolutely continuous. Moreover, Erdős (1940) [2] also proved that there exists $a < 1$ such that the distribution ν_λ is absolutely continuous with respect to Lebesgue measure for (Lebesgue) a.e. $\lambda \in (a, 1)$. Then P. Erdős asked:

Is this statement true with $a = \frac{1}{2}$?

Received by the editors February 11, 1997.

1991 *Mathematics Subject Classification*. Primary 26A30, 28A78, 28A80.

Key words and phrases. Bernoulli convolution, equivalent measures.

The first author's research was supported by NSF Grant DMS-9502952. The second author's research was partially supported by grants F19099 and T19104 from the OTKA Foundation.

This exciting problem remained open for more than fifty years. Then Solomyak (1995) [6] gave a positive answer (see also Peres, Solomyak (1996a) [4] for a shorter proof). Namely,

Theorem 1 (Solomyak).

$$\nu_\lambda \ll m \text{ for Lebesgue a.e. } \lambda \in \left(\frac{1}{2}, 1\right),$$

where m is Lebesgue measure.

Answering a problem of the first author, posed to the Conference on Fractals and Stochastics (1994, Finsterbergen), we prove that that ν_λ is even equivalent to Lebesgue measure for a.e. $\lambda \in [\frac{1}{2}, 1]$. Using Solomyak's theorem, it is enough to prove that Lebesgue measure is either absolutely continuous or singular with respect to ν_λ for each λ . Actually we prove this statement for a more general family of measures. Furthermore, Peres, Solomyak (1996b) [5] proved that if the probabilities of choosing the signs $+$ and $-$ in Y_λ are $(p, 1-p)$ where $p \in [1/3, 2/3]$, then $\nu_\lambda \ll m$ holds for a.e. $\lambda \in [p^p(1-p)^{1-p}, 1]$. Using this, it follows from our result that even in this non-symmetric case the distributions are not only absolutely continuous but equivalent to Lebesgue measure for a.e. $\lambda \in [p^p(1-p)^{1-p}, 1]$. (For smaller λ the distributions are singular.)

We thank Yuval Peres for some useful conversations.

2. NOTATION

For an arbitrary $\lambda \in (\frac{1}{2}, 1)$ we define the 'projection' $\Pi_\lambda : \{-1, 1\}^{\mathbf{N}} \rightarrow [\frac{-1}{1-\lambda}, \frac{1}{1-\lambda}]$ by $\Pi_\lambda(\mathbf{i}) = \sum_{k=0}^{\infty} i_k \lambda^k$. Let μ be any Borel probability measure on $\{-1, 1\}^{\mathbf{N}}$ for which

$$(1) \quad \mu(B) > 0 \implies \mu\{(i, B)\} > 0$$

holds for all $B \subset \{-1, 1\}^{\mathbf{N}}$ and $i \in \{-1, 1\}$, where $(i, B) := \{(i, \mathbf{j}) \in \{-1, 1\}^{\mathbf{N}} : \mathbf{j} \in B\}$. For example μ may be any Bernoulli measure on $\{-1, 1\}^{\mathbf{N}}$ with probabilities $(p, 1-p)$, $0 < p < 1$. The 'push down measure' of μ is $\alpha_{\lambda, \mu}(B) := \mu(\Pi_\lambda^{-1}(B))$. We denote the interval $[\frac{-1}{1-\lambda}, \frac{1}{1-\lambda}]$ by I . Further, we define $S_i : I \rightarrow I$, $S_i(x) := \lambda x + i$ for $(i = -1, 1)$. The iterates of S_i are

$$S_{i_1 \dots i_n}(x) := S_{i_1} \circ \dots \circ S_{i_n}(x).$$

The image of I under $S_{i_1 \dots i_n}$ is called $I_{i_1 \dots i_n}$. The inverse of $S_{i_1 \dots i_n}$ is defined **only** on $I_{i_1 \dots i_n}$. So $S_{i_1 \dots i_n}^{-1}(A) := S_{i_1 \dots i_n}^{-1}(A \cap I_{i_1 \dots i_n})$. Then $S_i^{-1}(x) = \frac{1}{\lambda}x - \frac{i}{\lambda}$ for $x \in I_i$ ($i = -1, 1$). We denote the Lebesgue measure of a set A by $m(A)$.

3. THE THEOREM AND ITS CONSEQUENCES

Theorem 2. *Either $m \ll \alpha_{\lambda, \mu}$ or $m \perp \alpha_{\lambda, \mu}$.*

If μ is the Bernoulli measure with probabilities $(\frac{1}{2}, \frac{1}{2})$ then $\nu_\lambda = \alpha_{\lambda, \mu}$. Using Solomyak's Theorem, we obtain

Consequence 1. *For almost all $\lambda \in (\frac{1}{2}, 1)$, ν_λ is equivalent to Lebesgue measure.*

Clearly, any Bernoulli measure μ with probabilities $(p, 1-p)$, satisfies (1) (if $p \neq 0$). Thus,

Consequence 2. Let η_λ be the distribution of the random series $Z_\lambda = \sum \pm \lambda^n$, where the signs are chosen independently with probabilities $(p, 1 - p)$ and $0 < \lambda < 1$. Then either $m \ll \eta_\lambda$ or $m \perp \eta_\lambda$.

Let η_λ be as above. Then η_λ is singular for all $\lambda < p^p(1 - p)^{1-p}$ (see Peres, Solomyak (1996b) [5, Theorem 2 (a)]). Also Peres, Solomyak (1996b) [5, Corollary 1.4] proved that for $p \in [1/3, 2/3]$ and for a.e. $\lambda \in [p^p(1 - p)^{1-p}, 1]$, $\eta_\lambda \ll m$. Thus, using our previous consequence we obtain

Consequence 3. Let η_λ be the distribution of the random series $Z_\lambda = \sum \pm \lambda^n$, where the signs are chosen independently with probabilities $(p, 1 - p)$. Then for each $p \in [1/3, 2/3]$ and for almost every $\lambda \in [p^p(1 - p)^{1-p}, 1]$, the distribution η_λ is equivalent to Lebesgue measure.

4. LEMMAS AND PROOFS

To prove Theorem 2 we need two lemmas.

Lemma 1. Let $A \subset I$. Then $\alpha_{\lambda,\mu}(A) = 0 \implies \alpha_{\lambda,\mu}(S_i^{-1}(A)) = 0$ ($i = -1, 1$).

Proof. First observe that

$$(2) \quad \Pi_\lambda^{-1}(A) = \{(-1, \Pi_\lambda^{-1}(S_{-1}^{-1}(A)))\} \cup \{(1, \Pi_\lambda^{-1}(S_1^{-1}(A)))\}.$$

This is so, since for $i = -1, 1$

$$\begin{aligned} \mathbf{j} \in \Pi_\lambda^{-1}(S_i^{-1}(A)) &\iff \sum_{k=0}^\infty j_k \lambda^k \in S_i^{-1}(A) \iff \sum_{k=0}^\infty j_k \lambda^k \in \frac{1}{\lambda}A - \frac{i}{\lambda} \\ &\iff i + \sum_{k=0}^\infty j_k \lambda^{k+1} \in A \iff (i, \mathbf{j}) \in \Pi_\lambda^{-1}(A). \end{aligned}$$

To get a contradiction we assume that there exists a set A such that $\alpha_{\lambda,\mu}(A) = 0$ and $\alpha_{\lambda,\mu}(S_i^{-1}(A)) = \mu(\Pi_\lambda^{-1}(S_i^{-1}(A))) > 0$ holds for some $i \in \{-1, 1\}$.

Then from (1), it follows that $\mu((i, \Pi_\lambda^{-1}(S_i^{-1}(A)))) > 0$. Using (2), we find that $\mu(\Pi_\lambda^{-1}(A)) = \alpha_{\lambda,\mu}(A) > 0$. This contradiction proves our lemma. \square

Let $C \subset I$ be an arbitrary fixed Borel set. Let $C_0 := C$ and

$$C_{-(k+1)} := (S_{-1}^{-1}(C_{-k}) \cup S_1^{-1}(C_{-k})).$$

Then the ‘backward orbit’ of C in I is

$$(3) \quad \Lambda_- := \bigcup_{k \geq 0} C_{-k}.$$

Lemma 2. For any $C \subset I$, the set Λ_- defined above is either a set of zero measure or a full measure subset of I with respect to Lebesgue measure.

Proof. Let $\bar{\Lambda}_- := I \setminus \Lambda_-$. Obviously, it is enough to prove the statement of Lemma 2 for the set $\bar{\Lambda}_-$ instead of Λ_- . Observe that

$$(4) \quad x \in \bar{\Lambda}_- \implies S_i(x) \in \bar{\Lambda}_-$$

holds, since $S_i(x) \notin \bar{\Lambda}_- \implies \exists k \geq 0$ such that $S_i(x) \in C_{-k} \cap I_i$. Then $x = S_i^{-1}(S_i(x)) \in C_{-(k+1)} \subset \Lambda_-$. Iterate (4) to obtain

$$(5) \quad S_{i_1 \dots i_n}(\bar{\Lambda}_-) \subset \bar{\Lambda}_-,$$

for each $n \in \mathbf{N}$ and $(i_1, \dots, i_n) \in \{-1, 1\}^n$. Suppose that $m(\overline{\Lambda}_-) > 0$. Then $d := \frac{m(\overline{\Lambda}_-)}{|I|}$ is positive. Using (5), we obtain that $m(\overline{\Lambda}_- \cap I_{i_1 \dots i_n}) \geq m(S_{i_1 \dots i_n}(\overline{\Lambda}_-)) = \lambda^n \cdot d \cdot |I|$. Thus

$$(6) \quad \frac{m(\overline{\Lambda}_- \cap I_{i_1 \dots i_n})}{|I_{i_1 \dots i_n}|} \geq d$$

holds for each $i_1 \dots i_n$.

On the other hand, let $J \subset I$ be an arbitrary interval. Then we can find n and $i_1 \dots i_n$ such that $I_{i_1 \dots i_n} \subset J$ and

$$(7) \quad \frac{|I_{i_1 \dots i_n}|}{|J|} \geq \frac{\lambda}{3}.$$

Now, from (6) and (7) together, it follows that

$$\frac{m(\overline{\Lambda}_- \cap J)}{|J|} \geq d \cdot \frac{\lambda}{3}.$$

That is, Λ_- has no density point. Thus $\overline{\Lambda}_-$ is a full measure subset of I . This completes the proof of Lemma 2. \square

Proof of Theorem 2. Suppose that $m \not\ll \alpha_{\lambda, \mu}$. Then there is a set $C \subset I$ such that $m(C) > 0$ and $\alpha_{\lambda, \mu}(C) = 0$. Define Λ_- by (3). Then $m(\Lambda_-) > 0$; thus it follows from Lemma 2 that Λ_- is a full measure subset of I with respect to Lebesgue measure. On the other hand, Lemma 1 implies that $\alpha_{\lambda, \mu}(\Lambda_-) = 0$. So $m \perp \alpha_{\lambda, \mu}$. This completes the proof of Theorem 2. \square

REFERENCES

- [1] P.Erdős (1939). On a family of symmetric Bernoulli convolutions, *Amer. J. Math.* **61**, 974-976. MR **1**:52a
- [2] P.Erdős (1940). On the smoothness properties of a family of Bernoulli convolutions, *Amer. J. Math.* **62**, 180-186. MR **1**:139e
- [3] A.M. Garsia (1962). Arithmetic properties of Bernoulli convolutions, *Trans. Amer. Math. Soc.* **102**, 409-432. MR **25**:1409
- [4] Y. Peres and B. Solomyak (1996a). Absolute continuity of Bernoulli convolutions, *Math. Research Letters* **3**:2, 231-239. MR **97f**:28006
- [5] Y. Peres and B. Solomyak (1996b). Self-similar measures and intersections of Cantor sets, *Trans. Amer. Math. Soc.*, to appear.
- [6] B. Solomyak (1995). On the random series $\sum \pm \lambda^i$ (an Erdős problem), *Annals of Math.* **142**, 611-625. MR **97d**:11125
- [7] A. Wintner (1935). On convergent Poisson convolutions, *Amer. J. Math.* **57**, 827-838.

DEPARTMENT OF MATHEMATICS, P. O. BOX 305118, UNIVERSITY OF NORTH TEXAS, DENTON, TEXAS 76203-5118

E-mail address: mauldin@dynamics.math.unt.edu

Current address, K. Simon: Institute of Mathematics, University of Miskolc, Miskolc-Egyetemvaros, H-3515 Hungary

E-mail address: matsimon@gold.uni-miskolc.hu