

## THE MULTIDIMENSIONAL $p$ -ADIC GREEN FUNCTION

NGUYEN MINH CHUONG AND NGUYEN VAN CO

(Communicated by Dennis A. Hejhal)

ABSTRACT. A proof much simpler than the one given by Bikulov (*Investigation of the  $p$ -adic Green function*, Teoret. Mat. Fiz. **87** (1991), 376–390) for properties of the 2-dimensional  $p$ -adic Green function is shown. By this method one can treat a multidimensional case, and some sharp estimates are obtained.

### 1. INTRODUCTION

In recent years there has been a growing interest in  $p$ -adic analysis (see for instance [1]–[8]). In [8] by V. S. Vladimirov, P. V. Volovich and E. I. Zelenov and in [1] by A. Kh. Bikulov, some properties of 1-dimensional and 2-dimensional  $p$ -adic Green functions were studied.

In this note, a proof much simpler than the one in [1] is shown. In this way a multidimensional case with some sharp estimates is proved.

Let us recall here the definition of the  $p$ -adic Green function. As for the classical Green function on the field of real numbers (even on a Riemannian manifold), the  $p$ -adic Green function can be defined as the solution of the equation

$$(*) \quad (\Delta_p + m^2)G(x) = \delta(x),$$

where  $m$  is a positive real,  $\delta(x)$  is the Dirac function,  $\Delta_p$  is an operator defined by

$$(**) \quad (\Delta_p \varphi)(x) = \int_{Q_p^n} |(y, y)|_p \tilde{\varphi}(y) \chi((x, y)) dy,$$

where  $y = (y_1, \dots, y_n) \in Q_p^n$ ,  $|(y, y)|_p = |y_1^2 + \dots + y_n^2|_p$ ,  $|\cdot|_p$  is the  $p$ -adic norm ([7], [8]),  $(x, y) = x_1 y_1 + \dots + x_n y_n$ ,  $\varphi(x)$  is the test function on  $Q_p^n$  ( $\varphi(x) \in \mathcal{D}(Q_p^n)$ ),  $\tilde{\varphi}(y)$  denotes the  $p$ -adic Fourier transform of  $\varphi(x)$ ,  $\chi(x)$  is the additive character on  $Q_p$ , and  $dy$  denotes the Haar measure on  $Q_p^n$  (see [1], [2], [3], [8]).

By applying the  $p$ -adic Fourier transform to equality (\*), from equalities (\*), (\*\*) we get

$$(|(y, y)|_p + m^2) \tilde{G}(-y) = \tilde{\delta}(-y) = 1,$$

and the  $p$ -adic inverse Fourier transform gives an equivalent definition of the  $p$ -adic

---

Received by the editors July 1, 1997.

1991 *Mathematics Subject Classification*. Primary 11S80; Secondary 81T99.

This paper was supported by the National Natural Science Foundation of Vietnam.

Green function as follows:

$$G(x) = \int_{Q_p^n} \frac{\chi((x, y))dy}{|(y, y)|_p + m^2},$$

which we use in the present paper.

Note that even though the  $p$ -adic Green function can be defined in the usual way, its properties are essentially different from respective properties of the classical Green one. This statement can be easily seen by the theorems in the following sections.

### 2. 2-DIMENSIONAL CASE

In this section we study properties of the 2-dimensional  $p$ -adic Green function

$$G(x) = \int_{Q_p^2} \frac{\chi((x, y))dy}{|(y, y)|_p + m^2},$$

where  $m$  is a positive real.

If we set  $z = (x, x)$ , then  $|z|_p = |x_1^2 + x_2^2|_p$  and we obtain

**Theorem 1.** *The Green function  $G(x)$  has the following properties: For  $p \equiv 3 \pmod{4}$*

- i)  $G(x) = (1 - \frac{1}{p^2}) \sum_{k=0}^{+\infty} \frac{p^{-2k}}{m^2|z|_p + p^{-2k}} - \frac{1}{m^2|z|_p + p^2}$ , for  $z \neq 0$ ;
- ii)  $G(x) = G(|z|_p) > 0, \forall z \neq 0$ ;
- iii)

$$\begin{aligned} & \frac{1}{p^2} \ln \frac{m^2|z|_p + 1}{m^2|z|_p} + \frac{p^2 - 1}{p^2} \frac{1}{m^2|z|_p + 1} - \frac{1}{m^2|z|_p + p^2} \leq G(|z|_p) \\ & \leq \ln \frac{m^2|z|_p + 1}{m^2|z|_p} - \frac{1}{m^2|z|_p + p^2}. \end{aligned}$$

*Proof.* i) Since  $p \equiv 3 \pmod{4}$  we get  $|z|_p = |x_1^2 + x_2^2|_p = \max(|x_1|_p^2, |x_2|_p^2)$  (see [8]), so from  $x \neq 0$ , it implies that  $z \neq 0$  and

$$\begin{aligned} G(|z|_p) &= \sum_{\gamma=-\infty}^{+\infty} \int_{|y|_p=p^\gamma} \frac{1}{p^{2\gamma} + m^2} \chi((x, y))dy \\ &= \sum_{\gamma=-\infty}^{+\infty} \frac{1}{p^{2\gamma} + m^2} \cdot \begin{cases} p^{2\gamma}(1 - \frac{1}{p^2}), & \text{for } |x|_p \leq p^{-\gamma}, \\ -p^{2(\gamma-1)}, & \text{for } |x|_p = p^{-\gamma+1}, \\ 0, & \text{for } |x|_p \geq p^{-\gamma+2}. \end{cases} \end{aligned}$$

By  $x \neq 0$ , we set  $|x|_p = p^N$ , so  $|z|_p = p^{2N}$ , and

$$\begin{aligned} G(|z|_p) &= \sum_{\gamma=-\infty}^{+\infty} \frac{1}{p^{2\gamma} + m^2} \cdot \begin{cases} p^{2\gamma}(1 - \frac{1}{p^2}), & \text{for } \gamma \leq -N, \\ -p^{2(\gamma-1)}, & \text{for } \gamma = -N + 1, \\ 0, & \text{for } \gamma \geq -N + 2. \end{cases} \\ &= \sum_{\gamma=-\infty}^{-N} \frac{1}{p^{2\gamma} + m^2} p^{2\gamma}(1 - \frac{1}{p^2}) - \frac{p^{-2N}}{p^{2(-N+1)} + m^2}. \end{aligned}$$

Setting  $k = -\gamma - N$  and  $p^{2N} = |z|_p$  we get

$$G(|z|_p) = \left(1 - \frac{1}{p^2}\right) \sum_{k=0}^{+\infty} \frac{p^{-2k}}{m^2|z|_p + p^{-2k}} - \frac{1}{m^2|z|_p + p^2}.$$

□

ii) It is easy to see

$$G(|z|_p) = \sum_{k=0}^{+\infty} \frac{p^{-2k}}{m^2|z|_p + p^{-2k}} - \sum_{k=0}^{+\infty} \frac{p^{-2k-2}}{m^2|z|_p + p^{-2k}} - \frac{p^0}{m^2|z|_p + p^0 \cdot p^2}$$

or

$$\begin{aligned} G(|z|_p) &= \sum_{k=0}^{+\infty} p^{-2k} \left[ \frac{1}{m^2|z|_p + p^{-2k}} - \frac{1}{m^2|z|_p + p^{-2k}p^2} \right] \\ &= \sum_{k=0}^{+\infty} \frac{(p^2 - 1)p^{-4k}}{(m^2|z|_p + p^{-2k})(m^2|z|_p + p^{-2k}p^2)} > 0, \quad \forall z \neq 0. \end{aligned}$$

iii) Set  $m^2|z|_p = a > 0, p^{-2} = q$ . Obviously  $0 < q < 1$  and from i) of the theorem

$$\begin{aligned} (1) \quad G(|z|_p) &= \sum_{k=0}^{+\infty} \frac{1}{a + q^k} (q^k - q^{k+1}) - \frac{1}{a + p^2} \\ &= - \sum_{k=0}^{+\infty} \frac{1}{a + q^k} \int_k^{k+1} d(q^x) - \frac{1}{a + p^2}. \end{aligned}$$

By  $0 < q < 1$

$$- \int_k^{k+1} d(q^x) = - \int_k^{k+1} q^x \ln q dx > 0, \quad \frac{1}{a + q^k} \leq \frac{1}{a + q^x}, \quad \frac{1}{a + q^k} \geq \frac{1}{a + q^{x-1}}.$$

So

$$- \frac{1}{a + q^k} \int_k^{k+1} q^x \ln q dx \leq - \int_k^{k+1} \frac{q^x \ln q dx}{a + q^x}.$$

In view of (1) we get

$$\begin{aligned} G(|z|_p) &\leq - \sum_{k=0}^{\infty} \int_k^{k+1} \frac{q^x \ln q dx}{a + q^x} - \frac{1}{a + p^2} \\ &= \ln \frac{m^2|z|_p + 1}{m^2|z|_p} - \frac{1}{m^2|z|_p + p^2}, \end{aligned}$$

that is the second inequality in iii) is proved.

For proving the first inequality in iii) we do as follows:

$$\begin{aligned} G(|z|_p) &= (1 - q) \sum_{k=0}^{+\infty} \frac{q^k}{a + q^k} - \frac{1}{a + p^2} \\ &\geq \frac{1 - q}{a + 1} - \frac{1}{a + p^2} - q \int_0^{\infty} \frac{q^t \ln q dt}{a + q^t} \\ &= \frac{1 - p^{-2}}{m^2|z|_q + 1} - \frac{1}{m^2|z|_p + p^2} + \frac{1}{p^2} \ln \frac{a + 1}{a}. \quad \square \end{aligned}$$

*Remark.* The estimates in iii) can be made sharper as follows:

$$\begin{aligned}
 G(|z|_p) &= -\sum_{k=0}^{+\infty} \frac{1}{a+q^k} \int_k^{k+1} q^x \ln q dx - \frac{1}{a+p^2} \\
 &= \frac{1-q}{a+1} - \frac{1}{a+p^2} - \sum_{k=1}^{\infty} \frac{1}{a+q^k} \int_k^{k+1} q^x \ln q dx \\
 &< \frac{1-q}{a+1} - \frac{1}{a+p^2} - \sum_{k=1}^{\infty} \int_k^{k+1} \frac{q^x \ln q}{a+q^x} dx \\
 &= \frac{1-q}{a+1} - \frac{1}{a+p^2} - \int_1^{\infty} \frac{q^x \ln q dx}{a+q^x} \\
 &= \frac{1-p^{-2}}{a+1} - \frac{1}{a+p^2} + \ln \frac{a+p^{-2}}{a}.
 \end{aligned}$$

For the inequality  $>$ , obviously

$$G(|z|_p) > \frac{1-q}{a+1} + \frac{1-q}{a+q} - \frac{1}{a+p^2} + \frac{1}{p^2} \ln \frac{a+q}{a}.$$

**Theorem 2.** For  $p \equiv 1 \pmod{4}$ ,  $p$  is a prime number, we have

$$\begin{aligned}
 G(x) &= G(|z|_p) \\
 &= \sum_{n=0}^{+\infty} \left(1 - \frac{1}{p}\right) p^{-n} \left[ \frac{(n+1)\left(1 - \frac{1}{p}\right)}{p^{-n} + m^2|z|_p} - \frac{2}{p^{-n}p + m^2|z|_p} \right] + \frac{1}{p^2 + m^2|z|_p},
 \end{aligned}$$

where

$$G(x) = \iint_{Q_p^2} \frac{\chi((x, y)) dy}{|y_1^2 + y_2^2|_p + m^2}.$$

*Proof.* Since  $p \equiv 1 \pmod{4}$  there exists  $\tau \in Q_p$  such that  $\tau^2 = -1$ . Let us denote  $\tau = \sqrt{-1}$ . We get

$$y_1^2 + y_2^2 = (y_1 + \tau y_2)(y_1 - \tau y_2).$$

Setting

$$t = y_1 + \tau y_2, \quad \bar{t} = y_1 - \tau y_2$$

we have  $dy_1 dy_2 = dt d\bar{t}$  and  $(x, y) = at + \bar{a}\bar{t}$  with  $a = \frac{1}{2}(x_1 + \frac{1}{\tau}x_2)$ ,  $\bar{a} = \frac{1}{2}(x_1 - \frac{1}{\tau}x_2)$ .

So  $|a\bar{a}| = |z|_p$ , and

$$\begin{aligned}
 (2) \quad G(x) &= \int_{Q_p^2} \frac{\chi(at)\chi(\bar{a}\bar{t}) dt d\bar{t}}{|t|_p |\bar{t}|_p + m^2} \\
 &= \lim_{N \rightarrow +\infty} \int_{|\bar{t}| \leq p^N} \chi(\bar{a}\bar{t}) d\bar{t} \int_{|t|_p \leq p^N} \frac{1}{|t|_p |\bar{t}|_p + m^2} \chi(at) dt.
 \end{aligned}$$

Setting  $|a|_p = p^k$ , with sufficiently large  $N$  such that  $|a|_p > p^{-N}$ , i.e.  $-k < N$ , we obtain

$$\begin{aligned}
 I_1 &= \int_{|t|_p \leq p^N} \frac{1}{|t|_p |\bar{t}|_p + m^2} \chi(at) \\
 (3) \quad &= \sum_{\gamma=-\infty}^N \frac{1}{p^\gamma |\bar{t}|_p + m^2} \cdot \begin{cases} (1 - \frac{1}{p})p^\gamma, & \gamma \leq -k, \\ -p^{\gamma-1}, & \gamma = -k + 1, \\ 0, & \gamma \geq -k + 2, \end{cases} \\
 &= \sum_{\gamma=-\infty}^{-k} (1 - \frac{1}{p})p^\gamma \frac{1}{p^\gamma |\bar{t}|_p + m^2} - \frac{p^{-k}}{p^{-k+1} |\bar{t}|_p + m^2}.
 \end{aligned}$$

Obviously (2), (3) with  $|\bar{t}|_p = p^\beta$ ,  $|\bar{a}|_p = p^h > p^{-N}$  give

$$\begin{aligned}
 (4) \quad G(x) &= \sum_{\gamma=k}^{+\infty} \sum_{\beta=h}^{+\infty} (1 - \frac{1}{p})^2 \frac{p^{-(\gamma+\beta)}}{p^{-(\gamma+\beta)} + m^2} \\
 &+ \sum_{\gamma=k}^{+\infty} (\frac{1}{p} - 1) \frac{p^{-\gamma-h}}{p^{-\gamma-h+1} + m^2} \\
 &+ \sum_{k=h}^{+\infty} (\frac{1}{p} - 1) \frac{p^{-\beta-k}}{p^{-\beta-k+1} + m^2} + \frac{1}{p^2 + m^2 |z|_p}.
 \end{aligned}$$

It is not difficult to calculate the first term  $I_2$  in (4)

$$I_2 = \sum_{n=0}^{+\infty} (n+1) (1 - \frac{1}{p})^2 \frac{p^{-n}}{p^{-n} + m^2 |z|_p}$$

and to see that, in (4) the second term  $I_3$  is equal to the third term  $I_4$  and

$$I_3 = I_4 = \sum_{n=0}^{+\infty} (\frac{1}{p} - 1) \frac{p^{-n}}{pp^{-n} + m^2 |z|_p}.$$

Therefore

$$\begin{aligned}
 G(x) &= G(|z|_p) \\
 &= \sum_{n=0}^{+\infty} \left[ \frac{(n+1)(1 - \frac{1}{p})^2 p^{-n}}{p^{-n} + m^2 |z|_p} + \frac{2(\frac{1}{p} - 1)p^{-n}}{p^{-n}p + m^2 |z|_p} \right] + \frac{1}{p^2 + m^2 |z|_p}.
 \end{aligned}$$

□

### 3. $n$ -DIMENSIONAL CASE

We now study the properties of the  $n$ -dimensional  $p$ -adic Green function.

**Theorem 3.** *Let  $n$  be a positive integer and  $p$  be a prime number satisfying*

$$a_1^2 + a_2^2 + \dots + a_n^2 \not\equiv 0 \pmod{p}, \quad \forall a_1, a_2, \dots, a_n \in \{1, 2, \dots, p-1\}.$$

*Then the Green function defined by*

$$(5) \quad G(x) = \int_{Q_p^n} \frac{\chi((x, y)) dy}{|(y, y)|_p + m^2}$$

is calculated as follows:

$$G(x) = G(|z|_p) = \left( \sum_{k=0}^{+\infty} \frac{p^{-nk} \left(1 - \frac{1}{p^n}\right)}{m^2 |z|_p + p^{-2k}} - \frac{1}{m^2 |z|_p + p^2} \right) \frac{1}{|z|_p^{\frac{n-2}{2}}},$$

where  $z = x_1^2 + \dots + x_n^2$ ,  $|(y, y)|_p = |y_1^2 + \dots + y_n^2|_p, z \neq 0$ .

*Proof.* By the assumptions on  $p$ , if  $|y|_p = p^\gamma$ , then  $|(y, y)|_p = |y|_p^2 = p^{2\gamma}$ . Consequently

$$\begin{aligned} G(x) &= \sum_{\gamma=-\infty}^{+\infty} \int_{|y|_p=p^\gamma} \frac{\chi((x, y)) dy}{p^{2\gamma} + m^2} \\ &= \sum_{\gamma=-\infty}^{+\infty} \frac{1}{p^{2\gamma} + m^2} \cdot \begin{cases} p^{n\gamma} \left(1 - \frac{1}{p^n}\right), & \text{for } \gamma \leq -N \text{ (} |x|_p = p^N \text{)}, \\ -p^{n(\gamma-1)}, & \text{for } \gamma = -N + 1, \\ 0, & \text{for } \gamma \geq -N + 2, \end{cases} \\ &= \sum_{\gamma=-\infty}^{-N} \frac{p^{n\gamma} \left(1 - \frac{1}{p^n}\right)}{p^{2\gamma} + m^2} - \frac{1}{p^{2(-N+1)} + m^2} p^{-nN} \\ &= \left(1 - \frac{1}{p^n}\right) \frac{1}{|z|_p^{\frac{n-2}{2}}} \sum_{k=0}^{+\infty} \frac{p^{-nk}}{m^2 |z|_p + p^{-2k}} - \frac{1}{|z|_p^{\frac{n-2}{2}} (p^2 + m^2 |z|_p)}. \end{aligned}$$

□

**Theorem 4.** Consider the same assumptions as in Theorem 3. Then the Green function  $G(|z|_p)$  defined by (5) has the following properties:

- a)  $G(|z|_p) > 0, \forall z \neq 0$ ;
- b)

$$G(|z|_p) = \frac{p^2 - 1}{1 - p^{-(n+2)}} \cdot \frac{1}{m^4 |z|_p^{1+\frac{n}{2}}} + O\left(\frac{1}{|z|_p^{1+\frac{n}{2}}}\right)$$

as  $m^2 |z|_p \rightarrow +\infty$ ;

- c)

$$\frac{1}{|z|_p^{\frac{n-2}{2}}} \left( \frac{1-q}{a+1} - \frac{1}{a+p^2} + qI_n \right) < G(|z|_p) < \frac{1}{|z|_p^{\frac{n-2}{2}}} \left( -\frac{1}{a+p^2} + I_n \right),$$

where  $a = m^2 |z|_p > 0, q = \frac{1}{p^n}$ ,

$$(6a) \quad I_n = n \left[ \frac{1}{n-2} - \frac{a}{n-4} + \frac{a^2}{n-6} - \dots + \frac{(-1)^{k-1} a^{k-1}}{2} \ln \frac{a+1}{a} \right],$$

if  $n = 2k, k$  is an integer  $\geq 1$  and

$$(6b) \quad I_n = n \left[ \frac{1}{n-2} - \frac{a}{n-4} + \frac{a^2}{n-6} - \dots + (-1)^{k-1} a^{k-1} \frac{1}{\sqrt{a}} \arctan \frac{1}{\sqrt{a}} \right],$$

if  $n = 2k - 1, k$  is an integer  $\geq 1$  (for  $k = 1$  there are no  $n - 2j$  terms in  $I_n$ ).

*Proof.* a) From Theorem 3 it follows that

$$\begin{aligned} G(|z|_p) &= \frac{1}{|z|_p^{\frac{n-2}{2}}} \left[ \sum_{k=0}^{+\infty} \frac{p^{-nk}(1-p^{-n})}{m^2|z|_p + p^{-2k}} - \frac{1}{m^2|z|_p + p^2} \right] \\ &= \frac{1}{|z|_p^{\frac{n-2}{2}}} \left[ \sum_{k=0}^{+\infty} \frac{p^{-nk}}{m^2|z|_p + p^{-2k}} - \sum_{k=1}^{+\infty} \frac{p^{-nk}}{m^2|z|_p + p^2p^{-2k}} - \frac{p^{-n \cdot 0}}{m^2|z|_p + p^2p^{-2 \cdot 0}} \right]. \end{aligned}$$

Consequently

$$\begin{aligned} (7) \quad G(|z|_p) &= \frac{1}{|z|_p^{\frac{n-2}{2}}} \sum_{k=0}^{+\infty} \left[ \frac{p^{-nk}}{m^2|z|_p + p^{-2k}} - \frac{p^{-nk}}{m^2|z|_p + p^2p^{-2k}} \right] \\ &= \frac{1}{|z|_p^{\frac{n-2}{2}}} \sum_{k=0}^{+\infty} (p^2 - 1) \frac{p^{-(n+2)k}}{(m^2|z|_p + p^{-2k})(m^2|z|_p + p^2p^{-2k})} > 0. \end{aligned}$$

b) Setting  $t = \frac{1}{m^2|z|_p}$  we have

$$\begin{aligned} (8) \quad \frac{1}{m^2|z|_p + p^{-2k}} &= t \left( \frac{1}{1 + p^{-2k}t} \right) = t(1 - p^{-2k}t + O(t)) \\ &= \frac{1}{m^2|z|_p} - \frac{p^{-2k}}{m^4|z|_p^2} + O\left(\frac{1}{|z|_p^2}\right), \end{aligned}$$

as  $m^2|z|_p \rightarrow +\infty$ .

Similarly

$$(9) \quad \frac{1}{m^2|z|_p + p^2p^{-2k}} = \frac{1}{m^2|z|_p} - \frac{p^2p^{-2k}}{m^4|z|_p^2} + O\left(\frac{1}{|z|_p^2}\right).$$

By (8), (9) and (7)

$$\begin{aligned} G(|z|_p) &= \frac{1}{|z|_p^{\frac{n-2}{2}}} \sum_{k=0}^{+\infty} p^{-nk} \left[ -\frac{p^{-2k}}{m^4|z|_p^2} + \frac{p^2p^{-2k}}{m^4|z|_p^2} + O\left(\frac{1}{|z|_p^2}\right) \right] \\ &= \frac{1}{|z|_p^{\frac{n-2}{2}}} \sum_{k=0}^{+\infty} \frac{(p^2 - 1)p^{-(n+2)k}}{m^4|z|_p^2} + O\left(\frac{1}{|z|_p^{1+\frac{n}{2}}}\right) \\ &= \frac{1}{|z|_p^{\frac{n-2}{2}}} \cdot \frac{p^2 - 1}{1 - p^{-(n+2)}} \cdot \frac{1}{m^4|z|_p^2} + O\left(\frac{1}{|z|_p^{1+\frac{n}{2}}}\right) \\ &= \frac{p^2 - 1}{1 - p^{-(n+2)}} \cdot \frac{1}{m^4|z|_p^{1+\frac{n}{2}}} + O\left(\frac{1}{|z|_p^{1+\frac{n}{2}}}\right). \end{aligned}$$

c) By Theorem 3

$$(10) \quad G(|z|_p) = \frac{1}{|z|_p^{\frac{n-2}{2}}} \left[ \sum_{k=0}^{+\infty} \frac{p^{-nk}(1-p^{-n})}{m^2|z|_p + p^{-2k}} - \frac{1}{m^2|z|_p + p^2} \right].$$

Setting  $a = m^2|z|_p > 0$ ,  $q = p^{-n}$  we get

$$\begin{aligned}
 (11) \quad s &= \sum_{k=0}^{+\infty} \frac{p^{-nk}(1-p^{-n})}{m^2|z|_p + p^{-2k}} = \sum_{k=0}^{+\infty} (1-q) \frac{q^k}{a + q^{\frac{2k}{n}}} \\
 &= \sum_{k=0}^{+\infty} \frac{1}{a + q^{\frac{2k}{n}}} (q^k - q^{k+1}) \\
 &= - \sum_{k=0}^{+\infty} \frac{1}{a + q^{\frac{2k}{n}}} \int_k^{k+1} d(q^x).
 \end{aligned}$$

By  $k \leq x \leq k+1$ ,  $0 < q < 1$

$$\frac{1}{a + q^{\frac{2(x-1)}{n}}} \leq \frac{1}{a + q^{\frac{2k}{n}}} \leq \frac{1}{a + q^{\frac{2x}{n}}}.$$

But

$$- \int_k^{k+1} d(q^x) > 0,$$

consequently

$$(12) \quad - \int_k^{k+1} \frac{d(q^x)}{a + q^{\frac{2(x-1)}{n}}} < - \frac{1}{a + q^{\frac{2k}{n}}} \int_k^{k+1} d(q^x) < - \int_k^{k+1} \frac{d(q^x)}{a + q^{\frac{2x}{n}}}.$$

By (11), (12) it is easy to see

$$(13) \quad \frac{1-q}{a+1} - \sum_{k=1}^{+\infty} \int_k^{k+1} \frac{d(q^x)}{a + q^{\frac{2(x-1)}{n}}} < s < - \sum_{k=0}^{+\infty} \int_k^{k+1} \frac{d(q^x)}{a + q^{\frac{2x}{n}}}.$$

From (13) it follows that

$$(14) \quad \frac{1-q}{a+1} - q \int_0^{+\infty} \frac{d(q^x)}{a + q^{\frac{2x}{n}}} < s < - \int_0^{+\infty} \frac{d(q^x)}{a + q^{\frac{2x}{n}}}.$$

Setting

$$I_n = - \int_0^{+\infty} \frac{d(q^x)}{a + q^{\frac{2x}{n}}}$$

we obtain

$$I_n = - \int_1^0 \frac{nt^{n-1} dt}{a + t^2} = n \int_0^1 \frac{t^{n-1} dt}{a + t^2}.$$

Let  $n \geq 3$ . Then

$$I_n = n \int_0^1 \left( t^{n-3} - a \frac{t^{n-3}}{a + t^2} \right) dt = n \left( \frac{1}{n-2} - a I_{n-2} \right).$$

By induction it is not difficult to prove

$$(15a) \quad I_n = n \left[ \frac{1}{n-2} - \frac{a}{n-4} + \frac{a^2}{n-6} - \dots + \frac{(-1)^{k-1} a^{k-1}}{2} \ln \frac{a+1}{a} \right],$$

if  $n = 2k$

$$(15b) \quad I_n = n \left[ \frac{1}{n-2} - \frac{a}{n-4} + \frac{a^2}{n-6} - \dots + \frac{(-1)^{k-1} a^{k-1}}{\sqrt{a}} \arctan \frac{1}{\sqrt{a}} \right],$$

if  $n = 2k - 1$ ,

$k \geq 1$ ; for  $k = 1$  it is assumed there are no  $n - 2j$  terms in  $I_n$ .

From (15a), (15b) and (10) with

$$G(|z|_p) = \frac{1}{|z|_p^{\frac{n-2}{2}}} \left[ s - \frac{1}{a+p^2} \right]$$

we get the estimates

$$\frac{1}{|z|_p^{\frac{n-2}{2}}} \left[ \frac{1-q}{a+1} - \frac{1}{a+p^2} + qI_n \right] < G(|z|_p) < \frac{1}{|z|_p^{\frac{n-2}{2}}} \left[ -\frac{1}{a+p^2} + I_n \right].$$

□

**Corollary 1.** For  $n = 1$  we obtain

$$\begin{aligned} (1 - \frac{1}{p}) \frac{\sqrt{|z|_p}}{m^2|z|_p + 1} - \frac{\sqrt{|z|_p}}{m^2|z|_p + p^2} + \frac{1}{mp} \arctan \frac{1}{m\sqrt{|z|_p}} &< G(|z|_p) \\ &< -\frac{\sqrt{|z|_p}}{m^2|z|_p + p^2} + \frac{1}{m} \arctan \frac{1}{m\sqrt{|z|_p}}. \end{aligned}$$

**Corollary 2.** For  $n = 2$  we have the estimates *iii*) in Theorem 1.

ACKNOWLEDGMENTS

The authors would like to thank the referee and the editor for helpful suggestions by which the paper is improved.

REFERENCES

- [1] A. Kh. Bikulov, *Investigation of the  $p$ -adic Green function*, Teoret. Mat. Fiz. (in Russian), **87** (1991), 376-390. MR **92g**:81202
- [2] Nguyen Minh Chuong, Nguyen Van Co and Tran Quoc Binh,  *$p$ -adic distributions and Sobolev spaces*, Preprint 96/10, Hanoi Institute of Mathematics.
- [3] Nguyen Minh Chuong and Nguyen Van Co, *A new proof for  $p$ -adic Green function*, Preprint 1997, Hanoi Institute of Mathematics.
- [4] Nguyen Minh Chuong, Nguyen Minh Tri and Le Quang Trung, *Theory of Partial Differential Equations*, Sci. & Techn. Publ. House, Hanoi (1995).
- [5] B. M. Dwork,  *$p$ -adic bound up regular singular points*, Duke. Math. J. **62**(1991), 689-720. MR **93c**:12009
- [6] A. Yu. Khrennikov,  *$p$ -adic valued distributions in mathematical physics*, Kluwer Academic Publishers, Dordrecht, Boston, London (1994), MR **96c**:46080

- [7] N. I. Koblitz, *p-adic numbers, p-adic analysis and zeta functions*, Springer, New-York (1984), MR **86c**:11086
- [8] V. S. Vladimirov, I. V. Volovich, E. I. Zelenov, *p-adic analysis and mathematical physics* (Russian), Nauka, Moscow (1994), MR **96f**:11156

HANOI INSTITUTE OF MATHEMATICS, P.O. BOX 631 BOHO, 10 000 HANOI, VIETNAM  
*E-mail address*: [nmchuong@thevinh.ac.vn](mailto:nmchuong@thevinh.ac.vn)

DEPARTMENT OF MATHEMATICS, HANOI STATE UNIVERSITY, MAI DICH, TU LIEM, HANOI, VIETNAM