

**A FIXED POINT THEOREM
AND ITS APPLICATION TO INTEGRAL EQUATIONS
IN MODULAR FUNCTION SPACES**

A. AIT TALEB AND E. HANEHALY

(Communicated by Dale Alspach)

ABSTRACT. In this paper we present a fixed point theorem of Banach type in modular space. We give an application of this result to a nonlinear integral equation in Musielak-Orlicz space.

0. INTRODUCTION

It is well known that one of the standard proofs of Banach's fixed point theorem is based on Cantor's theorem in complete metric spaces [3, 4]. To this end, using some convenient constants in the contraction assumption, we present a generalization of Banach's fixed point theorem in some classes of modular spaces, where the modular is s -convex, having the Fatou property and satisfying the Δ_2 -condition.

As an application we study the existence of a solution for an integral equation of Lipschitz type in a Musielak-Orlicz space.

We begin by recalling some basic concepts of modular spaces; for more information, we refer to the books by Musielak [8] and Kozłowski [7].

Definition 0-1. Let X be an arbitrary vector space over K ($= \mathbf{R}$ or \mathbf{C}).

a) A functional $\rho : X \rightarrow [0, +\infty]$ is called modular if:

i) $\rho(x) = 0 \iff x = 0$.

ii) $\rho(\alpha x) = \rho(x)$ for $\alpha \in K$ with $|\alpha| = 1$, $\forall x \in X$.

iii) $\rho(\alpha x + \beta y) \leq \rho(x) + \rho(y)$ if $\alpha, \beta \geq 0, \alpha + \beta = 1, \forall x, y \in X$.

b) If iii) is replaced by:

iii') $\rho(\alpha x + \beta y) \leq \alpha^s \rho(x) + \beta^s \rho(y)$ for $\alpha, \beta \geq 0, \alpha^s + \beta^s = 1$ with an $s \in]0, 1]$, then the modular ρ is called an s -convex modular; and if $s = 1$, ρ is called convex modular.

c) A modular ρ defines a corresponding modular space, i.e. the space X_ρ given by

$$X_\rho = \{x \in X \mid \rho(\lambda x) \rightarrow 0 \text{ as } \lambda \rightarrow 0\}.$$

Remarks. 1) Note that in general there is no reason to expect the subadditivity of a modular ρ . Nevertheless, in view of iii) from Definition 0-1 the inequality $\rho(x + y) \leq \rho(2x) + \rho(2y)$ holds.

Received by the editors January 5, 1996 and, in revised form, October 30, 1997.

1991 *Mathematics Subject Classification.* Primary 46A80, 47H10, 45G10, 46E30.

Key words and phrases. Modular space, fixed point, integral equation.

2) If ρ is convex modular, the modular space X_ρ can be equipped with a norm called the Luxemburg norm defined by:

$$|x|_\rho = \inf\{\alpha > 0; \rho(\frac{x}{\alpha}) \leq 1\}.$$

3) As a classical example, we would like to mention the Musielak-Orlicz space denoted by L^φ [8] and the modular function space denoted by L^ρ [7].

Definition 0-2. Let X_ρ be a modular space.

- a) A sequence $(x_n)_n$ in X_ρ is said to be:
 - i) ρ -convergent to x if $\rho(x_n - x) \rightarrow 0$ as $n \rightarrow +\infty$.
 - ii) ρ -Cauchy if $\rho(x_n - x_m) \rightarrow 0$ as n and $m \rightarrow +\infty$.
- b) X_ρ is ρ -complete if any ρ -Cauchy sequence is ρ -convergent.
- c) A subset $B \subset X_\rho$ is said to be ρ -closed if for any sequence $(x_n)_n \subset B$ with $x_n \rightarrow x$, then $x \in B$. \bar{B}^ρ denotes the closure of B in the sense of ρ .
- d) A subset $B \subset X_\rho$ is called ρ -bounded if:

$$\delta_\rho(B) = \sup_{x,y \in B} \rho(x - y) < +\infty,$$
 where $\delta_\rho(B)$ is called the ρ -diameter of B .
- e) We say that ρ has the Fatou property if:

$$\rho(x - y) \leq \liminf \rho(x_n - y_n)$$

whenever

$$x_n \xrightarrow{\rho} x \text{ and } y_n \xrightarrow{\rho} y.$$

f) ρ is said to satisfy the Δ_2 -condition if: $\rho(2x_n) \rightarrow 0$ as $n \rightarrow +\infty$ whenever $\rho(x_n) \rightarrow 0$ as $n \rightarrow +\infty$.

I. FIXED POINT THEOREM

I-1. Theorem I-1. Let X_ρ be a ρ -complete modular space. Assume that ρ is an s -convex modular satisfying the Δ_2 -condition and having the Fatou property. Let B be a ρ -closed subset of X_ρ and $T : B \rightarrow B$ a mapping such that:

$$(*) \quad \exists c, k \in \mathbf{R}^+ : c > \max(1, k), \rho(c(Tx - Ty)) \leq k^s \rho(x - y) \quad \forall x, y \in B.$$

Then T has a fixed point.

Remarks. 1) It is natural to introduce the constants c and k in the assumption of strict contraction in modular spaces. Note also that Theorem I-1 and its proof become more simple in the particular case where $s = 1$ (ρ is convex) and $c = 2 > k > 0$; see [1].

2) The contraction (*) in Theorem I-1 is also true for any constant c_0 such that $1 < c_0 \leq c$:

$$\begin{aligned} \rho(c_0(Tx - Ty)) &= \rho\left(\frac{c_0}{c}c(Tx - Ty)\right) \\ &\leq \frac{c_0^s}{c^s}\rho(c(Tx - Ty)) \leq k_0^s \rho(x - y) \end{aligned}$$

where $k_0 = \frac{c_0}{c}k < c_0$, since $\frac{k}{c} < 1$.

I-2. Proof of Theorem I-1. 1st step: The proof of Theorem I-1 is based on the next result which is the modular formulation of Cantor’s theorem:

Theorem I-2. *Let X_ρ be a ρ -complete modular space. Let $(F_n)_n$ be a decreasing sequence of nonempty, ρ -closed subset of X_ρ with $\delta_\rho(F_n) \rightarrow 0$ as $n \rightarrow +\infty$. Then $\bigcap_n F_n$ is reduced to one point.*

The proof of this result uses the same ideas as in complete metric spaces and it suffices to replace the distance by the modular ρ .

2nd step. Let $(\epsilon_n)_n$ be a decreasing sequence of positive numbers such that $\epsilon_n \rightarrow 0$ as $n \rightarrow +\infty$, and consider the sets defined by:

$$M_{\epsilon_n} = \{x \in B | \rho(L(x - Tx)) \leq \epsilon_n\},$$

where $L = \max\{c, 2\alpha\}$ and α is the s -conjugate of c , i.e. $\frac{1}{c^s} + \frac{1}{\alpha^s} = 1$. Then the sequence (M_{ϵ_n}) has the following property:

1) $M_{\epsilon_n} \neq \emptyset, \forall n$.

We assume without any loss of generality that: $\exists x \in B$ such that $\rho(x - Tx) < +\infty$; then for $p \in \mathbf{N}^*$ we have:

$$\begin{aligned} \rho(c(T^{p+1}x - T^p x)) &\leq k^s \rho(T^p x - T^{p-1}x) \\ &= k^s \rho\left(\frac{1}{c}(T^p x - T^{p-1}x)\right) \\ &\leq \left(\frac{k^s}{c^s}\right) \rho(T^{p-1}x - T^{p-2}x), \end{aligned}$$

and by induction we deduce:

$$\rho(c(T^{p+1}x - T^p x)) \leq \left(\frac{k}{c}\right)^{s(p-1)} \rho(x - Tx).$$

Since $\frac{k}{c} < 1$, we have $\rho(c(T^{p+1}x - T^p x)) \rightarrow 0$ as $p \rightarrow +\infty$. Thus by (Δ_2)

$$\rho(L(T^{p+1}x - T^p x)) \rightarrow 0 \quad \text{as } p \rightarrow +\infty.$$

Hence: $\exists q \in \mathbf{N}^*$ such that $\rho(L(T^{q+1}x - T^q x)) \leq \epsilon_n$. Then: $y = T^q x \in M_{\epsilon_n}$.

2) M_{ϵ_n} is ρ -closed.

Let (x_p) be a sequence in M_{ϵ_n} . Assume that $(x_p)_p$ is ρ -convergent to $x \in X_\rho$, $(x_p)_p \subset B$ and B is ρ -closed; it follows that $x \in B$.

$\rho(x_p - x) \rightarrow 0$ as $p \rightarrow +\infty$, by (Δ_2) , $\rho(L(x_p - x)) \rightarrow 0$ as $p \rightarrow +\infty$.

On the other hand $\rho(c(Tx_p - Tx)) \leq k^s \rho(x_p - x) \leq k^s \rho(L(x_p - x))$; then

$$\rho(c(Tx_p - Tx)) \rightarrow 0 \quad \text{as } p \rightarrow +\infty.$$

Again by (Δ_2) $\rho(L(Tx_p - Tx)) \rightarrow 0$ as $p \rightarrow +\infty$. The Fatou property implies that:

$$\rho(L(Tx - x)) \leq \liminf \rho(L(Tx_p - x_p)) \leq \epsilon_n.$$

Therefore $x \in M_{\epsilon_n}$ and hence M_{ϵ_n} is ρ -closed.

3) $\delta_\rho(M_{\epsilon_n}) \rightarrow 0$ as $n \rightarrow +\infty$.

Let $x, y \in M_{\epsilon_n}$; we have:

$$\rho(x - y) = \rho\left(\frac{\alpha(x - Tx)}{\alpha} + \frac{c(Tx - Ty)}{c} + \frac{\alpha(Ty - y)}{\alpha}\right)$$

where

$$\begin{aligned}\alpha^s &= \frac{c^s}{c^s - 1} \\ &\leq \frac{1}{\alpha^s} \rho(\alpha(x - Tx) + \alpha(Ty - y)) + \frac{1}{c^s} \rho(c(Tx - Ty)) \\ &\leq \frac{1}{2^s \alpha^s} \rho(2\alpha(x - Tx)) + \frac{1}{2^s \alpha^s} \rho(2\alpha(Ty - y)) + \frac{1}{c^s} \rho(c(Tx - Ty)) \\ &\leq \frac{1}{2^s \alpha^s} (\rho(L(x - Tx)) + \rho(L(Ty - y))) + \frac{1}{c^s} \rho(c(Tx - Ty)).\end{aligned}$$

Then

$$\rho(x - y) \leq \frac{1}{2^s \alpha^s} 2\epsilon_n + \frac{k^s}{c^s} \rho(x - y).$$

Hence,

$$\rho(x - y) \leq \frac{1}{2^s \alpha^s} 2\epsilon_n \frac{c^s}{c^s - k^s}$$

and we deduce that

$$\delta_\rho(M_{\epsilon_n}) = \sup_{x, y \in M_{\epsilon_n}} \rho(x - y) \leq \epsilon_n 2^{1-s} \frac{c^s - 1}{c^s - k^s},$$

and

$$\delta_\rho(M_{\epsilon_n}) \rightarrow 0 \text{ as } n \rightarrow +\infty.$$

4) (M_{ϵ_n}) is decreasing. This is an immediate consequence of the fact that (ϵ_n) is decreasing.

It follows that the conditions of Cantor's theorem are satisfied and hence we have: $\bigcap_n M_{\epsilon_n} = \{x_0\}$.

But $x_0 \in M_{\epsilon_n} \forall n \Rightarrow \rho(L(x_0 - Tx_0)) \leq \epsilon_n \rightarrow 0$ as $n \rightarrow +\infty \Rightarrow Tx_0 = x_0$.

Remark. We are unable to prove whether the conclusion of Theorem I-1 is true if we have $c = 1$ and $0 < k < 1$. To this end, recall the following results by Khamsi-Kozłowski-Reich:

Theorem I-3 ([5]). *Let ρ be a modular function satisfying the Δ_2 -condition and let B be a $|\cdot|_\rho$ -closed subset of L_ρ .*

$T : B \rightarrow B$ is a mapping such that: $\exists 0 < k < 1$ such that

$$\rho(Tf - Tg) \leq k\rho(f - g) \quad \forall f, g \in B.$$

Then: T has a fixed point if one of the following assumptions is satisfied:

i) $\exists f_0 \in B; \sup_n (2T^n f_0) < \infty$.

ii) B is ρ -bounded.

If *ii)* is satisfied, the fixed point is unique. Note also that the modular ρ in Theorem I-3 has the Fatou property and the Δ_2 -condition as in Theorem I-1. Also by (Δ_2) we have: B is ρ -closed $\Leftrightarrow B$ is $|\cdot|_\rho$ -closed.

On the other hand the strict inequality (*) in Theorem I-1 implies the inequality of Theorem I-3

$$\rho(Tx - Ty) = \rho\left(\frac{1}{c}(c(Tx - Ty))\right) \leq \frac{1}{c^s} \rho(c(Tx - Ty)) \leq \frac{k^s}{c^s} \rho(x - y).$$

Consequently with some reinforced assumptions in Theorem I-1, namely the s -convexity of ρ and the strict contraction (*), we prove the existence of a fixed point

for T without restrictive conditions concerning the domain of T : i) or ii) in Theorem I-3 .

The following example shows that Theorem I-1 can be more appropriate for applications.

II. APPLICATION

II-1. General frame. Consider the following integral equation:

$$(I) \quad u(t) = \exp(-t)f + \int_0^t \exp(s - t)Tu(s)ds,$$

where:

- i) $T : B \rightarrow B$ with B a ρ -closed, convex subset of a Musielak-Orlicz space L^φ satisfying the Δ_2 -condition.
- ii) T is ρ -Lipschitz:

$$\exists \gamma > 0, \rho(Tu - Tv) \leq \gamma\rho(u - v) \quad u, v \in B.$$

- iii) $f \in B$.

Theorem II-1. *Under these conditions, for all $A > 0$ the integral equation (I) has a solution $u \in C^\varphi = C([0, A], L^\varphi)$. C^φ is the modular space of continuous mappings from $[0, A]$ into L^φ*

By iterative techniques, Khamsi [6] has shown this result under supplementary conditions: B is ρ -bounded and T is ρ -Lipschitz with constant $\gamma = 1$.

To delete all restrictive assumptions on the Lipschitz constant γ we introduce the space C^φ equipped with a convenient modular.

Our method follows the standard technique of the resolution of differential equations or integral equations of Lipschitz type in Banach spaces. See Deimling [2, p. 39].

II-2. Functional frame: The modular space C^φ .

Definition 2-1. $u : I \rightarrow L^\varphi$, where $I = [0, A]$, is said to be continuous at $t_0 \in I$ if: for $t_n \in I$, $t_n \rightarrow t_0$ as $n \rightarrow +\infty \Rightarrow \rho(u(t_n) - u(t_0)) \rightarrow 0$ as $n \rightarrow +\infty$.

Since ρ satisfies the Δ_2 -condition, it is equivalent to:

$$t_n \rightarrow t_0 \text{ as } n \rightarrow +\infty \Rightarrow |u(t_n) - u(t_0)|_\rho \rightarrow 0 \text{ as } n \rightarrow +\infty.$$

Let $C^\varphi = C(I, L^\varphi)$ be the space of all continuous mappings from $I = [0, A]$ into L^φ .

Proposition 2-1. *Suppose that the modular ρ satisfies (Δ_2) , and $B \subset L^\varphi$ is a ρ -closed, convex subset of L^φ . For $a \geq 0$ let $\rho_a(u) = \sup_{t \in I} \exp(-at)\rho(u(t))$ for $u \in C^\varphi$. Then:*

- 1) (C^φ, ρ_a) is a modular space, and ρ_a is a convex modular satisfying the Fatou property and the Δ_2 -condition.
- 2) C^φ is ρ_a -complete.
- 3) $C_0^\varphi = C(I, B)$ is a ρ_a -closed, convex subset of C^φ .

Proof. 1) i) C^φ is a real vector space.

Let $u, v \in C^\varphi, t_0 \in I$. Then for $t_n \in I$ such that $t_n \rightarrow t_0$ as $n \rightarrow +\infty$ we have:

$$\rho\left(\frac{(u+v)(t_n) - (u+v)(t_0)}{2}\right) \leq \frac{1}{2}\rho(u(t_n) - u(t_0)) + \frac{1}{2}\rho(v(t_n) - v(t_0)).$$

Hence, $\rho\left(\frac{(u+v)(t_n)-(u+v)(t_0)}{2}\right) \rightarrow 0$ as $n \rightarrow +\infty$. And by Δ_2 :

$$\rho((u+v)(t_n) - (u+v)(t_0)) \rightarrow 0 \text{ as } n \rightarrow +\infty,$$

which implies that $(u+v)$ is continuous at t_0 .

Again by Δ_2 $\rho(\lambda(u(t_n) - u(t_0))) \rightarrow 0$ as $n \rightarrow +\infty$ for all $\lambda \in \mathbf{R}$; then λu is also continuous at t_0 .

ii) ρ_a is well defined.

Since ρ satisfies the (Δ_2) -condition, the domain of ρ is $\{f \in L^\varphi, \rho(f) < +\infty\} = L^\varphi$, and since ρ is convex and $|\cdot|_\rho$ -continuous at 0 it follows that ρ is $|\cdot|_\rho$ -continuous on L^φ . See Zeidler [9, p. 383].

Consequently for all $u \in C^\varphi$, $\rho_a(u)$ has a meaning.

iii) ρ_a is a convex modular.

This is a simple consequence of the fact that ρ is a convex modular.

iv) ρ_a satisfies the Fatou property.

Let $(u_n)_n$ (resp $(v_n)_n$) be a sequence in C^φ , ρ_a -convergent to u (resp. to v) $\in C^\varphi$. Then $\forall t \in I$, $\rho(u_n(t) - u(t)) \rightarrow 0$ as $n \rightarrow +\infty$ and $\rho(v_n(t) - v(t)) \rightarrow 0$ as $n \rightarrow +\infty$.

Since ρ satisfies the Fatou property, we have:

$$\rho(u(t) - v(t)) \leq \liminf \rho(u_n(t) - v_n(t)) \quad \forall t \in I,$$

$$\exp(-at)\rho(u(t) - v(t)) \leq \liminf \exp(-at)\rho(u_n(t) - v_n(t)) \leq \liminf \rho_a(u_n - v_n)$$

and then

$$\rho_a(u - v) \leq \liminf \rho_a(u_n - v_n).$$

v) ρ_a has the (Δ_2) -condition.

Since ρ satisfies the (Δ_2) -condition, one has:

$$\begin{aligned} \rho_a(u_n) \rightarrow 0 \text{ as } n \rightarrow +\infty &\Leftrightarrow \forall t \in I, \exp(-at)\rho(u_n(t)) \rightarrow 0 \text{ as } n \rightarrow +\infty \\ &\Leftrightarrow \forall t \in I, \exp(-at)\rho(2u_n(t)) \rightarrow 0 \text{ as } n \rightarrow +\infty \\ &\Leftrightarrow \rho_a(2u_n) \rightarrow 0 \text{ as } n \rightarrow +\infty. \end{aligned}$$

2) It is known that $(L^\varphi, |\cdot|_\rho)$ is a Banach space. Then the space $(C^\varphi, |\cdot|_{\rho_a})$ is also a Banach space. If $(u_n)_n$ is a ρ_a -Cauchy sequence in C^φ , by the Δ_2 -condition it is $|\cdot|_{\rho_a}$ -Cauchy. Hence: $\exists u \in C^\varphi$ such that $u_n \rightarrow u$ in $(C^\varphi, |\cdot|_{\rho_a})$.

Consequently $(u_n) \rightarrow u$ in (C^φ, ρ_a) . It follows that C^φ is ρ_a -complete.

3) $C_0^\varphi = C(I, B)$ is convex and ρ_a -closed.

The convexity of C_0^φ and its ρ_a -closedness are clearly obtained by the convexity and ρ -closedness of B in L^φ .

II-3. Proof of Theorem II-1. Define the operator S over C_0^φ by:

$$\forall u \in C_0^\varphi, \quad Su(t) = \exp(-t) f + \int_0^t \exp(s-t) Tu(s) ds, \quad \forall t \in I.$$

1st step: We show that $S : C_0^\varphi \rightarrow C_0^\varphi$.

i) Su is continuous from I into $(L^\varphi, |\cdot|_\rho)$.

Let $t_n, t_0 \in I$ with $t_n \rightarrow t_0$ as $n \rightarrow +\infty$.

T is ρ -Lipschitz $\rho(Tu(t_n) - Tu(t_0)) \leq \gamma\rho(u(t_n) - u(t_0))$. Since u is ρ -continuous, Tu is ρ -continuous at t_0 and by (Δ_2) , Tu is $|\cdot|_\rho$ -continuous at t_0 . Hence Su is $|\cdot|_\rho$ -continuous at t_0 .

ii) $Su(t) \in B, \quad \forall t \in I.$

It is well known that in the Banach space $(L^\varphi, |\cdot|_\rho)$

$$\int_0^t \exp(s-t) Tu(s) ds \in \left(\int_0^t \exp(s-t) ds \right) \overline{\text{co}}\{Tu(s), 0 \leq s \leq t\} \\ \in (1 - \exp(-t)) \overline{\text{co}}B,$$

where $\overline{\text{co}}B$ is the closed convex hull of B in $(L^\varphi, |\cdot|_\rho)$. But B is convex and ρ -closed; thus, $\overline{\text{co}}B = \overline{B} \subset \overline{B^\rho} = B$. Hence, $Su(t) \in \exp(-t) B + (1 - \exp(-t)) B \subseteq B \quad \forall t \in I$.

2nd step: For $u, v \in C_0^\varphi$ and $\lambda > 0$ we have:

$$\lambda(Su(t) - Sv(t)) = \int_0^t \lambda \exp(s-t) (Tu(s) - Tv(s)) ds.$$

Lemma 3-1 ([6]). *Let $x \in C^\varphi$ and $0 < \lambda \leq \frac{\exp A}{\exp(A)-1}$. Then:*

$$\rho\left(\int_0^t \lambda \exp(s-t) x(s) ds\right) \leq \lambda \frac{\exp(at) - \exp(-t)}{1+a} \rho_a(x).$$

Note that in [6] this result was been shown with $\lambda = 1$ and $a = 0$.

Proof. Let $T = \{t_0, t_1, t_2, \dots, t_n\}$ be any subdivision of $[0, t]$.

$$\sum_{i=0}^{n-1} \lambda(t_{i+1} - t_i) \exp(t_i - t) x(t_i)$$

is $|\cdot|_\rho$ -convergent, and consequently, ρ -convergent to $\int_0^t \lambda \exp(s-t) x(s) ds$ in L^φ when $|T| = \sup\{|t_{i+1} - t_i|, i = 0, \dots, n-1\} \rightarrow 0$ as $n \rightarrow +\infty$.

By Fatou we have:

$$\rho\left(\int_0^t \lambda \exp(s-t) x(s) ds\right) \leq \liminf \rho\left(\sum_{i=0}^{n-1} \lambda(t_{i+1} - t_i) \exp(t_i - t) x(t_i)\right).$$

On the other hand:

$$\sum_{i=0}^{n-1} \lambda(t_{i+1} - t_i) \exp(t_i - t) \leq \int_0^t \lambda \exp(s-t) ds \\ = (1 - \exp(-t))\lambda \leq \lambda(1 - \exp(-A)).$$

Since $0 < \lambda < \frac{\exp(A)}{\exp(A)-1}$, it follows that $\sum_{i=0}^{n-1} \lambda(t_{i+1} - t_i) \exp(t_i - t) \leq 1$, and the convexity of ρ implies:

$$\rho\left(\sum_{i=0}^{n-1} \lambda(t_{i+1} - t_i) \exp(t_i - t) x(t_i)\right) \leq \sum_{i=0}^{n-1} \lambda(t_{i+1} - t_i) \exp(t_i - t) \rho(x(t_i)) \\ = \sum_{i=0}^{n-1} \lambda(t_{i+1} - t_i) \exp(t_i - t) \exp(at_i) \exp(-at_i) \rho(x(t_i)) \\ \leq \sum_{i=0}^{n-1} \lambda(t_{i+1} - t_i) \exp((1+a)t_i - t) \rho_a(x) \\ \leq \left(\int_0^t \lambda \exp((1+a)s - t) ds\right) \rho_a(x).$$

Then:

$$\rho\left(\int_0^t \lambda \exp(s-t) x(s) ds\right) \leq \lambda \frac{\exp(at) - \exp(-t)}{1+a} \rho_a(x).$$

3rd step:

$$\rho(\lambda(Su(t) - Sv(t))) \leq \lambda \frac{\exp(at) - \exp(-t)}{1+a} \rho_a(Tu - Tv).$$

But

$$\rho_a(Tu - Tv) = \sup_{t \in I} \exp(-at) \rho(Tu(t) - Tv(t)) \leq \gamma \rho_a(u - v).$$

Then:

$$\begin{aligned} \exp(-at) \rho(\lambda(Su(t) - Sv(t))) &\leq \lambda \frac{\gamma}{1+a} (1 - \exp((-t)(1+a))) \rho_a(u - v) \\ &\leq \lambda \frac{\gamma}{1+a} (1 - \exp((-A)(1+a))) \rho_a(u - v), \quad \forall t \in I. \end{aligned}$$

Hence

$$\rho_a(\lambda(Su - Sv)) \leq \lambda \frac{\gamma}{1+a} (1 - \exp((-A)(1+a))) \rho_a(u - v).$$

We consider $\lambda, 1 < \lambda \leq \frac{\exp(A)}{\exp(A)-1}$. Then S has a fixed point if:

$$\lambda \frac{\gamma}{1+a} (1 - \exp((-A)(1+a))) < \lambda \Leftrightarrow \frac{\gamma}{1+a} (1 - \exp((-A)(1+a))) < 1.$$

The last inequality is satisfied if we take for example $a \geq \gamma$.

In the end, by Theorem I-1, S has a fixed point which is a solution of the integral equation (I).

REFERENCES

- [1] A. Ait Taleb, Points fixes et Applications aux equations integrales dans les espaces modulaires. These de 3me cycle, Departement de Mathematiques et Informatique, Rabat (1996).
- [2] K. Deimling, Nonlinear Functional Analysis, Springer Verlag, (1985). MR **86j**:47001
- [3] K. Goebel-S. Reich, Uniform convexity, Hyperbolic Geometry and nonexpansive mappings, Dekker, (1984). MR **86d**:58012
- [4] K. Goebel-W.A. Kirk, Topics in metric fixed point theory, Cambridge University Press, Cambridge (1990). MR **92c**:47070
- [5] M.A. Khamsi-W.M. Kozłowski-S. Reich, Fixed point theory in modular Function spaces. Nonlinear Analysis, theory, methods and Applications. vol.14. N^o 11 (1990). 935-953. MR **91d**:47042
- [6] M.A. Khamsi, Nonlinear semigroups in modular Function space thèse d' état. Departement de Mathematiques, Rabat (1994). MR **93g**:47085
- [7] W.M. Kozłowski, Modular function spaces. Dekker (1988). CMP 98:02
- [8] J. Musielak, Orlicz spaces and modular spaces, Lecture notes in Mathematics, vol 1034, Springer Verlag (1983). MR **85m**:46028
- [9] E. Zeidler, Nonlinear Functional Analysis and its Applications tome III, Springer Verlag (1985). MR **90b**:49005

DEPARTMENT OF MATHEMATICS AND INFORMATICS, FACULTE DES SCIENCES, UNIVERSITE MOHAMMED V, BP. 1014, RABAT, MOROCCO