

FINITE-DIMENSIONAL LEFT IDEALS IN SOME ALGEBRAS ASSOCIATED WITH A LOCALLY COMPACT GROUP

M. FILALI

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. Let G be a locally compact group, let $L^1(G)$ be its group algebra, let $M(G)$ be its usual measure algebra, let $L^1(G)^{**}$ be the second dual of $L^1(G)$ with an Arens product, and let $LUC(G)^*$ be the conjugate of the space $LUC(G)$ of bounded, left uniformly continuous, complex-valued functions on G with an Arens-type product. We find all the finite-dimensional left ideals of these algebras. We deduce that such ideals exist in $L^1(G)$ and $M(G)$ if and only if G is compact, and in $L^1(G)^{**}$ (except those generated by right annihilators of $L^1(G)^{**}$) and $LUC(G)^*$ if and only if G is amenable.

1. INTRODUCTION

Let G be a locally compact group, $L^1(G)$ be its group algebra, and $M(G)$ be its usual measure algebra. Other Banach algebras (usually larger than $L^1(G)$ and $M(G)$) can also be associated with G . For instance, the second dual $L^1(G)^{**}$ of $L^1(G)$ is a Banach algebra with an Arens product. One can also consider the space $LUC(G)$ of the bounded left uniformly continuous functions on G , or the space $WAP(G)$ of the weakly almost periodic functions on G . An Arens-type product can be introduced into the conjugate of each of these spaces of functions and make them into Banach algebras. Let \mathcal{A} denote any of these algebras. Our main concern in this paper is with the finite-dimensional left ideals in \mathcal{A} . We start in Theorem 1 by giving examples of such ideals. These examples are obtained with the help of the U -invariant elements of \mathcal{A}^n where U is a continuous and bounded representation of G on \mathbb{C}^n . This notion has already been introduced in earlier papers with $n = 1$. In [3], the so-called χ -invariant elements (where χ is a character of G) were introduced in order to determine the minimal left ideals of these algebras when G is abelian. In [4], these type of elements were referred to as λ -invariant elements with $\lambda \in \mathbb{T}$, and were used to solve some linear equations in $\ell^\infty(\mathbb{Z})^*$, then to determine the finite-dimensional left ideals in this case. In [5], the U -invariance was used to study the minimal ideals in these algebras. In Theorems 2 and 3, we determine all the finite-dimensional left ideals in \mathcal{A} and show that they are in general of the form given in Theorem 1. This is a generalization of the result obtained in [4, 2.7(a)]. We deduce in the corollary which follows that such ideals exist in $LUC(G)^*$ and in $L^1(G)^{**}$ (apart from those generated by right annihilators of $L^1(G)^{**}$) if and only

Received by the editors October 28, 1997.

1991 *Mathematics Subject Classification*. Primary 43A10, 22D15.

Key words and phrases. Locally compact group, Arens product, representation, amenable, U -invariant, finite-dimensional left ideal.

if G is amenable. In $L^1(G)$ and $M(G)$, the ideals of finite-dimension exist if and only if G is compact.

2. PRELIMINARIES

Let G be a locally compact group with a left Haar measure λ , and a Haar modular function Δ . For measurable functions f and ϕ on G and for $s \in G$, we write

$$f * \phi(s) = \int_G f(t)\phi(t^{-1}s) d\lambda(t)$$

whenever the integral exists. We shall be concerned with the following Banach algebras which are related to G . We begin by recalling, of course, the most intimate ones: the group algebra $L^1(G)$ and the measure algebra $M(G)$. The first one is the Banach algebra of all measurable complex-valued functions ϕ on G satisfying

$$\int_G |\phi(s)| d\lambda(s) < \infty.$$

The product of two elements ϕ and ψ in $L^1(G)$ is $\phi * \psi$. The second one is the Banach algebra of all bounded, regular, Borel measures on G . By the Riesz representation theorem (see [8, Chapter 3]), we shall regard a measure in $M(G)$ as an element of $C_0(G)^*$, where $C_0(G)$ is the space of continuous complex-valued functions on G vanishing at infinity. The product of two elements μ and ν of $M(G)$ is given then by

$$(\mu\nu)(f) = \int_G \int_G f(st) d\mu(s)d\nu(t) = \int_G \int_G f(st) d\nu(t)d\mu(s) \quad \text{for } f \in C_0(G).$$

Furthermore, for each $\phi \in L^1(G)$, we define $\lambda_\phi \in M(G)$ by

$$\lambda_\phi(f) = \int_G f(s)\phi(s) d\lambda(s) \quad \text{for } f \in C_0(G).$$

Then under the map $\phi \rightarrow \lambda_\phi$, we will regard $L^1(G)$ as a subalgebra of $M(G)$. In fact, $L^1(G)$ is a closed two-sided ideal of $M(G)$ (see for example [8, Theorem 19.18]). The other algebras which we shall consider are defined in the following way. Let $L^\infty(G)$ be the Banach space of all measurable complex-valued functions that are bounded almost everywhere with respect to λ , and for each $\phi \in L^\infty(G)$, let $\hat{\phi}$ be the function defined on G by $\hat{\phi}(s) = \Delta(s^{-1})\phi(s^{-1})$. The Banach space $L^1(G)^{**}$ becomes a Banach algebra with the first Arens product. This product is obtained by first letting

$$f_\nu(\phi) = \nu(\hat{\phi} * f) \quad \text{for all } \nu \in L^1(G)^{**}, f \in L^\infty(G) \text{ and } \phi \in L^1(G).$$

Then, for μ and ν in $L^1(G)^{**}$,

$$(\mu\nu)(f) = \mu(f_\nu) \quad \text{for all } f \in L^\infty(G).$$

Let $C(G)$ denote the space of all bounded, complex-valued, continuous functions on G . The left translate of a function f on G by $s \in G$ is defined by $f_s(t) = f(st)$ for all $t \in G$. Let $LUC(G)$ be the space of left uniformly continuous functions in $C(G)$, i.e.,

$$LUC(G) = \{f \in C(G) : s \mapsto f_s : G \rightarrow C(G) \text{ is norm continuous}\}.$$

Then $LUC(G)^*$ is also a Banach algebra under the product

$$\begin{aligned}(\mu\nu)(f) &= \mu(f_\nu) \quad \text{for all } f \in LUC(G), \quad \text{where} \\ f_\nu(s) &= \nu(f_s) \quad \text{for all } s \in G\end{aligned}$$

(the function f_ν is easily seen to be in $LUC(G)$). Note that the product in $M(G)$ (and so in $L^1(G)$) is defined in the same way with $C_0(G)$ instead of $LUC(G)$.

More generally, one can start with a norm closed, conjugate closed subspace F of $C(G)$ containing the constant functions and having the property that the functions f_s and f_μ are in F for all $f \in F$, $s \in G$ and $\mu \in F^*$ (the functions f_s and f_μ are defined as earlier in $LUC(G)$). Following [2, Definition 2.2.10], such an F is said to be *admissible*; the space F^* also becomes a Banach algebra under the product

$$(\mu\nu)(f) = \mu(f_\nu) \quad \text{for all } f \in F.$$

For more details, the reader is directed to [2, pages 72-78]. As we have already seen, $LUC(G)$ is admissible. Other examples are the space $WAP(G)$ of weakly almost periodic functions on G , and the space $AP(G)$ of almost periodic functions on G . These spaces are

$$\begin{aligned}WAP(G) &= \{f \in C(G) : f_G \text{ is weakly relatively compact}\}, \\ AP(G) &= \{f \in C(G) : f_G \text{ is norm relatively compact}\}, \quad \text{where} \\ f_G &= \{f_s : s \in G\}.\end{aligned}$$

Let \mathcal{A} denote each of the Banach algebras $L^1(G)$, $M(G)$, $L^1(G)^{**}$, and F^* , where F is an admissible subspace of $C(G)$ with $AP(G)F \subseteq F$. Apart from $L^1(G)$, the algebra \mathcal{A} is the dual of some Banach space of functions of G , which we shall denote by \mathcal{F} . In the case of $L^1(G)$, we let $\mathcal{F} = C_0(G)$. When $\mathcal{A} = F^*$ or $M(G)$, the group G may be embedded continuously into \mathcal{A} by the mapping $e : G \rightarrow \mathcal{A}$ which is defined by

$$e(s)(f) = f(s) \quad \text{for all } s \in G \text{ and } f \in \mathcal{F}.$$

We recall that an element μ of \mathcal{A} is *left invariant* if

$$\mu(f_s) = \mu(f) \quad \text{for all } f \in \mathcal{F} \text{ and } s \in G.$$

When $L^1(G) * \mathcal{F} \subseteq \mathcal{F}$, we say that $\mu \in \mathcal{A}$ is *topologically left invariant* if

$$\mu(\phi * f) = \left(\int_G \phi(s) d\lambda(s) \right) \mu(f) \quad \text{for all } \phi \in L^1(G) \text{ and } f \in \mathcal{F}.$$

The notions of left invariance and topological left invariance are equivalent when $\mathcal{F} \subseteq LUC(G)$. But this is not so if $\mathcal{F} = L^\infty(G)$. We say that \mathcal{F} is *amenable* if there is a non-zero left invariant (or equivalently a topologically left invariant) element in \mathcal{A} . When G is a compact topological group, $C_0(G) = C(G)$ is amenable since $\lambda \in C(G)^*$. The spaces $WAP(G)$ and $AP(G)$ are always amenable. But this is not so for $LUC(G)$ and $L^\infty(G)$; for example, when G is the free group on two generators (see [2, Example 3.4(e)]). So we say that the group G is *amenable* if $L^\infty(G)$, or equivalently $LUC(G)$, is amenable. See [2] or [7].

We shall also need representations of G on \mathbb{C}^n . Recall that a *representation* of G on a Hilbert space H is a homomorphism of G into the semigroup of bounded operators on H . We say that U is *continuous* when the function $s \mapsto U(s)\bar{x}$ is continuous on G for each $\bar{x} \in H$. We say that U is *irreducible* when $\{0\}$ and H are the only invariant (closed) subspaces under all $U(s)$, i.e., there is no (closed)

subspace E other than $\{0\}$ and H satisfying $U(s)E \subseteq E$ for all $s \in G$. We also recall that there is a one-to-one correspondence between the representations of $L^1(G)$ on H and those of G on H . This is given by the formula

$$\langle U(\phi)\bar{x}, \bar{y} \rangle = \int_G \langle U(s)\bar{x}, \bar{y} \rangle \phi(s) d\lambda(s) \quad \phi \in L^1(G), \bar{x}, \bar{y} \in H;$$

see [8, Section 22]. Since we shall be concerned solely with representations U on $H = \mathbb{C}^n$, we fix a basis $\{\bar{x}_1, \bar{x}_2, \dots, \bar{x}_n\}$ for \mathbb{C}^n , and correspond to U the matrix representation $U = (u_{ij})_{i,j=1}^n$, where u_{ij} are the coordinate functions defined on G by

$$u_{ij}(s) = \langle U(s)\bar{x}_i, \bar{y}_j \rangle.$$

Note that the letter U is used to denote the representations of G , $L^1(G)$, and their corresponding matrices, but we promise the reader that this will cause no confusion.

3. LEFT IDEALS OF FINITE-DIMENSION

The notion of χ -invariance, where χ is a continuous character of G was introduced in [3] to determine the minimal left ideals of \mathcal{A} when G is abelian. In this section, we generalize this notion by defining the U -invariant vectors of \mathcal{A}^n , where U is a representation of G on \mathbb{C}^n . The U -invariance is essential to determine the finite-dimensional left ideals of these algebras. Some of the arguments in Theorems 1 and 2 have already been in [1]; they are, however, simplified here with the help of the U -invariance. Before we state our main definition, we introduce the following notations.

Notations. Let $\mu \in \mathcal{A}$, $f \in \mathcal{F}$, $\bar{\mu} = (\mu_i)_{i=1}^n$ be a column vector in \mathcal{A}^n , and A be an $n \times n$ matrix with entries a_{ij} ($i, j = 1, \dots, n$) in \mathcal{F} . Then we write $\mu(A) = (\mu(a_{ij}))_{i,j=1}^n$, $\bar{\mu}(f)$ is the column vector $(\mu_i(f))_{i=1}^n$ of \mathbb{C}^n , $(A)\bar{\mu}$ is the column vector $(\sum_{j=1}^n \mu_j(a_{ij}))_{i=1}^n$ of \mathbb{C}^n (note that this is obtained with the matrix multiplication relative to the product given by the duality between \mathcal{A} and \mathcal{F}), and $\mu\bar{\mu}$ is the column vector $(\mu\mu_i)_{i=1}^n$ of \mathcal{A}^n . Furthermore, when A and B are $n \times n$ matrices whose entries are measurable functions on G then, whenever the integrals exist, we write

$$\begin{aligned} A \cdot B &= \int_G A(t)B(t) d\lambda(t), \\ A * B(s) &= \int_G A(t)B(t^{-1}s) d\lambda(t), \\ A\tilde{*}B(s) &= \int_G B(t^{-1}s)A(t) d\lambda(t). \end{aligned}$$

Definition. We say that a vector $\bar{\mu}$ of \mathcal{A}^n is U -invariant if there exists a continuous and bounded representation U of G on \mathbb{C}^n such that

$$\bar{\mu}(f_s) = U(s^{-1})\bar{\mu}(f) \quad \text{for each } f \in \mathcal{F} \text{ and } s \in G.$$

When $\mathcal{A} = L^1(G)^{**}$, we say that $\bar{\mu}$ be topologically U -invariant if

$$\bar{\mu}(\phi * f) = U(\phi)\bar{\mu}(f) \quad \text{for each } f \in L^\infty(G) \text{ and } \phi \in L^1(G).$$

Let \tilde{U} be defined on G by $\tilde{U}(s) = U(s^{-1})$. A direct computation leads to the following lemma. We omit the proof.

Lemma 1. Let A be an $n \times n$ matrix with entries in \mathcal{F} , and for $s \in G$, let A_s be the matrix whose entries are the left translates of those of A by s . Let Φ be an $n \times n$ matrix with entries in $L^1(G)$.

- (1) If $\bar{\mu} \in \mathcal{A}^n$ is U -invariant, then $(A_s)\bar{\mu} = (A\tilde{U}(s))\bar{\mu}$.
- (2) If $A = L^1(G)^{**}$ and $\bar{\mu} \in \mathcal{A}^n$ is topologically U -invariant, then $(\Phi * A)\bar{\mu} = (\Phi \cdot A(\cdot)U)\bar{\mu}$ and $(\Phi \bar{*} A)\bar{\mu} = (A(\Phi \cdot U))\bar{\mu}$, where $\Phi \cdot A(\cdot)U$ is the matrix-valued function defined on G by $\Phi \cdot A(s)U = \int_G \Phi(t)A(s)U(t) d\lambda(t)$.

Lemma 2. Let U be a continuous and bounded representation of G on \mathbb{C}^n , and let I be the identity representation. Let $\bar{\mu}$ and $\bar{\nu}$ be in \mathcal{A}^n such that $\bar{\mu}(f) = (fU)\bar{\nu}$ for all $f \in \mathcal{F}$; or equivalently, $\bar{\nu}(f) = (f\tilde{U})\bar{\mu}$ for all $f \in \mathcal{F}$. Then

- (1) $\bar{\nu}$ is I -invariant if and only if $\bar{\mu}$ is U -invariant,
- (2) when $\mathcal{A} = L^1(G)^{**}$, $\bar{\nu}$ is topologically I -invariant if and only if $\bar{\mu}$ is topologically U -invariant.

Remark. Observe that $(fU)\bar{\nu}$ and $(f\tilde{U})\bar{\mu}$ are well defined in the lemma. This is due to the fact that the coordinate functions of U and \tilde{U} are almost periodic (which is easy to verify, or see [1, Lemma 1]) and the extra assumption that $AP(G)F \subseteq F$.

Proof of Lemma 2. Let $\bar{\nu} \in \mathcal{A}^n$ be I -invariant, $s \in G$ and $f \in \mathcal{F}$. Then, by Lemma 1,

$$\bar{\mu}(f_s) = (f_s U)\bar{\nu} = (\tilde{U}(s)f_s U_s)\bar{\nu} = \tilde{U}(s)(fU)\bar{\nu} = \tilde{U}(s)\bar{\mu}(f).$$

So $\bar{\mu}$ is U -invariant. For the converse, let $\bar{\mu}$ be U -invariant, $s \in G$ and $f \in \mathcal{F}$. Then

$$\bar{\nu}(f_s) = (f_s \tilde{U})\bar{\mu} = (f_s \tilde{U}_s U(s))\bar{\mu} = (f \tilde{U} U(s) \tilde{U}(s))\bar{\mu} = (f \tilde{U})\bar{\mu} = \bar{\nu}(f).$$

So $\bar{\nu}$ is I -invariant.

Statement (2) follows with the help of statement (2) of Lemma 1. Let $\bar{\nu}$ be topologically I -invariant, $\phi \in L^1(G)$ and $f \in L^\infty(G)$. We remark first that

$$\begin{aligned} (\phi * f)U(s) &= U(s) \int_G \phi(t)f(t^{-1}s) d\lambda(t) \\ &= \int_G U(t)U(t^{-1}s)\phi(t)f(t^{-1}s) d\lambda(t) = (\phi U) * (fU)(s). \end{aligned}$$

Therefore,

$$\begin{aligned} \bar{\mu}(\phi * f) &= ((\phi * f)U)\bar{\nu} = ((\phi U) * (fU))\bar{\nu} = ((\phi U) \cdot (fU)(\cdot)I)\bar{\nu} \\ &= \left(\int_G U(s)\phi(s) d\lambda(s) \right) (fU)\bar{\nu} = U(\phi)\bar{\mu}(f), \end{aligned}$$

and so $\bar{\mu}$ is topologically U -invariant.

For the converse, let $\bar{\mu}$ be topologically U -invariant, $f \in L^\infty(G)$ and $\phi \in L^1(G)$. Then, for each $s \in G$,

$$\begin{aligned} (\phi * f)\tilde{U}(s) &= \tilde{U}(s) \int_G \phi(t)f(t^{-1}s) d\lambda(t) \\ &= \int_G \tilde{U}(t^{-1}s)\tilde{U}(t)\phi(t)f(t^{-1}s) d\lambda(t) = (\phi \tilde{U})\bar{*}(f\tilde{U})(s), \end{aligned}$$

and so

$$\begin{aligned} \bar{\nu}(\phi * f) &= ((\phi * f) \tilde{U}) \bar{\mu} = ((\phi \tilde{U}) \tilde{*} (f \tilde{U})) \bar{\mu} = ((f \tilde{U})((\phi \tilde{U}) \cdot U)) \bar{\mu} \\ &= \left((f \tilde{U}) \int_G \phi(s) \tilde{U}(s) U(s) d\lambda(s) \right) \bar{\mu} \\ &= \left(\int_G \phi(s) d\lambda(s) \right) (f \tilde{U}) \bar{\mu} = \left(\int_G \phi(s) d\lambda(s) \right) \bar{\nu}(f). \end{aligned}$$

So $\bar{\nu}$ is topologically I -invariant. □

Theorem 1. *Let G be a locally compact group. Let $\bar{\mu} \in \mathcal{A}^n$ and M be the linear span of the coordinates $\mu_1, \mu_2, \dots, \mu_n$ of $\bar{\mu}$. Let U be a continuous and bounded representation of G on \mathbb{C}^n . Then M is a left ideal of \mathcal{A} of dimension less or equal to n in each of the following cases:*

- (1) F is amenable, $\mathcal{A} = F^*$ and $\bar{\mu}$ is U -invariant.
- (2) G is amenable, $\mathcal{A} = L^1(G)^{**}$ and $\bar{\mu}$ is topologically U -invariant.
- (3) G is compact, $\mathcal{A} = L^1(G)$ or $M(G)$, and $\bar{\mu}$ is U -invariant.

Furthermore, M is minimal and of dimension n when U is irreducible.

Proof. We consider only the first two statements. The proof of statement (3) is similar. Let $\mathcal{A} = F^*$, and let $\bar{\mu}$ be U -invariant. That M is a left ideal follows directly from the lemma above. Let μ be arbitrary in F^* and $f \in F$. Then $(\mu \bar{\mu})(f) = \mu(f_{\bar{\mu}})$, where $f_{\bar{\mu}}(s) = \bar{\mu}(f_s) = \tilde{U}(s)\bar{\mu}(f)$, and so

$$\mu \bar{\mu}(f) = \mu(\tilde{U}\bar{\mu}(f)) = \mu(\tilde{U})\bar{\mu}(f).$$

Thus, $\mu \bar{\mu} = \mu(\tilde{U})\bar{\mu}$, which means obviously that M is a left ideal of F^* .

For statement (2), let $\mu \in L^1(G)^{**}$ and $f \in L^\infty(G)$. Then $(\mu \bar{\mu})(f) = \mu(f_{\bar{\mu}})$, where

$$\begin{aligned} f_{\bar{\mu}}(\phi) &= \bar{\mu}(\hat{\phi} * f) = U(\hat{\phi})\bar{\mu}(f), \text{ and} \\ U(\hat{\phi}) &= \int_G U(s)\hat{\phi}(s) d\lambda(s) = \int_G \tilde{U}(s)\phi(s) d\lambda(s) = \tilde{U}(\phi) \text{ for all } \phi \in L^1(G). \end{aligned}$$

Thus, $(\mu \bar{\mu})(f) = \mu(f_{\bar{\mu}}) = \mu(\tilde{U})\bar{\mu}$, and so M is a left ideal of $L^1(G)^{**}$. That M is of dimension less or equal to n is clear in each case.

Suppose now that U is irreducible, and let us prove that M is minimal. We start with the algebra F^* . Let $\mu \in M$ be arbitrary, and write $\mu = \sum_{i=1}^n x_i \mu_i = \bar{\mu} \underline{x} = \bar{x} \underline{\mu}$, where $\underline{x} = (x_1, x_2, \dots, x_n)$ is a non-zero vector of \mathbb{C}^n . For each vector $\bar{y} \in \mathbb{C}^n$, we can find s_1, s_2, \dots, s_k in G and $\alpha_1, \alpha_2, \dots, \alpha_k$ in \mathbb{C} such that

$$\sum_{i=1}^n \alpha_i \tilde{U}(s_i) \bar{x} = \bar{y}$$

since \tilde{U} is also irreducible. It follows that

$$\begin{aligned} \sum_{i=1}^n \alpha_i e(s_i) \mu &= \sum_{i=1}^n \alpha_i e(s_i) (\bar{\mu} \underline{x}) = \sum_{i=1}^n \alpha_i (e(s_i) \bar{\mu}) \underline{x} \\ &= \sum_{i=1}^n \alpha_i (\tilde{U}(s_i) \bar{\mu}) \underline{x} = \sum_{i=1}^n \alpha_i (\tilde{U}(s_i) \bar{x}) \underline{\mu} = \bar{y} \underline{\mu} \end{aligned}$$

(remember that $e(s) \in F^*$ and $e(s)(f) = f(s)$ for $s \in G$ and $f \in F$). This means first that $F^* \mu = M$, and so M is minimal. Secondly, if $\bar{y} \in \mathbb{C}^n$ is such that $\bar{y}\underline{\mu} \neq \bar{0}$, then this argument shows also that

$$\sum_{i=1}^n \alpha_i e(s_i) \bar{\mu} \underline{x} = \bar{y} \underline{\mu} \neq 0,$$

and implies that $\bar{\mu} \underline{x} \neq 0$. So the elements $\mu_1, \mu_2, \dots, \mu_n$ are linearly independent and M is of dimension n .

In $L^1(G)^{**}$, the corresponding representation U of $L^1(G)$ is also irreducible, and so we can find, for each $\bar{y} \in \mathbb{C}^n$, $\phi \in L^1(G)$ such that $U(\phi)\bar{x} = \bar{y}$. As above, this shows that M is minimal and is of dimension n . \square

Remark. Statement (1) means that G needs to be amenable if $F = LUC(G)$.

Theorem 2. *Let \mathcal{A} be $L^1(G)$, $M(G)$ or F^* , and let M be a left ideal of \mathcal{A} of dimension n . Then there exist m vectors $\bar{\mu}^i \in \mathcal{A}^{n_i}$ and m irreducible, unitary, bounded and continuous representations U^i ($i = 1, 2, \dots, m$) of G such that*

- (1) *each $\bar{\mu}^i$ is U^i -invariant,*
- (2) *for each $i = 1, 2, \dots, m$, the coordinates of $\bar{\mu}^i$ span a minimal left ideal M_i of \mathcal{A} of dimension n_i , and*
- (3) $M = M_1 \oplus M_2 \oplus \dots \oplus M_m$.

Proof. In the case of $\mathcal{A} = L^1(G)$, we regard M as a left ideal of $M(G)$. This is possible because M is closed and $L^1(G)$ is a closed ideal of $M(G)$. We start with a set of elements $\mu_1, \mu_2, \dots, \mu_n$ which generate M , and let $\bar{\mu} = (\mu_i)_{i=1}^n$. Then, for each $\mu \in \mathcal{A}$ and for each $i = 1, 2, \dots, n$, there exist $a_{i1}(\mu), a_{i2}(\mu), \dots, a_{in}(\mu) \in \mathbb{C}$ such that

$$\mu \mu_i = a_{i1}(\mu) \mu_1 + a_{i2}(\mu) \mu_2 + \dots + a_{in}(\mu) \mu_n.$$

Put $A(\mu) = (a_{ij}(\mu))_{i,j=1}^n$. Then, for μ and ν in \mathcal{A} ,

$$A(\mu\nu)\bar{\mu} = (\mu\nu)\bar{\mu} = \mu(A(\nu)\bar{\mu}) = A(\nu)(\mu\bar{\mu}) = A(\nu)A(\mu)\bar{\mu}.$$

Hence, $A(\mu\nu) = A(\nu)A(\mu)$, i.e., A is an antirepresentation of \mathcal{A} . Moreover, for each $1 \leq j \leq n$, let $f \in \mathcal{F}$ be such that $\mu_j(f) = 1$ and $\mu_k(f) = 0$ for $k \neq j$. Then, for each $1 \leq i \leq n$, $\mu \mu_i(f) = a_{ij}(\mu)$, so

$$|a_{ij}(\mu)| \leq \|\mu \mu_i\| \|f\| \leq \|\mu\| \|\mu_i\| \|f\|,$$

which implies that A is bounded. Since the product in \mathcal{A} is $\sigma(\mathcal{A}, \mathcal{F})$ -continuous on the left side, one can also see that the functions $\mu \mapsto a_{ij}(\mu)$ are $\sigma(\mathcal{A}, \mathcal{F})$ -continuous. When $\mathcal{A} = L^1(G)$, we let V be the antirepresentation of G associated to A ; and when $\mathcal{F} = F$ is an admissible subspace of $C(G)$, we let

$$V(s) = (a_{ij}(e(s)))_{i,j=1}^n.$$

Then \tilde{V} is a bounded and continuous representation of G in each case. Furthermore, for $f \in \mathcal{F}$, we have

$$\bar{\mu}(f_s) = e(s)\bar{\mu}(f) = A(e(s))\bar{\mu}(f) = V(s)\bar{\mu}(f),$$

i.e., $\bar{\mu}$ is \tilde{V} -invariant. Now it is easy to verify (or see [1, Lemma 3]) that \tilde{V} is in fact equivalent to a unitary representation U in the sense that $P\tilde{V}(s) = U(s)P$ for all $s \in G$, where P is an invertible operator on \mathbb{C}^n . (This result is also true for the infinite-dimensional representations when G is amenable; see [9] or [7].) Put

$\bar{\gamma} = P\bar{\mu}$. It is clear that the coordinates of $\bar{\gamma}$ also generate the ideal M . We have also

$$\bar{\gamma}(f_s) = P\bar{\mu}(f_s) = PV(s)\bar{\mu}(f) = \tilde{U}(s)P\bar{\mu}(f) = \tilde{U}(s)\bar{\gamma}(f),$$

and so $\bar{\gamma}$ is U -invariant. Since U is unitary, it follows by [8, 21.40(a)] that U is a direct sum of continuous, irreducible representations U^1, U^2, \dots , and U^m . This means that \mathbb{C}^n is the direct sum of some invariant subspaces H_i , and each U^i is the restriction of U to H_i ($i = 1, 2, \dots, m$). For each $f \in \mathcal{F}$, we write $\bar{\gamma}(f) = \sum_i \bar{\mu}^i(f)$. Then

$$\sum_{i=1}^m \bar{\mu}^i(f_s) = \bar{\gamma}(f_s) = \tilde{U}(s)\bar{\gamma}(f) = \sum_{i=1}^m \tilde{U}^i(s)\bar{\mu}^i(f).$$

It follows that, for each $i = 1, 2, \dots, m$, $\bar{\mu}^i(f_s) = \tilde{U}^i(s)\bar{\mu}^i(f)$. This yields statement (1). Statement (2) follows from Theorem 1. Statement (3) is clear. \square

Remark. In $L^1(G)^{**}$, the situation is slightly different. In fact, one can also produce finite-dimensional left ideals with the use of the right annihilators of $L^1(G)^{**}$. These are elements μ in $L^1(G)^{**}$ which satisfy $L^1(G)^{**}\mu = \{0\}$; see [6]. In such a situation, the representation \tilde{V} is trivial in the proof above.

Theorem 3. *Let M be a left ideal of $L^1(G)^{**}$ of dimension n , and suppose that M contains l linearly independent right annihilators $\gamma_1, \gamma_2, \dots, \gamma_l$ of $L^1(G)^{**}$. Then there exist m vectors $\bar{\mu}^i \in (L^1(G)^{**})^{n_i}$ and m irreducible, unitary, bounded and continuous representations U^i ($i = 1, 2, \dots, m$) of G such that*

- (1) each $\bar{\mu}^i$ is topologically U^i -invariant,
- (2) for each $i = 1, 2, \dots, m$, the coordinates of $\bar{\mu}^i$ span a minimal left ideal M_i of $L^1(G)^{**}$ of dimension n_i , and
- (3) $M = \mathbb{C}\gamma_1 \oplus \mathbb{C}\gamma_2 \oplus \dots \oplus \mathbb{C}\gamma_l \oplus M_1 \oplus M_2 \oplus \dots \oplus M_m$.

Proof. We take $n - l$ linearly independent elements $\mu_1, \mu_2, \dots, \mu_{n-l}$ in M which are not right annihilators of $L^1(G)^{**}$, let $\bar{\mu} = (\mu_i)_{i=1}^{n-l}$. Then, form the matrices $A(\mu)$ such that $\mu\bar{\mu} = A(\mu)\bar{\mu}$ for $\mu \in L^1(G)^{**}$, restrict A to $L^1(G)$, and let \tilde{V} be the corresponding representation of G . Then, for $\phi \in L^1(G)$ and $f \in L^\infty(G)$, we have

$$\bar{\mu}(\phi * f) = \bar{\mu}(\hat{\phi} * f) = \hat{\phi}\bar{\mu}(f) = A(\hat{\phi})\bar{\mu}(f) = V(\hat{\phi})\bar{\mu}(f) = \tilde{V}(\phi)\bar{\mu}(f),$$

and so $\bar{\mu}$ is topologically \tilde{V} -invariant. The proof is completed as that of Theorem 2. \square

Corollary. *Let G be a locally compact group. Then*

- (1) finite-dimensional (left) ideals exist in $M(G)$ and $L^1(G)$ if and only if G is compact,
- (2) finite-dimensional left ideals exist in $LUC(G)^*$ if and only if G is amenable,
- (3) finite-dimensional left ideals which are not generated by right annihilators of $L^1(G)^{**}$ exist in $L^1(G)^{**}$ if and only if G is amenable.

Proof. This follows from Lemma 2 and Theorems 2 and 3. \square

Remark. The finite-dimensional right ideals in $WAP(G)^*$ are determined in the same way because the two Arens product coincide in this case; see [2, Section 4.2]. When G is compact, one proceeds also in the same way to find these ideals in $L^1(G)$ and $M(G)$. These facts were already observed for the minimal right ideals in [1,

Section 4]. However, in [4, Remark 2.7(b)], we have proved that the non-trivial right ideals are all of infinite dimension in $LUC(\mathbb{Z})^* = \ell^\infty(\mathbb{Z})^*$, where \mathbb{Z} is the additive group of the integers. In [1, Section 4], we have given a class of locally compact abelian groups, which includes \mathbb{Z} , for which the non-trivial right ideals are all of infinite dimension in $LUC(G)^*$. Now we can prove that, for a locally compact abelian group G , the finite-dimensional right ideals exist in $LUC(G)^*$ if and only if G is compact. We hope to publish this result in another paper.

The author wishes to thank the referee for the very careful reading the paper was given.

REFERENCES

- [1] J. W. Baker and M. Filali, *Minimal ideals in group algebras and their biduals*, Math. Proc. Cambridge Philos. Soc. **120** (1996), 475–488. CMP 96:12
- [2] J. F. Berglund, H. D. Jungheun and P. Milnes, *Analysis on Semigroups: Function Spaces, Compactifications, Representations*, Wiley, New York, 1989. MR **91b**:43001
- [3] M. Filali, *The ideal structure of some Banach algebras*, Math. Proc. Cambridge Philos. Soc. **111** (1992), 567–576. MR **93e**:22005
- [4] M. Filali, *Linear equations in $B(\mathbb{Z})^*$* , Proc. Roy. Soc. Edinburgh Sect. A **123A** (1993), 1001–1009. MR **95j**:39035
- [5] M. Filali, *Minimal left ideals in the group algebra and its second conjugate*, Mathematica Japonica (to appear).
- [6] E. E. Granirer, *The radical of $L^\infty(G)^*$* , Proc. Amer. Math. Soc. **41** (1973), 321–324. MR **48**:4646
- [7] F. P. Greenleaf, *Invariant means on topological groups and their applications*, Van Nostrand, New York, 1969. MR **40**:4776
- [8] E. Hewitt and K. A. Ross, *Abstract harmonic analysis I*, Springer-Verlag, Berlin,, 1963, 1970. MR **28**:158
- [9] B. Sz.-Nagy, *Uniformly bounded linear transformations in Hilbert space*, Acta Math. (Szeged) **11** (1947), 152–157.

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF OULU, SF 90570 FINLAND
E-mail address: mfilali@cc.oulu.fi