

## OSSA'S THEOREM AND ADAMS COVERS

ROBERT R. BRUNER

(Communicated by Ralph Cohen)

ABSTRACT. We show that Ossa's theorem splitting  $ku \wedge BV$  for elementary abelian groups  $V$  follows from general facts about  $ku \wedge BZ/2$  and Adams covers. For completeness, we also provide the analogous results for  $ko \wedge BV$ .

### 1. INTRODUCTION

The connective complex  $K$ -homology of  $BV$ ,  $V$  elementary abelian, has been computed by Ossa [4], and, by different techniques, by Johnson and Wilson [1]. A key step in both proofs is the equivalence of  $ku$ -module spectra

$$ku \wedge BZ/2 \wedge BZ/2 \simeq ku \wedge \Sigma^2 BZ/2 \vee \Sigma^2 H(\mathbb{F}_2[u, v])$$

where  $H(\mathbb{F}_2[u, v])$  is the generalized Eilenberg-MacLane spectrum on the graded vector space  $\mathbb{F}_2[u, v]$  with  $|u| = |v| = 2$ .

The purpose of this note is to show that this equivalence follows from two more general facts of independent interest about such smash products. We also provide analogous results in the real case, and the splitting

$$ko \wedge BZ/2 \wedge BZ/2 \simeq ko\langle 1 \rangle \wedge BZ/2 \vee \Sigma^2 H(\mathbb{F}_2[u, v^2]),$$

where  $ko\langle 1 \rangle$  is the connected cover of  $ko$ , that is, the fiber of the Thom map  $ko \rightarrow H\mathbb{Z}$ .

The calculation of  $ku \wedge BV$  and  $ko \wedge BV$  from these splittings follows from formal properties of products, and the reader is referred to [4] or [1] for details. Results analogous to the complex case hold for both the real and complex cases at odd primes.

I would like to thank Stephan Stolz for introducing me to the essential cofiber sequence (1) as the simplest way to calculate  $ku_* BZ/2$  and  $ko_* BZ/2$ . Its use for this purpose is probably due to Mark Mahowald or Don Davis.

### 2. GENERALITIES

As  $BZ/2$  is a 2-local spectrum, we shall localize at 2 throughout.

*Notation 2.1.* Let  $H = H\mathbb{F}_2$  and let 'cohomology' mean mod 2 cohomology. Let  $\mathcal{A}$  be the mod 2 Steenrod algebra,  $\mathcal{A}(i)$  the subalgebra generated by  $Sq^1, \dots, Sq^{2^i}$ , and  $E(1) = E[Q_0, Q_1]$  the exterior subalgebra of  $\mathcal{A}(1)$  generated by  $Q_0 = Sq^1$  and  $Q_1 = [Sq^2, Q_0]$ .

---

Received by the editors October 22, 1997.

1991 *Mathematics Subject Classification.* Primary 55P10, 55N20; Secondary 55N15, 55S10.

Let  $P = BZ/2$  and let  $R$  be the suspension of the subcomplex of the stunted projective space  $P_{-1}^\infty$  omitting the 0-cell. (Thus,  $H^*R = \Sigma F_{-1,1}$  in the notation of [2].)

The bottom cell of  $R$  is in dimension 0, and the fiber of the inclusion of the bottom cell is  $P$ . This gives the key cofiber sequence

$$(1) \quad P \longrightarrow S \longrightarrow R.$$

**Lemma 2.2.**

$$ko \wedge R \simeq \bigvee_{i \geq 0} \Sigma^{4i} H\mathbb{Z}$$

and

$$ku \wedge R \simeq \bigvee_{i \geq 0} \Sigma^{2i} H\mathbb{Z}.$$

*Proof.* It suffices to prove the first equivalence, since  $ku \simeq ko \wedge C\eta$ . In [2] it is shown that

$$(2) \quad \mathcal{A} \otimes_{\mathcal{A}(1)} H^*R \cong \bigoplus_{i \geq 0} \mathcal{A} \otimes_{\mathcal{A}(0)} \Sigma^{4i} \mathbb{F}_2.$$

Since  $H^*(ko \wedge R) \cong (\mathcal{A} \otimes_{\mathcal{A}(1)} \mathbb{F}_2) \otimes H^*R \cong \mathcal{A} \otimes_{\mathcal{A}(1)} H^*R$ , the isomorphism (2) implies that the Adams spectral sequence

$$\text{Ext}_{\mathcal{A}}(H^*(ko \wedge R), \mathbb{F}_2) \implies \pi_*(ko \wedge R)$$

is isomorphic, by a standard change of rings argument, to

$$\bigoplus_{i \geq 0} \text{Ext}_{\mathcal{A}(0)}(\Sigma^{4i} \mathbb{F}_2, \mathbb{F}_2) \implies \pi_*(ko \wedge R).$$

This must collapse at  $E_2$  because it is zero except in total degrees congruent to 0 modulo 4. If we let  $x_i : ko \wedge R \rightarrow \Sigma^{4i} H\mathbb{Z}$  be a cohomology class dual to the generator of  $\text{Ext}^{0,4i}(H^*(ko \wedge R), \mathbb{F}_2)$ , then the collapse of the Adams spectral sequence implies that

$$\bigvee_{i \geq 0} x_i : ko \wedge R \longrightarrow \bigvee_{i \geq 0} \Sigma^{4i} H\mathbb{Z}$$

induces an isomorphism in homotopy, and is thus a homotopy equivalence. □

**Theorem 2.3.** *For any spectrum  $X$  there are cofiber sequences*

$$ko \wedge P \wedge X \xrightarrow{i} ko \wedge X \xrightarrow{p} \bigvee_{i \geq 0} \Sigma^{4i} H\mathbb{Z} \wedge X$$

and

$$ku \wedge P \wedge X \xrightarrow{i} ku \wedge X \xrightarrow{p} \bigvee_{i \geq 0} \Sigma^{2i} H\mathbb{Z} \wedge X$$

in which the homomorphisms  $p^*$  are epimorphisms in cohomology.

*Proof.* Smash the cofiber sequence (1) with  $ko \wedge X$  or  $ku \wedge X$  and apply Lemma 2.2. Since tensor products preserve epimorphisms, by the Kunn eth theorem it is sufficient to note that  $H^*R \rightarrow H^*S$  is an epimorphism. □

Thus, smashing  $ko \wedge X$  or  $ku \wedge X$  with  $P$  has the effect of taking a ‘generalized Adams cover’. This is not in general an actual  $H\mathbb{Z}$  Adams cover because  $H\mathbb{Z} \wedge X$  is not in general a coproduct of  $H\mathbb{Z}$ ’s. Further, this cover contains a large generalized Eilenberg-Mac Lane summand, so is far from minimal.

### 3. THE COMPLEX CASE

In the complex case we have a much smaller generalized Adams cover.

**Proposition 3.1.** *For any spectrum  $X$  there is a cofiber sequence*

$$ku \wedge \Sigma^2 X \xrightarrow{u} ku \wedge X \xrightarrow{p} H\mathbb{Z} \wedge X,$$

where  $u \in ku_2$  is the multiplicative generator of  $ku_*$  and  $p^*$  is an epimorphism in cohomology.

*Proof.* Multiplication by the Bott class  $u \in ku_2$  gives a cofiber sequence

$$\Sigma^2 ku \xrightarrow{u} ku \longrightarrow H\mathbb{Z}$$

in which the map  $ku \longrightarrow H\mathbb{Z}$  induces on  $H^*$  the quotient

$$\mathcal{A} \otimes_{E(1)} \mathbb{F}_2 \longleftarrow \mathcal{A} \otimes_{A(0)} \mathbb{F}_2.$$

This remains an epimorphism upon tensoring with  $H^*X$ . □

We would like to claim that this smaller generalized Adams cover of  $ku \wedge X$  splits off the one in Theorem 2.3. However, there is no comparison theorem for  $H\mathbb{Z}$  Adams resolutions. For example, the map  $ku \longrightarrow ku \wedge R$  of Theorem 2.3 does not factor through the map  $ku \longrightarrow H\mathbb{Z}$  of Proposition 3.1. The standard theorems which assert the existence of a comparison theorem would require that  $H\mathbb{Z}^*(H\mathbb{Z})$  be projective over  $\pi_*H\mathbb{Z}$ , which it is not. However, with coefficients in a field, such a comparison theorem does exist and allows us to compare the cofibrations above.

**Theorem 3.2.** *If  $H^*X$  is free over  $E[\beta]$  and  $ku_*X \longrightarrow H\mathbb{Z}_*X$  is onto, then*

$$ku \wedge P \wedge X \simeq ku \wedge \Sigma^2 X \vee \bigvee_{i>0} \Sigma^{2i-1} H\mathbb{Z} \wedge X.$$

*Proof.* If  $H^*X$  is  $E[\beta]$ -free, then  $H\mathbb{Z}^*X$  is a mod 2 vector space, so  $H\mathbb{Z} \wedge X$  is a wedge of  $H = H\mathbb{F}_2$ ’s. Hence, the sequences in Theorems 2.3 and 3.1 are both beginnings of  $H\mathbb{F}_2$ -Adams resolutions. As such, the comparison theorem gives maps between them which cover the identity of  $ku \wedge X$ .

$$\begin{array}{ccccc} ku \wedge \Sigma^2 X & \longrightarrow & ku \wedge P \wedge X & \longrightarrow & ku \wedge \Sigma^2 X \\ \downarrow & & \downarrow & & \downarrow \\ ku \wedge X & = & ku \wedge X & = & ku \wedge X \\ \downarrow & & \downarrow & & \downarrow \\ H\mathbb{Z} \wedge X & \longrightarrow & \bigvee_{i \geq 0} \Sigma^{2i} H\mathbb{Z} \wedge X & \longrightarrow & H\mathbb{Z} \wedge X. \end{array}$$

Since  $ku_*X \longrightarrow H\mathbb{Z}_*X$  is onto, the bottom row must be the identity in homotopy, hence an equivalence. It follows that we have a splitting

$$ku \wedge P \wedge X \simeq ku \wedge \Sigma^2 X \vee \bigvee_{i>0} \Sigma^{2i-1} H\mathbb{Z} \wedge X$$

since the left map in the bottom row may be taken to be the inclusion of the  $i = 0$  summand. □

**Corollary 3.3.**

$$ku \wedge P \wedge P \simeq ku \wedge \Sigma^2 P \vee \bigvee_{i,j>0} \Sigma^{2i+2j-2} H\mathbb{F}_2.$$

*Proof.* This follows from Theorem 3.2 and the equivalence

$$HZ \wedge P \simeq \bigvee_{j>0} \Sigma^{2j-1} H\mathbb{F}_2$$

since it is easy to see that  $ku_*P \rightarrow HZ_*P$  is onto. □

*Remark 3.4.* We get the dimensions right if we write the generalized Eilenberg-Mac Lane summand as the double suspension of  $H(\mathbb{F}_2[u, v])$ . However, the geometric origin of the splitting is clearer with the following parameterization. We have  $H^*(P \wedge P) = \langle x_1 x_2 \rangle$ , the ideal generated by  $x_1 x_2$  in  $H^*(P \times P)_+ = \mathbb{F}_2[x_1, x_2]$ . As an  $E(1)$ -module, each  $x_1^{2i-1} x_2^{2j-1}$  generates a copy of  $E(1)$  and these form an  $E(1)$ -free submodule

$$\bigoplus_{i,j>0} \langle x_1^{2i-1} x_2^{2j-1} \rangle \subset H^*(P \wedge P).$$

Since  $E(1)$  is a Frobenius algebra, this submodule is a direct summand and it is easy to check that the quotient is isomorphic to  $H^*(\Sigma^2 P)$ . For example, we can write

$$H^*(P \wedge P) \cong x_1^2(x_2) \oplus \bigoplus_{i,j>0} \langle x_1^{2i-1} x_2^{2j-1} \rangle.$$

Since  $Q_0$  and  $Q_1$  act trivially on  $x_1^2$ , the first summand is  $H^*(\Sigma^2 P)$  as an  $E(1)$ -module. If all one wants is to understand  $ku \wedge P \wedge P$ , this splitting of cohomology and Margolis’s theorem, that  $\mathcal{A}$ -free submodules in cohomology correspond to  $H\mathbb{F}_2$  wedge summands ([3]), are sufficient.

*Remark 3.5.* In general, an Adams cover of  $ku \wedge X$  is not just  $ku \wedge \Sigma^2 X$ . For example,  $X = BZ/4$  is composed of mod 4 Moore spaces, and  $ku_*BZ/4 \rightarrow HZ_*BZ/4$  is onto, presumably the mildest weakening of the hypotheses of Theorem 3.2. We have the following homotopy groups for  $ku \wedge BZ/4$  and its first three Adams covers, where we write  $a + b + c$  for the group  $\mathbb{Z}/a \oplus \mathbb{Z}/b \oplus \mathbb{Z}/c$  and 0 for the trivial group.

Adams cover	$\pi_1$	$\pi_3$	$\pi_5$	$\pi_7$	$\pi_9$	$\pi_{11}$
0	4	2+8	2+2+16	4+2+32	4+4+64	8+4+128
1	2	2+4	2+2+8	4+2+16	4+4+32	8+4+64
2	0	2	2+4	2+2+8	4+2+16	4+4+32
3	0	0	2	2+4	2+2+8	4+2+16

From this it is clear that the first Adams cover is not merely a suspension, but it is plausible that the  $i^{th}$  Adams cover is the double suspension of the  $(i - 1)^{st}$  for  $i > 1$ .

4. THE REAL CASE

We record the following trivial consequence of the definitions for reference.

**Proposition 4.1.** *For any spectrum  $X$  there is a cofiber sequence*

$$ko\langle 1 \rangle \wedge X \rightarrow ko \wedge X \xrightarrow{p^*} HZ \wedge X,$$

where  $p^*$  is an epimorphism in cohomology.

As in the complex case, with coefficients in a field the comparison theorem allows us to compare the preceding cofibration with that in Theorem 2.3.

**Theorem 4.2.** *If  $H^*X$  is free over  $E[\beta]$  and  $ko_*X \rightarrow H\mathbb{Z}_*X$  is onto, then*

$$ko \wedge P \wedge X \simeq ko\langle 1 \rangle \wedge X \vee \bigvee_{i>0} \Sigma^{4i-1} H\mathbb{Z} \wedge X.$$

*Proof.* The proof is entirely analogous to the complex case. □

*Remark 4.3.* It might be better to replace  $ko\langle 1 \rangle \wedge X$  by  $(ko \wedge X)\langle 1 \rangle$ , a potentially smaller Adams cover of  $ko \wedge X$ . For example,

$$ko\langle 1 \rangle \wedge P \simeq (ko \wedge P)\langle 1 \rangle \vee \bigvee_{i>0} \Sigma^{4i} H\mathbb{F}_2.$$

**Theorem 4.4.**

$$\begin{aligned} ko \wedge P \wedge P &\simeq ko\langle 1 \rangle \wedge P \vee \bigvee_{i,j>0} \Sigma^{4i+2j-2} H\mathbb{F}_2 \\ &\simeq (ko \wedge P)\langle 1 \rangle \vee \bigvee_{i>0,j\geq 0} \Sigma^{4i+2j-2} H\mathbb{F}_2 \end{aligned}$$

where  $(ko \wedge P)\langle 1 \rangle$  is the minimal Adams cover of  $ko \wedge P$ . The cohomology of  $(ko \wedge P)\langle 1 \rangle$  is tensored up from an  $\mathcal{A}(1)$ -module with Poincare series  $t^3/(1-t) + t^2(1+t)(1+t^3)$ .

*Proof.* Theorem 4.2 does not apply, so we must resort to an ad hoc argument in this case. We decompose  $H^*(P \wedge P)$  as a module over  $\mathcal{A}(1)$  into an  $\mathcal{A}(1)$ -free summand, producing the generalized Eilenberg-MacLane spectrum upon smashing with  $ko$ , and the  $\mathcal{A}(1)$ -module which is the kernel of the minimal projective cover of  $H^*P$  in  $\mathcal{A}(1)\text{-Mod}$ . □

*Remark 4.5.* The homotopy type of  $ko \wedge P^{\wedge n}$  modulo generalized Eilenberg-MacLane summands depends upon the congruence class of  $n$  modulo 4. This is an easy consequence of the homotopy of the Adams covers of  $ko \wedge P$ , and is the maximal number of different homotopy types since the 4<sup>th</sup> Adams cover of  $ko$  is  $\Sigma^8 ko$ .

REFERENCES

[1] David Copeland Johnson and W. Stephen Wilson, "On a Theorem of Ossa" *Proc. Amer. Math. Soc.* **125** (1997), 3753–3755. MR **98b**:55003  
 [2] W. H. Lin, D. Davis, M. E. Mahowald and J. F. Adams, "Calculation of Lin's Ext Groups", *Math. Proc. Camb. Phil. Soc.* **87** (1980), 459–469. MR **81e**:55025  
 [3] H. R. Margolis, "Eilenberg-MacLane Spectra", *Proc. Amer. Math. Soc.* **43** (1974), 409–415. 1973. MR **49**:6239  
 [4] Eric Ossa, "Connective K Theory of Elementary Abelian Groups", *Proceedings of the 1987 Osaka Conference on Transformation groups*, Springer Lecture Notes in Mathematics, V. **1375**, 269–275. MR **90h**:55009

DEPARTMENT OF MATHEMATICS, WAYNE STATE UNIVERSITY, DETROIT, MICHIGAN 48202  
*E-mail address:* rrb@math.wayne.edu