

CONVOLUTION WITH AFFINE ARCLENGTH MEASURES IN THE PLANE

DANIEL M. OBERLIN

(Communicated by Christopher D. Sogge)

ABSTRACT. We obtain an estimate for the $L^{3/2,1}(\mathbb{R}^2) - L^3(\mathbb{R}^2)$ norm of a certain convolution operator.

Let ϕ be real-valued and smooth on an interval $(a, b) \subseteq \mathbb{R}$. Define the measure λ on \mathbb{R}^2 by

$$\int_{\mathbb{R}^2} f \, d\lambda = \int_a^b f(t, \phi(t)) |\phi''(t)|^{1/3} dt.$$

We are interested in the $L^p(\mathbb{R}^2) - L^q(\mathbb{R}^2)$ mapping properties of the operator given by convolution with λ . The study of this operator was initiated by Drury ([D]), who used complex interpolation and certain integral estimates to obtain the optimal result

$$\|\lambda * f\|_3 \leq C \|f\|_{3/2}$$

when ϕ satisfies certain regularity conditions. Drury's approach was refined and extended by Choi in [C]. The purpose of this note is to prove an almost-sharp estimate for a large class of functions ϕ :

Theorem. *Suppose $\phi'' > 0$ and $\phi^{(3)} \geq 0$ on (a, b) . Then*

$$(1) \quad \|\lambda * \chi_E\|_3 \leq 12^{1/3} \|\chi_E\|_{3/2}$$

for any measurable $E \subseteq \mathbb{R}^2$.

Proof. Define $\gamma(s) = (s, \phi(s))$. Inequality (1) is equivalent to

$$\int_{\mathbb{R}^2} \int_a^b \int_a^b \int_a^b \prod_1^3 \left[\chi_E(x - \gamma(s_i)) \phi''(s_i)^{1/3} \right] ds_1 ds_2 ds_3 dx \leq 12 |E|^2$$

and so will follow from

$$\int_{\mathbb{R}^2} \int_a^b \int_{s_3}^b \int_{s_3}^b \prod_1^3 \left[\chi_E(x - \gamma(s_i)) \phi''(s_i)^{1/3} \right] ds_1 ds_2 ds_3 dx \leq 4 |E|^2.$$

Received by the editors February 16, 1998.

1991 *Mathematics Subject Classification.* Primary 42B15.

The author was partially supported by a grant from the National Science Foundation.

By Fubini's theorem this is equivalent to

$$\int_{\mathbb{R}^2} \chi_E(x) \int_a^b \left(\int_t^b \chi_E(x + \gamma(t) - \gamma(s)) \phi''(s)^{1/3} ds \right)^2 \phi''(t)^{1/3} dt dx \leq 4 |E|^2.$$

Thus, it is enough to establish the inequality

$$(2) \quad \int_a^b \left(\int_t^b \chi_E(\gamma(t) - \gamma(s)) \phi''(s)^{1/3} ds \right)^2 \phi''(t)^{1/3} dt \leq 4 |E|.$$

The convexity of the graph of ϕ shows that the change of variables

$$(s, t) \rightarrow \gamma(s) - \gamma(t) = (s - t, \phi(s) - \phi(t))$$

is one-to-one. Thus

$$\int_a^b \int_a^b \chi_E(\gamma(t) - \gamma(s)) |\phi'(s) - \phi'(t)| ds dt \leq |E|$$

and (2) will follow from the inequality

$$(3) \quad \phi''(t)^{1/3} \left(\int_t^b \chi_A(s) \phi''(s)^{1/3} ds \right)^2 \leq 4 \int_t^b \chi_A(s) (\phi'(s) - \phi'(t)) ds$$

if $A \subseteq (t, b)$. To prove (3) we let $|A_u|$ stand for the (one-dimensional) Lebesgue measure of $A \cap (u, b)$ whenever $t \leq u \leq b$. Then

$$(4) \quad \int_t^b \chi_A(s) (\phi'(s) - \phi'(t)) ds = \int_t^b \chi_A(s) \int_t^s \phi''(u) du ds = \int_t^b \phi''(u) |A_u| du.$$

Also

$$\begin{aligned} \int_t^b \chi_A(s) \phi''(s)^{1/3} ds &= \int_t^b \chi_A(s) \phi''(s)^{1/3} |A_s|^{1/3} |A_s|^{-1/3} ds \\ &\leq \left(\int_t^b \chi_A(s) \phi''(s) |A_s| ds \right)^{1/3} \left(\int_t^b \chi_A(s) |A_s|^{-1/2} ds \right)^{2/3}. \end{aligned}$$

Thus, it follows from (4) that

$$(5) \quad \left(\int_t^b \chi_A(s) \phi''(s)^{1/3} ds \right)^3 \leq \left(\int_t^b \chi_A(s) (\phi'(s) - \phi'(t)) ds \right) \left(\int_t^b \chi_A(s) |A_s|^{-1/2} ds \right)^2.$$

If $0 \leq \rho \leq |A|$, then $|\{s \in A : |A_s| \leq \rho\}| = \rho$, and so

$$\int_t^b \chi_A(s) |A_s|^{-1/2} ds = \int_0^{|A|} y^{-1/2} dy = 2 |A|^{1/2}.$$

With this and the fact that ϕ'' is nondecreasing, (5) yields (3) to complete the proof.

REFERENCES

- [C] Y. Choi, Convolution operators with affine arclength measures on plane curves, *J. Korean Math. Soc.* 36 (1999), 193–207. CMP 99:09
 [D] S. W. Drury, Degenerate curves and harmonic analysis, *Math. Proc. Camb. Phil. Soc.* 108 (1990), 89–96. MR 91h:42021

DEPARTMENT OF MATHEMATICS, FLORIDA STATE UNIVERSITY, TALLAHASSEE, FLORIDA 32306-4510

E-mail address: oberlin@math.fsu.edu