

WEAKLY COMPACT COMPOSITION OPERATORS BETWEEN ALGEBRAS OF BOUNDED ANALYTIC FUNCTIONS

PABLO GALINDO, MIKAEL LINDSTRÖM, AND RAY RYAN

(Communicated by Steven R. Bell)

ABSTRACT. We prove a characterization (up to the approximation property) of weakly compact composition operators $C_\phi : H^\infty(B_F) \rightarrow H^\infty(B_E)$ in terms of their inducing analytic maps $\phi : B_E \rightarrow B_F$.

Let E denote a complex Banach space with open unit ball B_E and let $\phi : B_E \rightarrow B_F$ be an analytic map, where F is also a complex Banach space. We will consider the composition operator C_ϕ defined by $C_\phi(f) = f \circ \phi$, acting from the uniform algebra $H^\infty(B_F)$ of all bounded analytic functions on B_F into $H^\infty(B_E)$.

In [U] A. Ülger proved that every weakly compact homomorphism from a log-modular uniform algebra into any uniform algebra is compact. This was also later proved in [GL] by a slightly different proof. In his paper A. Ülger also asks whether every weakly compact homomorphism between uniform algebras is compact without the logmodularity condition. An example showing that this is not generally true was given in [AGL]. In fact it was shown that the uniform algebra $H^\infty(B_E)$ when E is the Tsirelson space and $\phi(x) = \frac{x}{2}$ give rise to a weakly compact homomorphism which is not compact. In this note we characterize (modulo the approximation property) the weakly compact composition operators C_ϕ which makes it possible, in a general way, to produce noncompact weakly compact composition operators C_ϕ . As a byproduct of our technique we characterize the completely continuous composition operators. In [GG] M. González and J. Gutiérrez have studied weakly compact composition operators between the Fréchet algebras $H_b(B_E)$ of analytic functions of bounded type endowed with the topology of uniform convergence on B_E -bounded sets.

Preliminaries. The reader is referred to [D] and [M2] for background information on analytic functions on an infinite dimensional Banach space. The algebra $H^\infty(B_E)$ is a Banach algebra with the natural norm $\|f\| = \sup_{x \in B_E} |f(x)|$. This algebra, which is a natural generalization of the classical algebra $H^\infty(\Delta)$ of analytic functions on the complex open disk Δ , has been studied in [ACG]. A homomorphism between Banach algebras is a continuous linear multiplicative map. By an operator we mean a continuous linear map from a Banach space into another Banach space. The space of all operators from E into F is denoted by $L(E, F)$. We denote

Received by the editors March 10, 1998.

1991 *Mathematics Subject Classification.* Primary 46J15; Secondary 46E15, 46G20.

Key words and phrases. Weakly compact operator, composition operator, bounded analytic functions on the open unit ball.

Research of the first author was partially supported by DGICYT(Spain) pr. 91-0326.

the adjoint operator of $T \in L(E, F)$ by $T^t : F^* \rightarrow E^*$. We say that $T \in L(E, F)$ is (weakly) compact if T maps bounded sets in E into relatively (weakly) compact sets in F . If T maps the closed unit ball of E onto a conditionally weakly compact set, T is called a *Rosenthal operator*. Clearly every weakly compact operator is Rosenthal. The operator T is called *completely continuous*, if it maps every weakly convergent sequence of E into a norm convergent one of F . Every compact operator is completely continuous. A subset $A \subset E$ is called a *Dunford-Pettis set*, if for every weakly null sequence $(l_n) \subset E^*$ and every sequence $(x_n) \subset A$ we have that $l_n(x_n) \rightarrow 0$, when $n \rightarrow \infty$. If E^* has the Schur property, then every bounded set in E is a Dunford-Pettis set.

Let $P(E)$ denote the algebra of all continuous polynomials on E . We denote by $(E, \sigma(E, P(E)))$ (respectively $(B_E, \sigma(E, H^\infty(B_E)))$) the set E (respectively B_E) endowed with the weakest topology making all $p \in P(E)$ (respectively $f \in H^\infty(B_E)$) continuous. The topology $\sigma(E, P(E))$ is a regular Hausdorff topology such that $(E, \|\cdot\|) \geq (E, \sigma(E, P(E))) \geq (E, \sigma(E, E^*))$. Thus it follows that $\sigma(E, P(E))$ is angelic, and consequently the concepts (relatively) countably compact, (relatively) sequentially compact and (relatively) compact all agree with respect to this topology. A Banach space E is called a Λ -space, if all null sequences in $(E, \sigma(E, P(E)))$ are norm convergent, and hence convergent sequences in $(E, \sigma(E, P(E)))$ are also norm convergent. All superreflexive spaces and ℓ_1 are Λ -spaces [JaP].

The space $H^\infty(B_E)$ is a dual space, i.e., there is a Banach space $G^\infty(B_E)$ such that $H^\infty(B_E) = G^\infty(B_E)^*$. This fact follows from a theorem of K. Ng [N] and has been pointed out by S. Dineen in his book [D] and developed by J. Mujica in [M1]. By τ_0 we denote the compact-open topology on $H^\infty(B_E)$. $G^\infty(B_E)$ is defined as the subspace of $H^\infty(B_E)^*$ of those functionals which are τ_0 continuous when restricted to the unit ball of $H^\infty(B_E)$ or equivalently to the bounded subsets. The correspondence $f \in H^\infty(B_E) \rightarrow T_f \in G^\infty(B_E)^*$ given by $T_f(u) = u(f)$, $u \in G^\infty(B_E)$, is an isometric isomorphism. The map $\delta : B_E \rightarrow G^\infty(B_E)$ is defined by $\delta(x) = \delta_x$, where $\delta_x(f) = f(x)$ for all $f \in H^\infty(B_E)$. The space $G^\infty(B_E)$ is a closed subspace of $H^\infty(B_E)^*$ and the proof of Theorem 2.1 in [M1] shows that the closed unit ball of $G^\infty(B_E)$ coincides with the closed, convex, balanced hull $\overline{\Gamma}\{\delta_x : x \in B_E\}$ of $\{\delta_x : x \in B_E\}$. In particular, we have that $G^\infty(B_E)$ is the closed span of $\{\delta_x : x \in B_E\}$ in $H^\infty(B_E)^*$.

In [AGL] the following characterization of compactness of the composition operator C_ϕ was obtained in the case $E = F$. The same proof works also in this more general situation.

Theorem. *Consider the composition operator $C_\phi : H^\infty(B_F) \rightarrow H^\infty(B_E)$. The following statements are equivalent:*

- (i) C_ϕ is compact;
- (ii) C_ϕ is weakly compact and $\phi(B_E)$ is relatively compact in F ;
- (iii) $\phi(B_E)$ lies strictly inside B_F and $\phi(B_E)$ is relatively compact in F .

Weak compactness of composition operators. We start with an elementary observation.

Lemma 1. *Let (x_α) be a net in E such that $\|x_\alpha\| \leq r$ for some $0 < r < 1$ and let $x \in B_E$. If $p(x_\alpha) \rightarrow p(x)$ for all $p \in P(E)$, then $f(x_\alpha) \rightarrow f(x)$ for all $f \in H^\infty(B_E)$. Hence for given $0 < r < 1$, we have that $\sigma(E, H^\infty(B_E))$ and $\sigma(E, P(E))$ coincide on $r\overline{B}_E$.*

Proof. Take $f \in H^\infty(B_E)$. It is a uniform limit on every sB_E , $0 < s < 1$, of the partial sums of its Taylor series. Thus there is a sequence $(p_n) \subset P(E)$ such that $p_n \rightarrow f$ uniformly on sB_E for every $0 < s < 1$. Let $\varepsilon > 0$. Hence there is n_0 such that $\sup_{\|x_\alpha\| \leq r} |p_{n_0}(x_\alpha) - f(x_\alpha)| < \frac{\varepsilon}{2}$ and $|p_{n_0}(x) - f(x)| < \frac{\varepsilon}{2}$, so for all α we have

$$|f(x_\alpha) - f(x)| \leq \varepsilon + |p_{n_0}(x_\alpha) - p_{n_0}(x)|.$$

Thus $f(x_\alpha) \rightarrow f(x)$, when $\alpha \rightarrow \infty$. \square

Proposition 2. *If $C_\phi : H^\infty(B_F) \rightarrow H^\infty(B_E)$ is Rosenthal or completely continuous, then $\phi(B_E)$ lies strictly inside B_F .*

Proof. If not, then there is a sequence $(x_j) \subset B_E$ such that $\|\phi(x_j)\| > 1 - \frac{1}{j}$ for all j . Hence $\lim \|\phi(x_j)\| = 1$. By the proof of Theorem 10.5 in [ACG] there is a $g \in H^\infty(B_F^{**})$ which satisfies $|g| < 1$ and such that $(g(\phi(x_k)))_k$ is interpolating for $H^\infty(\Delta)$, where Δ is the open unit disk in \mathbb{C} and (x_k) is a subsequence of (x_j) . Hence, by Theorem 2.1 in [G, p. 294], there is a sequence $(f_n) \subset H^\infty(\Delta)$ and a constant $M > 0$ such that

$$f_n(g(\phi(x_k))) = 0, \text{ if } n \neq k, \quad f_n(g(\phi(x_n))) = 1,$$

and

$$\sum_{n=1}^{\infty} |f_n(z)| \leq M \text{ for all } z \in \Delta.$$

Now we define a map $T : H^\infty(B_E) \rightarrow l^\infty$ by $T(f) = (f(x_k))_k$ and another map $S : l^\infty \rightarrow H^\infty(B_F)$ by $S((\xi_n)) = \sum_{n=1}^{\infty} \xi_n f_n \circ g$. These two maps are both well-defined, continuous and linear. Further it can be seen that $T \circ C_\phi \circ S = id_{l^\infty}$. Since id_{l^∞} is neither Rosenthal nor completely continuous, we get a contradiction. \square

A portion of the next proposition relies on the proof of Proposition 3 in [GGM].

Proposition 3. *Assume F has the approximation property. If $C_\phi : H^\infty(B_F) \rightarrow H^\infty(B_E)$ is a weakly compact composition operator, then $\phi(B_E)$ is relatively compact in $(B_F, \sigma(F, H^\infty(B_F)))$.*

Proof. Let us denote $\sigma_F = \sigma(F, H^\infty(B_F))$ and $\sigma_{G^\infty} = \sigma(G^\infty(B_F), H^\infty(B_F))$. By weak compactness of C_ϕ^t and the fact that $C_\phi^t(\delta_{B_E}) = \delta_{\phi(B_E)}$, we get that its weak closure in $H^\infty(B_F)^*$, $\overline{\delta_{\phi(B_E)}}$ $\subset G^\infty(B_F)$, endowed with the weak topology is compact. Hence $\overline{\delta_{\phi(B_E)}}$ is compact in $G^\infty(B_F)$ with the induced topology, i.e., in σ_{G^∞} . Let $(\phi(x_\alpha)) \subset \phi(B_E)$ be an arbitrary net. We will show that it has a σ_F converging subnet to a point in B_F and that will prove that $\phi(B_E)$ is a σ_F relatively compact set in B_F . Since $\overline{\delta_{\phi(B_E)}}^{\sigma_{G^\infty}}$ is a compact set, there is a subnet (which we denote the same) σ_{G^∞} converging to some $u \in \overline{\delta_{\phi(B_E)}}^{\sigma_{G^\infty}}$. Let $y = u|_{F^*} \in F^{**}$. Observe that by Proposition 2, $\phi(B_E) \subset rB_F$ for some $0 < r < 1$, hence $\|y\| \leq r$. We claim that $y \in F$ and $u = \delta_y$. Recall that u is $(H^\infty(B_F), \tau_0)$ continuous on bounded subsets of $H^\infty(B_F)$. An application of the Banach-Dieudonné theorem [M3, Theorem 2.1], for $P^{(m)}F$, $m \in \mathbb{N}$, leads to the $(P^{(m)}F, \tau_0)$ continuity of u . In particular $y \in F$ since the compact open topology on F^* belongs to the dual pair (F, F^*) . Then $u(P) = P(y)$ for all finite type polynomials and hence for all polynomials because of the approximation property. Moreover, for each $f \in H^\infty(B_F)$ the sequence of the Cesàro sums $(\sigma_m f)$ of its Taylor series τ_0 converges to f and constitutes a bounded set in $H^\infty(B_F)$ ([M1, 5.2.c]),

thus $u(f) = \lim_m u(\sigma_m f) = \lim_m \sigma_m f(y) = f(y)$. Therefore $\delta_{\phi(x_\alpha)} \rightarrow \delta_y$ in $(G^\infty(B_F), \sigma_{G^\infty})$ or in other words, $\phi(x_\alpha) \rightarrow y$ in (B_F, σ_F) . \square

Proposition 4. *If $\phi(B_E) \subset rB_F$ for some $0 < r < 1$ and $\phi(B_E)$ is relatively compact in $(B_F, \sigma(F, H^\infty(B_F)))$, then $C_\phi : H^\infty(B_F) \rightarrow H^\infty(B_E)$ is weakly compact.*

Proof. Consider $C_\phi^t : H^\infty(B_E)^* \rightarrow H^\infty(B_F)^*$. Since $H^\infty(B_E)^* = G^\infty(B_E)^{**}$ we consider the continuous restriction map $C_\phi^t|_{G^\infty} : G^\infty(B_E) \rightarrow G^\infty(B_F)^{**}$. We have that $G^\infty(B_E)$ is the closed span of $\{\delta_x : x \in B_E\}$ in $H^\infty(B_E)^*$ and $C_\phi^t(\delta_x) = \delta_{\phi(x)}$, so it follows that $C_\phi^t(G^\infty(B_E)) \subset \overline{\text{span}}\{\delta_{\phi(x)} : x \in B_E\} \subset G^\infty(B_F)$. Thus the restriction map $S = C_\phi^t|_{G^\infty}$ can be considered as a map from $G^\infty(B_E)$ into $G^\infty(B_F)$. We first show that $S : G^\infty(B_E) \rightarrow G^\infty(B_F)$ is weakly compact. Let \overline{B}_{G^∞} be the closed unit ball in $G^\infty(B_E)$. Then $S(\overline{B}_{G^\infty}) \subset \overline{\Gamma}\{\delta_{\phi(x)} : x \in B_E\}$. By Krein's theorem it is enough to show that $\{\delta_{\phi(x)} : x \in B_E\}$ is relatively $\sigma(G^\infty(B_F), H^\infty(B_F))$ -compact. Since $\phi(B_E) \subset rB_F$, it follows that

$$\overline{\phi(B_E)}^{\sigma(F, H^\infty(B_F))} \subset B_F,$$

which is $\sigma(F, H^\infty(B_F))$ -compact, so $\delta_{\phi(B_E)}$ is relatively $\sigma(G^\infty(B_F), H^\infty(B_F))$ -compact.

Observe that for $f \in H^\infty(B_F)$ and $x \in B_E$, we have $\langle S^t(f), \delta_x \rangle = \langle f, S(\delta_x) \rangle = \langle f, C_\phi^t(\delta_x) \rangle = \langle C_\phi(f), \delta_x \rangle$. Therefore, $S^t(f) = C_\phi(f)$, so $S^t = C_\phi$. Since $S^t : G^\infty(B_F)^* \rightarrow G^\infty(B_E)^*$ is weakly compact, $C_\phi : H^\infty(B_F) \rightarrow H^\infty(B_E)$ is weakly compact. \square

Collecting the above propositions and recalling Lemma 1, we get

Theorem 5. *The composition operator $C_\phi : H^\infty(B_F) \rightarrow H^\infty(B_E)$ is weakly compact if (i) $\phi(B_E) \subset rB_F$ for some $0 < r < 1$ and (ii) $\phi(B_E)$ is relatively compact in $(F, \sigma(F, P(F)))$. The converse holds if moreover F has the approximation property.*

Corollary 6. *If F is a Λ -space with the approximation property, then every weakly compact composition operator $C_\phi : H^\infty(B_F) \rightarrow H^\infty(B_E)$ is compact.*

Corollary 7. *If $\phi(B_E)$ is a relatively weakly compact set strictly inside B_F , and $P(F) = P_{wsc}(F)$, where $P_{wsc}(F)$ is the subspace of $P(F)$ of all weakly sequentially continuous polynomials, then $C_\phi : H^\infty(B_F) \rightarrow H^\infty(B_E)$ is weakly compact.*

Proof. From any sequence in $\phi(B_E)$ we get a subsequence which is weakly convergent, hence $(F, \sigma(F, P(F)))$ -convergent. Thus $\phi(B_E)$ is relatively compact in the angelic space $(F, \sigma(F, P(F)))$. \square

Example 8. Should we have an infinite dimensional reflexive Banach space E , a Banach space F so that $P(F) = P_{wsc}(F)$ and an embedding $\phi : E \rightarrow F$ with $\|\phi\| < 1$, we will have found a weakly compact noncompact composition operator $C_\phi : H^\infty(B_F) \rightarrow H^\infty(B_E)$.

The Banach spaces (and their closed subspaces) c_0 , T^* , the original (reflexive) Tsirelson space, the Tsirelson*-James space, T_J^* , and $c_0 \times T^*$ have property P_α of Pelczynski for all $\alpha < 1$, so they fulfill the assumption regarding F in the above corollary ([AF, Corollary 3]). Also any Banach space F with the Dunford-Pettis property satisfies that $P(F) = P_{wsc}(F)$ [R1]. Therefore we have a bunch of triads (E, F, ϕ) fulfilling the quoted conditions. For instance, taking E any separable reflexive Banach space, $F = C([0, 1])$ -where E may be isometrically embedded since

$C([0,1])$ is universal among the separable Banach spaces- and ϕ any contraction of the embedding. Another example is provided by taking $\phi : x \in T^* \rightsquigarrow rx \in c_0$ ($0 < r < 1$), which is linear, continuous and weakly compact but noncompact (the unit ball of T^* contains the sequence of units of c_0) by the original construction of Tsirelson as can be seen in [HHZ]. The same will happen if we replace T^* by ℓ_2 .

Also, if $E \subset T^*$ is an infinite dimensional Banach space and $\phi : x \in E \rightsquigarrow rx \in T^*$ ($0 < r < 1$), then $C_\phi : H^\infty(B_{T^*}) \rightarrow H^\infty(B_E)$ is weakly compact, but noncompact.

Corollary 9. *If $P(^nF)$ is reflexive for all n , F has the approximation property and $\phi(B_E)$ is strictly inside B_F , then $C_\phi : H^\infty(B_F) \rightarrow H^\infty(B_E)$ is weakly compact.*

Proof. Since $P(^nF)$ is reflexive for all n , F is reflexive and $P(F) = P_{wsc}(F)$ holds when F has the approximation property (see [AAD], [R]). \square

We next describe the completely continuous composition operators C_ϕ .

Proposition 10. *The composition operator $C_\phi : H^\infty(B_F) \rightarrow H^\infty(B_E)$ is completely continuous if and only if (i) $\phi(B_E) \subset rB_F$ for some $0 < r < 1$ and (ii) $\phi(B_E)$ is a Dunford-Pettis set in F .*

Proof. Assume first that C_ϕ is completely continuous. By Proposition 2, $\phi(B_E)$ is strictly inside B_F . Let $(l_n) \subset F^*$ be a weak-null sequence. The map $J : F^* \rightarrow H^\infty(B_F)$, $l \mapsto l|_{B_F}$, is weak-weak continuous, so $(J(l_n))$ is a weak-null sequence in $H^\infty(B_F)$. Hence (l_n) converges to zero uniformly on $\phi(B_E)$, so $\phi(B_E)$ is a Dunford-Pettis set in F .

Conversely, suppose that the assumptions are satisfied. Take a weak-null sequence $(f_n) \subset H^\infty(B_F)$. Then for every m the sequence of their Taylor polynomials at 0, $(P_m f_n)_n \subset P(^mF)$, is also weakly null. Moreover it follows from [GG, Theorem 2.2] that $\phi(B_E) \otimes \cdots \otimes \phi(B_E)$ is a Dunford-Pettis set in the projective tensor product $F \otimes_\pi \cdots \otimes_\pi F$, thus the sequence $(P_m f_n)_n$ converges uniformly to 0 on $\phi(B_E)$.

Since $(\|f_n\|)$ is bounded, it follows from Cauchy inequalities that on every ball of radius less than 1 the approximation of each f_n by its Taylor series $\sum P_m f_n$ can be chosen independently of n . That is, for a given $\epsilon > 0$, there is $k \in \mathbb{N}$ such that $|\sum_{m=0}^k P_m f_n(x) - f_n(x)| < \epsilon$ for all $n \in \mathbb{N}$ and $x \in rB_F$. Once we have found k we also find, by the above, $n_0 \in \mathbb{N}$ so that $|P_m f_n(z)| < \frac{\epsilon}{k+1}$ for $n > n_0$ and for all $z \in \phi(B_E)$.

So, if $n > n_0$, then $|C_\phi(f_n)(x)| = |f_n(\phi(x))| \leq |\sum_{m=0}^k P_m f_n(\phi(x)) - f_n(\phi(x))| + |\sum_{m=0}^k P_m f_n(\phi(x))| \leq \|\sum_{m=0}^k P_m f_n - f_n\|_{rB_F} + \|\sum_{m=0}^k P_m f_n\|_{\phi(B_E)} < 2\epsilon$ for all $x \in B_E$. Hence $(\|C_\phi(f_n)\|)$ converges to 0. \square

Corollary 11. *Assume that F^* has the Schur property. The composition operator $C_\phi : H^\infty(B_F) \rightarrow H^\infty(B_E)$ is completely continuous if and only if $\phi(B_E)$ lies strictly inside B_F .*

Corollary 12. *If F has the Dunford-Pettis property as well as the approximation property, then every weakly compact composition operator $C_\phi : H^\infty(B_F) \rightarrow H^\infty(B_E)$ is completely continuous.*

Proof. By the Dunford-Pettis property of F , every weakly null sequence in F^* converges uniformly to 0 in $\phi(B_E)$ since it is a relatively weakly compact subset of

F by Proposition 3. Thus $\phi(B_E)$ is a Dunford-Pettis set lying strictly inside B_F (Proposition 2), so Proposition 10 leads to the result. \square

Remark 13. There is no characterization of Rosenthal composition operators similar to Proposition 10. Take $E = F = \ell_2$ and $\phi(x) = \frac{x}{2}$; then $\phi(B_E)$ is conditionally weakly compact. But $C_\phi : P(^2\ell_2) \rightarrow P(^2\ell_2)$ is not a Rosenthal operator, since if it were, the identity on $P(^2\ell_2)$, which coincides with $4C_\phi$, would also be a Rosenthal operator which is prevented by the fact that $P(^2\ell_2)$ contains a copy of ℓ^∞ [D1, Corollary 4].

Example 14. *A Rosenthal nonweakly compact composition operator.* Let $E = F = T_J^*$. Since E^{**} has the Radon-Nikodym property, E^* is an Asplund space, hence $P(^m E) = P_{w^*}(^m E^{**})$ is also Asplund ([V, Corollary 1.1]), so it does not contain any copy of ℓ_1 and therefore its bounded subsets are conditionally weakly compact. If we take $\phi(x) = \frac{x}{2}$, then C_ϕ cannot be weakly compact since $\phi(B_E) = \frac{B_E}{2}$ is not a weakly compact set with E a nonreflexive space.

On the other hand C_ϕ is a Rosenthal operator. Indeed: Put U for the unit ball in $H^\infty(B_E)$ and as in the proof of the above Proposition, for a given $\epsilon > 0$, there is $k \in \mathbb{N}$ such that $|\sum_{m=0}^k P_m f(x) - f(x)| < \epsilon$ for all $f \in U$ and $x \in \frac{1}{2}B_E$. So, $\|\sum_{m=0}^k \frac{P_m f}{2^m} - C_\phi(f)\| < \epsilon$ for all $f \in U$. Since each of the sets $K_m = \{\frac{P_m f}{2^m} : f \in U\}$ is conditionally weakly compact and $C_\phi(U) \subset \epsilon U + \sum_{m=0}^k K_m$, it follows that $C_\phi(U)$ is a conditionally weakly compact set (see [Di, Example 2, p. 237]).

REFERENCES

- [AAD] R. Alencar, R.M. Aron and S. Dineen, *A reflexive space of holomorphic functions in infinitely many variables*, Proc. Amer. Math. Soc. **90** (1984), 407-411. MR **85b**:46050
- [ACG] R.M. Aron, B.J. Cole and T.W. Gamelin, *Spectra of algebras of analytic functions on a Banach space*, J. reine angew. Math. **415** (1991), 51-93. MR **92f**:46056
- [AF] R. Alencar and K. Floret, *Weak-strong continuity of multilinear mappings and the Pelczynski -Pitt theorem*, J. of Math. Anal. and Appl. **206** (2) (1997), 532-546. MR **98h**:46045
- [AGL] R. Aron, P. Galindo and M. Lindström, *Compact homomorphisms between algebras of analytic functions*, Studia Math. **123** (3) (1997), 235-247. MR **98h**:46053
- [Di] J. Diestel, *Sequences and series in Banach spaces*, Springer, 1983. MR **85j**:46020
- [D] S. Dineen, *Complex analysis in locally convex spaces*, North Holland, 1981. MR **84b**:46050
- [D1] S. Dineen, *A Dvoretzky theorem for polynomials*, Proc. Amer. Math. Soc. **123** (1995), 2817-2821. MR **95k**:46021
- [GG] M. González and J. Gutiérrez, *Gantmacher type theorems for holomorphic mappings*, Math. Nachr. **186** (1997), 131-145. CMP 97:16
- [GGM] P. Galindo, D. Garcia and M. Maestre, *Entire functions of bounded type on Fréchet spaces*, Math. Nachr. **161** (1993), 185-198. MR **94k**:46086
- [GL] P. Galindo and M. Lindström, *Gleason parts and weakly compact homomorphisms between uniform Banach algebras*, Monatshefte für Math.
- [G] J. Garnett, *Bounded analytic functions*, Academic Press, 1981. MR **83g**:30037
- [HHZ] P. Habala, P. Hajek and V. Zizler, *Introduction to Banach spaces*, Charles University Prague, 1996.
- [JaP] J. A. Jaramillo and A. Prieto, *The weak-polynomial convergence in a Banach space*, Proc. Amer. Math. Soc. **118** (1993), 463-468. MR **93g**:46019
- [M1] J. Mujica, *Linearization of bounded holomorphic mappings on Banach spaces*, Trans. Amer. Math. Soc. **324** (1991), 867-887. MR **91h**:46088
- [M2] J. Mujica, *Complex analysis in Banach spaces*, North Holland, 1986. MR **88d**:46084
- [M3] J. Mujica, *Complex homomorphisms on the algebras of holomorphic functions on Fréchet spaces*, Math. Ann. **241** (1979), 73-82. MR **80j**:46074
- [N] K. Ng, *On a theorem of Dixmier*, Math. Scand. **29** (1971), 279-280. MR **49**:3504

- [R] R. Ryan, *Applications of topological tensor products in infinite dimensional holomorphy*, Ph. D. Thesis, Trinity College, Dublin (1980).
- [R1] R. Ryan, *Dunford-Pettis properties*, Bull. Acad. Polon. Sci. Ser. Sci. Math **27** (1979), 373-379. MR **80m**:46018
- [U] A. Ülger, *Some results about the spectrum of commutative Banach algebras under the weak topology and applications*, Monatshefte für Math. **121** (1996), 353-379. MR **98a**:46058
- [V] M. Valdivia, *Banach spaces of polynomials without copies of ℓ_1* , Proc. Amer. Math. Soc. **123** (1995), 3143-3150. MR **95m**:46070

DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, UNIVERSIDAD DE VALENCIA, 46100 BURJASOT, VALENCIA, SPAIN

E-mail address: `galindo@uv.es`

DEPARTMENT OF MATHEMATICS, ÅBO AKADEMI UNIVERSITY, FIN-20500 ÅBO, FINLAND

E-mail address: `mikael.lindstrom@abo.fi`

DEPARTMENT OF MATHEMATICS, UNIVERSITY COLLEGE GALWAY, GALWAY, IRELAND

E-mail address: `ray.ryan@ucg.ie`