

KD_∞ IS A CS-ALGEBRA

S. K. JAIN, P. KANWAR, S. MALIK, AND J. B. SRIVASTAVA

(Communicated by Ken Goodearl)

ABSTRACT. In this paper, it is shown that the group algebra KD_∞ is right CS if and only if $\text{char}(K) \neq 2$. Moreover, when $\text{char}(K) \neq 2$, then KD_∞ is also CS as a module over its center.

1. INTRODUCTION

Rings whose complement right ideals are direct summands are called right CS-rings. The class of CS-rings includes selfinjective rings, continuous rings etc. and have been of interest to many authors. However, there is hardly any literature on CS-group algebras. It is well known that the group algebra KG , where K is a field, is selfinjective if and only if G is a finite group. But the group algebra KG may be CS without the finiteness condition on the group G . For example if G is a torsion-free solvable-by-finite group, then KG is an Ore domain and hence is a CS-algebra. On the other hand if G is a finite group, then the group ring RG over the ring $R = M_n(\mathbb{Z})$ is not CS for any $n \geq 1$. It is, therefore, of interest to study when a given group algebra is CS. In this paper we study the group algebra $S = KD_\infty$ over a field K for its being CS or not. It is proved that S is a right CS-algebra if and only if $\text{char}(K) \neq 2$ (Theorem 3.6). It is further shown that the center $Z(S)$ of S is a Dedekind domain (Lemma 3.8) and that S is also a CS-module over $Z(S)$ (Theorem 3.9).

2. NOTATION AND PRELIMINARIES

Throughout, unless otherwise stated, K will denote a field and D_∞ , the infinite dihedral group, that is, the group generated by two elements a and b where a is of infinite order, b is of order 2 and $ab = ba^{-1}$. A module will always mean a right unital module. A nonzero submodule N of a module M is said to be essential in M , denoted by $N \subseteq_e M$, if, for every nonzero submodule L of M , $L \cap N \neq 0$. N is called closed in M if N has no proper essential extensions in M . A module M is said to be CS if every nonzero submodule of M is essential in a summand of M , or equivalently, if every closed submodule of M is a summand of M . CS-modules are also commonly known as extending modules ([1]). A ring R is called right CS if it is CS as a right module over itself.

Received by the editors April 10, 1998.

1991 *Mathematics Subject Classification*. Primary 16S14, 13B21, 13C10, 13F05.

Key words and phrases. Group algebra, CS-ring, Dedekind domain.

If A is an algebra over a ring R , then an element $u \in A$ is called integral over R if it satisfies a polynomial equation with coefficients in R and leading coefficient 1. A is called integral if all its elements are integral.

3. GROUP RING KD_∞

Throughout this section, S will denote the group algebra KD_∞ , and R will denote the group algebra KA where $A = \langle a \rangle$. It is well known that KA is a PID ([5], Exercise 2, p.28). Also, by ([6], Theorem 5.1 and [3], Proposition 9, p.165), S is a prime PI-ring. We begin with some lemmas which will be useful to prove our main result. Our first lemma is well known. We state it here without proof for convenience.

Lemma 3.1 ([1], Corollary 12.8). *For a commutative domain R , the following are equivalent:*

- (a) R is a Prüfer domain.
- (b) $(R \oplus R)_R$ is a CS-module.

Lemma 3.2. S is CS as a right R -module.

Proof. Since $S = R \oplus Rb \simeq R \times R$ and R is a Prüfer domain, the result follows by Lemma 3.1. \square

Lemma 3.3. *If $\text{char}(K) \neq 2$ and if U is a right ideal of S such that $S_R = U_R \oplus X_R$ for some R -submodule X of the right R -module S_R , then there exists a right ideal V of S such that $S = U \oplus V$.*

Proof. Let π_1, π_2 be the projections of S_R onto U_R and X_R respectively. Define $\gamma : S \rightarrow S$ by $\gamma(s) = \frac{1}{2}[\pi_2(s) + \pi_2(sb)b]$. Since $\text{char}(K) \neq 2$, γ is well-defined. Clearly, $\gamma(s_1 + s_2) = \gamma(s_1) + \gamma(s_2)$. Also,

$$\begin{aligned} \gamma(sa) &= \frac{1}{2}[\pi_2(sa) + \pi_2(sab)b] = \frac{1}{2}[\pi_2(s)a + \pi_2(sba^{-1})b] \\ &= \frac{1}{2}[\pi_2(s)a + \pi_2(sb)a^{-1}b] = \frac{1}{2}[\pi_2(s)a + \pi_2(sb)ba] = \gamma(s)a. \end{aligned}$$

Similarly, $\gamma(sb) = \gamma(s)b$. Thus $\gamma \in \text{Hom}_S(S, S)$. Let $V = \gamma(S)$. Then V is a right ideal of S . We will prove that $S = U \oplus V$. So, let $s \in S$. Write $s = (s - \gamma(s)) + \gamma(s)$. Since

$$\begin{aligned} s - \gamma(s) &= s - \frac{1}{2}[\pi_2(s) + \pi_2(sb)b] = \frac{1}{2}[(s - \pi_2(s)) + (s - \pi_2(sb)b)] \\ &= \frac{1}{2}[(s - \pi_2(s)) + (sb - \pi_2(sb))b] = \frac{1}{2}[\pi_1(s) + \pi_1(sb)b], \end{aligned}$$

and U is a right ideal of S , we have $s - \gamma(s) \in U$. Thus $S = U + V$. Also since for every $s \in S$, $s - \gamma(s) \in U$, we have $\gamma(s - \gamma(s)) = 0$, that is, $\gamma(s) = \gamma^2(s)$ for every $s \in S$.

To prove $U \cap V = (0)$, let $x \in U \cap V$. Then $x = \gamma(s)$ for some $s \in S$ and $\gamma(x) = 0$. Thus $\gamma^2(s) = 0$ and consequently, $x = \gamma(s) = \gamma^2(s) = 0$, as desired. \square

Lemma 3.4. *If U is a closed right ideal of S , then U is a closed submodule of the right R -module S_R .*

Proof. Suppose $x \in cl(U_R)$, the closure of U_R in S_R . Then $x \in S$ and $xE \subset U$ for some essential right ideal E of R . Consequently, $x(ES) \subset US \subset U$. Also ES is an essential right ideal of S ([5], Exercise 27, p. 467). Thus $x \in cl(U_S) = U$, because U is a closed right ideal of S . This completes the proof. \square

Lemma 3.5. *If $char(K) = 2$, then S has no nontrivial idempotents.*

Proof. For $\alpha = \sum k_i a^i \in R$, let $\alpha^* = \sum k_i a^{-i}$. Since $ab = ba^{-1}$, it follows that $ab = b\alpha^*$ for every $\alpha \in R$. Now if $\alpha + b\beta \in S$ is a nontrivial idempotent in S , then using $\alpha b = b\alpha^*$ and $(\alpha + b\beta)^2 = \alpha + b\beta$ we get $\alpha^2 + \beta^* \beta + b(\alpha^* \beta + \beta \alpha) = \alpha + b\beta$. Thus $\alpha^2 + \beta^* \beta = \alpha$ and $\alpha^* \beta + \beta \alpha = \beta$. Since R is a PID and $\alpha + b\beta$ is an idempotent in S , $\beta \neq 0$. Consequently, using R is a domain, the relation $\alpha^* \beta + \beta \alpha = \beta$ yields $\alpha^* + \alpha = 1$. Since $\alpha \in R$, $\alpha = \sum k_i a^i$ where $k_i \in K$. Since $\alpha^* + \alpha = 1$ and $char(K) = 2$, we have $0 = 1$, a contradiction. Thus S has no nontrivial idempotents. \square

Theorem 3.6. *S is a right CS-ring if and only if $char(K) \neq 2$.*

Proof. First assume that $char(K) \neq 2$. Let U be a closed right ideal of S . By Lemma 3.4, U_R is a closed submodule of the right R -module S_R . Since S_R is CS (Lemma 3.2), U_R is a direct summand of S_R . But then by Lemma 3.3, U is a summand of S . Hence S is a right CS-ring. Conversely, let S be right CS. If possible, let $char(K) = 2$. By Lemma 3.5, S has no nontrivial idempotents. Since S is right CS, every nonzero right ideal of S is essential in S . Thus S and hence the right maximal quotient ring $Q_{max}^r(S)$ of S is uniform. Since S is right nonsingular, it follows that $Q_{max}^r(S)$ is a division ring. Hence S is a domain, a contradiction because $(1 + b)^2 = 0$ and $1 + b \neq 0$. Thus $char(K) \neq 2$. \square

In what follows $Z(S)$ will denote the center of the ring S . Unless otherwise stated $char(K) \neq 2$ and $e = \frac{1}{2} + \frac{1}{2}b$. Notice that e is an idempotent in S . For $\alpha = \sum k_i a^i$, we will write $\alpha^* = \sum k_i a^{-i}$. In the following lemma we determine $Z(S)$.

Lemma 3.7. *For any field K , $Z(S) = \{\alpha \in R \mid \alpha = \alpha^*\}$.*

Proof. Clearly, $\{\alpha \in R \mid \alpha = \alpha^*\} \subset Z(S)$. To prove the reverse inclusion, let $s = \alpha + \beta b \in Z(S)$. Then $sx = xs$ for every $x \in S$. In particular, $sa = as$ and $sb = bs$. Now $sa = as$ gives $\alpha a + \beta ba = a\alpha + a\beta b$. Since $\alpha a = a\alpha$, we get $\beta a^{-1}b = a\beta b$, that is, $(a^2 - 1)\beta = 0$. Thus, $\beta = 0$. Consequently, $s = \alpha \in R$. Again as $sb = bs$, we have $ab = b\alpha = \alpha^*b$. Thus $\alpha = \alpha^*$ and the proof is complete. \square

Lemma 3.8. *$Z(S)$ is a Dedekind domain.*

Proof. Clearly, $Z(S) \simeq eZ(S) = eSe$. Since S is right noetherian, eSe is right noetherian ([7], Lemma 2.7.12). Thus $Z(S)$ is right noetherian. Since S is a prime PI ring, by ([4], Corollary 6.14, p.467), S is a finitely generated $Z(S)$ -module. Let $S = s_1Z(S) + s_2Z(S) + \dots + s_kZ(S)$ and let for $1 \leq i \leq k$, $s_i = \alpha_i + \beta_i b$ where $\alpha_i, \beta_i \in R$. Then $R \oplus Rb = (\alpha_1 + \beta_1 b)Z(S) + (\alpha_2 + \beta_2 b)Z(S) + \dots + (\alpha_k + \beta_k b)Z(S)$. Consequently, $R = \alpha_1Z(S) + \alpha_2Z(S) + \dots + \alpha_kZ(S)$, that is, R is a finitely generated noetherian $Z(S)$ -module. Thus by ([2], Theorem 17) R is integral over $Z(S)$, that is, $R|_{Z(S)}$ is an integral extension of $Z(S)$.

Let L denote the quotient field of $Z(S)$. We will show that $R \cap L = Z(S)$. Let $\gamma \in R \cap L$. Then $\gamma = \alpha\beta^{-1}$ for some $\alpha, \beta \in Z(S)$. Thus $\alpha = \gamma\beta$. Since

$\alpha, \beta \in Z(S)$, $\alpha = \alpha^*$, $\beta = \beta^*$. Hence $\alpha = \alpha^* = (\gamma\beta)^* = \beta^*\gamma^* = \beta\gamma^*$ so that $\gamma^* = \alpha\beta^{-1} = \gamma$. Since $\gamma \in R$, we have $\gamma \in Z(S)$. Hence $R \cap L = Z(S)$. Since R is a Dedekind domain, it follows, by ([8], Theorem 20, p.283), that $Z(S)$ is a Dedekind domain \square

Theorem 3.9. *R and S are CS as right $Z(S)$ -modules.*

Proof. Clearly, $Z(S) \cap aZ(S) = (0)$. Further, since $a^{-n} = (a^n + a^{-n}) - a^n$ and $a^n = a^{n-1}(a + a^{-1}) - a^{n-2}$ we have $R = Z(S) \oplus aZ(S) \simeq Z(S) \times Z(S)$. Also as $S = R \oplus bR$, we have $S \simeq Z(S) \times Z(S) \times Z(S) \times Z(S)$. By Theorem 3.8, $Z(S)$ is a Dedekind domain. The result now follows by Lemma 3.1. \square

Remark 1. The uniform dimension of R as a $Z(S)$ -module is 2 and that of S as a $Z(S)$ -module is 4.

ACKNOWLEDGMENT

S. K. Jain would like to thank Indian Institute of Technology, Delhi and its Department of Mathematics for their hospitality during his visit to the Institute in November–December 1997. P. Kanwar would also like to express his thanks to Professor J. B. Srivastava for providing him an opportunity to join in the project.

REFERENCES

- [1] Nguyen Viet Dung, Dinh Van Huynh, Patrick F. Smith and Robert Wisbauer, *Extending Modules*, Pitman, London, 1994. MR **96f**:16008
- [2] Irving Kaplansky, *Commutative Rings*, Allyn and Bacon, Inc., Boston, 1970. MR **40**:7234
- [3] Joachim Lambek, *Lectures on Rings and Modules*, Blaisdell Publishing Company, London, 1966. MR **34**:5857
- [4] J. C. McConnell and J. C. Robson, *Noncommutative Noetherian Rings*, Wiley-Interscience, NY, 1987. MR **89j**:16023
- [5] Donald S. Passman, *The Algebraic Structure of Group Rings*, John Wiley, NY, 1977. MR **81d**:16001
- [6] Donald S. Passman, *Infinite Group Rings*, Marcel Dekker, NY, 1971. MR **47**:3500
- [7] Louis Rowen, *Ring Theory, vol I*, Academic Press, NY, 1988. MR **89h**:16001
- [8] Oscar Zariski and Pierre Samuel, *Commutative Algebra, vol. I*, D. Van Nostrand, NY, 1958. MR **19**:833e

DEPARTMENT OF MATHEMATICS, OHIO UNIVERSITY, ATHENS, OHIO 45701

E-mail address: jain@math.ohiou.edu

E-mail address: pkanwar@math.ohiou.edu

Current address, P Kanwar: Division of Mathematics and Computer Science, Truman State University, Kirksville, Missouri 63501

DEPARTMENT OF MATHEMATICS, HINDU COLLEGE, DELHI-110007, INDIA

E-mail address: sbm@csec.ernet.in

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY, NEW DELHI-110016, INDIA

E-mail address: jbsrivas@maths.iitd.ernet.in