

THE GLOBALLY IRREDUCIBLE REPRESENTATIONS OF SYMMETRIC GROUPS

ALEXANDER KLESHCHEV AND ALEXANDER PREMET

(Communicated by Ronald M. Solomon)

ABSTRACT. Let K be an algebraic number field and \mathcal{O} be the ring of integers of K . Let G be a finite group and M be a finitely generated torsion free $\mathcal{O}G$ -module. We say that M is a *globally irreducible* $\mathcal{O}G$ -module if, for every maximal ideal \mathfrak{p} of \mathcal{O} , the $k_{\mathfrak{p}}$ G -module $M \otimes_{\mathcal{O}} k_{\mathfrak{p}}$ is irreducible, where $k_{\mathfrak{p}}$ stands for the residue field \mathcal{O}/\mathfrak{p} .

Answering a question of Pham Huu Tiep, we prove that the symmetric group Σ_n does not have non-trivial globally irreducible modules. More precisely we establish that if M is a globally irreducible $\mathcal{O}\Sigma_n$ -module, then M is an \mathcal{O} -module of rank 1 with the trivial or sign action of Σ_n .

1. INTRODUCTION AND GENERALITIES

The notion of global irreducibility was first introduced by J. G. Thompson in the course of constructing the sporadic group F_3 ([9, 10]). He observed that there exists an even unimodular \mathbb{Z} -lattice Λ of rank 248 with $\text{Aut}(\Lambda) \cong \mathbb{Z}_2 \times F_3$ such that the $\mathbb{F}_p F_3$ -module $\Lambda/p\Lambda$ is irreducible for every prime p . Other interesting examples of the same nature were discovered later, and different (closely related) notions of global irreducibility were introduced. A good survey on the subject is [11].

Let K be an algebraic number field and \mathcal{O} be the ring of integers of K . Let G be a finite group and M be a finitely generated torsion free $\mathcal{O}G$ -module. We say that M is *globally irreducible* if, for every maximal ideal \mathfrak{p} of \mathcal{O} , the $k_{\mathfrak{p}}$ G -module $M \otimes_{\mathcal{O}} k_{\mathfrak{p}}$ is irreducible, where $k_{\mathfrak{p}}$ denotes the residue field \mathcal{O}/\mathfrak{p} . This definition seems to “include” all versions of global irreducibility introduced before.

Let Σ_n denote the symmetric group on n letters. The main result of this paper is the following theorem which answers a question posed by Pham Huu Tiep.

Theorem A. *Let M be a globally irreducible $\mathcal{O}\Sigma_n$ -module. Then M is an \mathcal{O} -module of rank 1 with the trivial or sign action of Σ_n .*

Throughout the paper:

$\mathbb{N} = \{1, 2, \dots\}$ is the set of all natural numbers,

F is an arbitrary field of characteristic $p > 0$ (unless otherwise stated).

Received by the editors December 10, 1997 and, in revised form, April 15, 1998.

1991 *Mathematics Subject Classification.* Primary 20C30, 20C10.

Key words and phrases. Symmetric group, Specht module.

The authors thank G. Michler and A. Zalesskii who organized a conference on representations of finite groups in Bad-Honnef where this collaboration began, and the Volkswagen foundation for financial support. The first author was also supported by the NSF.

For $r, s \in \mathbb{N}$ we denote:

$$[r, s] = \{t \in \mathbb{N} \mid r \leq t \leq s\}; \quad (r, s] = \{t \in \mathbb{N} \mid r < t \leq s\};$$

$$[r, s) = \{t \in \mathbb{N} \mid r \leq t < s\}; \quad (r, s) = \{t \in \mathbb{N} \mid r < t < s\}.$$

Given a partition λ of n , we often gather together equal parts of λ and write

$$\lambda = (\lambda_1^{a_1}, \lambda_2^{a_2}, \dots, \lambda_k^{a_k}),$$

where $\lambda_1 > \lambda_2 > \dots > \lambda_k > 0$, $a_i > 0$, and $a_1\lambda_1 + a_2\lambda_2 + \dots + a_k\lambda_k = n$. A partition λ is called *p-regular* if $a_i < p$ for all i . We denote by λ' the partition conjugate to λ (the Young diagram of λ' is the transpose of the Young diagram of λ). The symbol (i, j) denotes the node in the i th row and the j th column of the Young diagram of λ , and $h(i, j) = h_\lambda(i, j)$ stands for the length of the hook of λ corresponding to the node (i, j) . Note that the notation for the *node* (i, j) should not be confused with the one for the *interval* (i, j) as defined above! In what follows we do not distinguish between partitions and their Young diagrams.

We introduce some notions from the representation theory of symmetric groups. Details can be found in [6]. Let $S_{\mathbb{Z}}^\lambda$ denote the Specht module over \mathbb{Z} corresponding to λ (it carries a canonical $\mathbb{Z}\Sigma_n$ -module structure). Given a field F set $S_F^\lambda = S_{\mathbb{Z}}^\lambda \otimes_{\mathbb{Z}} F$. It is well-known that if $\text{char } F = 0$, then S_F^λ is an irreducible $F\Sigma_n$ -module. In general, the irreducible $F\Sigma_n$ -module corresponding to a p -regular partition λ is denoted by D_F^λ . Note that D_F^λ is always a quotient of S_F^λ . Another fact which is going to be used without comments is that \mathbb{F}_p is the splitting field for the group Σ_n . So the question of the irreducibility of S_F^λ depends on the characteristic of F only, not on F itself.

We shall deduce Theorem A from the following

Theorem B. *The Specht module $S_{\mathbb{Z}}^\lambda$ is globally irreducible if and only if $\lambda = (n)$ or (1^n) .*

We now explain why Theorem B implies Theorem A.

Let M be a globally irreducible $\mathcal{O}\Sigma_n$ -module, and $M_K = M \otimes_{\mathcal{O}} K$. Then M_K is a direct sum of Specht modules over K (see [6, Theorem 4.12]). It follows that there exists a $\mathbb{Z}\Sigma_n$ -module $M_{\mathbb{Z}}$ such that $M_K \cong M_{\mathbb{Z}} \otimes_{\mathbb{Z}} K$ as $K\Sigma_n$ -modules. Moreover, $M_{\mathbb{Z}}$ can be chosen to be a direct sum of Specht modules over \mathbb{Z} . By a theorem of Brauer–Nesbitt, the $k_{\mathfrak{p}}\Sigma_n$ -modules $M \otimes_{\mathcal{O}} k_{\mathfrak{p}}$ and $M_{\mathbb{Z}} \otimes_{\mathbb{Z}} k_{\mathfrak{p}}$ have the same composition factors (see, e.g., [2, Theorem 82.1]). We claim that $M_{\mathbb{Z}}$ is a globally irreducible $\mathbb{Z}\Sigma_n$ -module isomorphic to a Specht module $S_{\mathbb{Z}}^\lambda$ for some partition λ of n . Indeed, if $\bar{M} = M_{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{F}_p$ is reducible for some prime p , then, for each maximal ideal \mathfrak{q} of \mathcal{O} such that $\mathfrak{q} \cap \mathbb{Z} = p\mathbb{Z}$, the $k_{\mathfrak{q}}\Sigma_n$ -module $M_{\mathbb{Z}} \otimes_{\mathbb{Z}} k_{\mathfrak{q}} \cong \bar{M} \otimes_{\mathbb{F}_p} k_{\mathfrak{q}}$ is reducible. Also, if $M_{\mathbb{Z}}$ is a direct sum of more than one Specht module over \mathbb{Z} , then, for each maximal ideal \mathfrak{p} of \mathcal{O} , the $k_{\mathfrak{p}}\Sigma_n$ -module $M_{\mathbb{Z}} \otimes_{\mathbb{Z}} k_{\mathfrak{p}}$ is reducible again. This establishes the claim in view of the global irreducibility of M and the result of Brauer–Nesbitt mentioned above.

Now suppose we have proved Theorem B. Then the \mathbb{Z} -rank of $M_{\mathbb{Z}}$ is 1. So $\dim M_K = 1$, i.e. M is an \mathcal{O} -module of rank 1. Hence the subgroup $A_n = \Sigma'_n$ acts trivially on M , and Theorem A follows.

2. SOME REPRESENTATION THEORY AND SOME NUMBER THEORY

The following result known as Carter’s conjecture was proved by James and Murphy [5, 7]. Recall that the p -exponent of an integer n is the largest power of the prime p dividing n .

Proposition 2.1. *Let λ be a p -regular partition of n . Then S_F^λ is irreducible if and only if the p -exponents of the hook lengths of the Young diagram of λ are constant along the columns.*


Our first lemma is a straightforward consequence of Proposition 2.1.

Lemma 2.2. *Assume that both λ and λ' are p -regular partitions of n . If there is a hook in the Young diagram of λ whose length is divisible by p , then S_F^λ is reducible.*

Proof. Suppose that S_F^λ is irreducible and some $h_\lambda(i, j)$ is divisible by p . In view of Proposition 2.1 all hook lengths in the j th column of the Young diagram of λ should be divisible by p . But this is impossible for the bottom hook in the j th column, since λ' is p -regular. \square

From now on we fix an arbitrary partition λ of n . Let

$$\lambda = (\lambda_1^{a_1}, \lambda_2^{a_2}, \dots, \lambda_k^{a_k}), \quad \lambda' = (\mu_1^{b_1}, \mu_2^{b_2}, \dots, \mu_k^{b_k}).$$



Put

$$A_s = \left(\sum_{i=1}^s a_i, \lambda_s \right), \quad s = 1, 2, \dots, k;$$

$$B_r = \left(\sum_{i=1}^{r-1} a_i + 1, \lambda_r + 1 \right), \quad r = 1, 2, \dots, k + 1$$

(λ_{k+1} is interpreted as 0). Then A_1, \dots, A_k and B_1, \dots, B_{k+1} are all the removable and the addable nodes for λ , respectively. We denote by $\lambda(r, s)$ the partition of n whose Young diagram is obtained from that of λ by removing A_s and adding B_r . This partition is *not* well defined exactly in the following cases:

- (a) $r = s, a_r = 1$;
- (b) $r = s + 1, b_s = 1$.

So when we write $\lambda(r, s)$ we assume that neither (a) nor (b) occurs. Clearly, $\lambda(r, s) \triangleright \lambda$ if $r \leq s$ and $\lambda(r, s) \triangleleft \lambda$ if $r > s$.

We make use of the so-called *p-ladders* introduced by James in [4]. Those are, roughly speaking, straight lines with slope $(p - 1)$. More precisely, the u -th p -ladder is defined as the following set of nodes:

$$L_u := \{(i, j) \mid i, j \in \mathbb{N}, i + (p - 1)(j - 1) = u\}.$$

If we slide the nodes of λ along the p -ladders as far up as they can go, then we obtain a new partition λ^R , called the *p-regularization* of λ . It is proved in [4] that λ^R is well defined and that λ is p -regular if and only if $\lambda = \lambda^R$.

Proposition 2.3 ([4]). *$D_F^{\lambda^R}$ occurs in S_F^λ with multiplicity 1, and if D_F^μ is a composition factor of S_F^λ , then $\mu \supseteq \lambda^R$.*

Lemma 2.4. *Suppose that λ and $\lambda(r, s)$ are p -regular. Let A_s belong to the v -th p -ladder, and B_r belong to the u -th p -ladder. Then $\lambda(r, s) \triangleright \lambda$ if and only if $u > v$.*

Proof. It was mentioned above that $\lambda(r, s) \triangleright \lambda$ if and only if $r \leq s$. By definition,

$$v = \sum_{i=1}^s a_i + (p - 1)(\lambda_s - 1) \quad \text{and} \quad u = \sum_{i=1}^{r-1} a_i + 1 + (p - 1)\lambda_r.$$

Now $u > v$ is equivalent to

$$(1) \quad \sum_{i=1}^s a_i - \sum_{i=1}^{r-1} a_i < (p - 1)(\lambda_r - \lambda_s) + p.$$

Suppose $r > s$ and $u > v$. Then

$$(2) \quad \sum_{i=s+1}^{r-1} a_i > (p - 1)(\lambda_s - \lambda_r) - p.$$

Note that $(p - 1)(\lambda_s - \lambda_r) \geq (p - 1)(r - s)$, and

$$\sum_{i=s+1}^{r-1} a_i \leq (r - 1 - s)(p - 1),$$

since λ is p -regular. Therefore (2) holds if and only if $a_i = p - 1$ for $s + 1 \leq i \leq r - 1$ and $\lambda_s - \lambda_r = r - s$. If $r = s + 1$, this gives $b_s = \lambda_s - \lambda_{s+1} = 1$, so $\lambda(r, s)$ is not defined. If $r > s + 1$, we get $a_{r-1} = p - 1$, and $b_{r-1} = 1$. Thus $\lambda(r, s)$ is not p -regular, giving a contradiction. Hence $r > s$ implies $u \leq v$.

Let $r \leq s$. Then

$$\sum_{i=r}^s a_i \leq (s - r + 1)(p - 1) < (s - r)(p - 1) + p \leq (\lambda_r - \lambda_s)(p - 1) + p,$$

which in combination with (1) forces $u > v$. Thus we have proved that $r > s$ implies $u \leq v$ and $r \leq s$ implies $u > v$, as desired. \square

Lemma 2.5. *Suppose A_s belongs to the v -th p -ladder and B_r belongs to the u -th p -ladder.*

- (i) *If $u = v$, then $\lambda(r, s)^R = \lambda^R$.*
- (ii) *If $u \neq v$, then $\lambda(r, s)^R$ is obtained from λ^R by moving the bottom removable node of λ^R on the p -ladder L_v to the top addable node for λ^R on the p -ladder L_u (and such nodes exist).*

Proof. This follows from the definition of p -regularization. □

Denote

$$A_{r,s} = \sum_{i=r}^s a_i, \quad B_{r,s} = \sum_{j=r}^s b_j.$$

(The sums are interpreted as 0 if $r > s$.)

Lemma 2.6. *Let $r \leq s$. Then $\lambda(r, s)^R \triangleright \lambda^R$ if and only if $A_{r,s} < p + (p - 1)B_{r,s-1}$.*

Proof. Let A_s belong to the v -th p -ladder and B_r belong to the u -th p -ladder. By Lemma 2.5, $\lambda(r, s)^R$ is obtained from λ^R by moving a removable node on the v -th p -ladder to an addable node on the u -th p -ladder. Since both λ^R and $\lambda(r, s)^R$ are p -regular, Lemma 2.4 implies that $\lambda(r, s)^R \triangleright \lambda^R$ if and only if $u > v$, which is equivalent to

$$A_{1,s} + (p - 1)(\lambda_s - 1) < A_{1,r-1} + 1 + (p - 1)\lambda_r.$$

Now the result follows from the equality $\lambda_r - \lambda_s = \sum_{j=r}^{s-1} b_j$. □

The following proposition is a special case of the main result of Carter–Payne [1].

Proposition 2.7 ([1]). (i) *Let $1 \leq r \leq s \leq k$. If p divides $A_{r,s} + B_{r,s-1}$, then $\text{Hom}_{F\Sigma_n}(S_F^{\lambda(r,s)}, S_F^\lambda) \neq 0$.*

(ii) *Assume that $k \geq 2$, $b_2 \geq 2$, and p is odd. Let*

$$\mu = (\lambda_1 + 2, \lambda_1^{a_1-1}, \lambda_2^{a_2-1}, \lambda_2 - 2, \lambda_3^{a_3}, \dots, \lambda_k^{a_k}).$$

If p divides $a_1 + a_2 + b_1 + 1$, then $\text{Hom}_{F\Sigma_n}(S_F^\mu, S_F^\lambda) \neq 0$.

Our further arguments rely heavily on the following consequence of Proposition 2.7.

Lemma 2.8. *Let λ be a partition of n .*

- (i) *Suppose there exist $r, s \in \mathbb{N}$ with $1 \leq r < s \leq k$ such that p divides $A_{r,s} + B_{r,s-1}$ and $A_{r,s} < (p - 1)B_{r,s-1} + p$. Then S_F^λ is reducible.*
- (ii) *Suppose that $k \geq 2$, $b_2 \geq 2$, $a_1 + a_2 < (p - 1)(b_1 + 2) + 1$, and p is odd. If p divides $a_1 + a_2 + b_1 + 1$, then S_F^λ is reducible.*

Proof. (i) By Proposition 2.7(i),

$$\text{Hom}_{F\Sigma_n}(S_F^{\lambda(r,s)}, S_F^\lambda) \neq 0.$$

By Lemma 2.6, our assumption guarantees that $\lambda(r, s)^R \triangleright \lambda^R$. If S_F^λ is irreducible, then it is isomorphic to $D_F^{\lambda^R}$ (see Proposition 2.3). On the other hand, by the same proposition, each composition factor D_F^μ of $S_F^{\lambda(r,s)}$ satisfies $\mu \triangleright \lambda(r, s)^R \triangleright \lambda^R$; hence the Hom-space above should be zero. This contradiction proves (i).

For (ii), let μ be as in Proposition 2.7(ii). Then the Young diagram of μ can be obtained from that of λ by moving two nodes from the $(a_1 + a_2)$ th row of the Young diagram of λ to the first row. Observe that μ is well defined as $b_2 \geq 2$. By Proposition 2.7(ii),

$$\text{Hom}_{F\Sigma_n}(S_F^\mu, S_F^\lambda) \neq 0.$$

Let $C = (a_1 + a_2, \lambda_2)$, $D = (a_1 + a_2, \lambda_2 - 1)$, $X = (1, \lambda_1 + 1)$, and $Y = (1, \lambda_1 + 2)$. The Young diagram of μ can be obtained from that of λ in two steps. First we

remove C and add X to get a partition ν . Then we remove D from the Young diagram of ν and add Y . By our assumption,

$$a_1 + a_2 + b_1 + 1 < p(b_1 + 1) + p.$$

As p divides $a_1 + a_2 + b_1 + 1$, we must have

$$a_1 + a_2 + b_1 + 1 \leq p(b_1 + 1)$$

whence $a_1 + a_2 < (p - 1)b_1 + p$. So $\nu^R \triangleright \lambda^R$ by Lemma 2.6 (one should keep in mind that $\nu = \lambda(1, 2)$). Applying Lemma 2.6 for the second time we obtain $\mu^R \triangleright \nu^R \triangleright \lambda^R$. Now we finish the proof as in part (i). \square

The next three lemmas are versions (and generalizations) of the well-known Chebyshev’s results on prime numbers.

Lemma 2.9. *Let $n \in \mathbb{N}$. Then the interval $(n, 2n + 1]$ contains at least two primes.*

Proof. It is known [3, p. 21] that, for $n \geq 21$, the number of primes on the interval $(n, 2n + 1]$ exceeds $3n/(5 \ln n)$. Clearly, $3n/(5 \ln n)$ is greater than 2 if $n \geq 21$. Now check the result for $n = 1, 2, \dots, 20$. \square

Lemma 2.10. *Let n be an integer greater than 1. Then $(n, 2n)$ contains a prime.*

Proof. This is the famous “Bertrand’s postulate” proved by Chebyshev (see, e.g., [3, p. 21]). \square

Lemma 2.11. *Let $n \in \mathbb{N}$. Then $(n, \frac{3}{2}n + 2]$ contains a prime.*

Proof. For $n \geq 25$, the interval $(n, \frac{6}{5}n)$ contains a prime (see [8, p. 192]). Now check the result for $n = 1, 2, \dots, 24$. \square

Remark. It is easy to see that Lemma 2.9 also follows from the fact that the interval $(n, \frac{6}{5}n)$ contains a prime ([8, p. 192]).

3. PROOF OF THEOREM B

Given a partition λ of n set

$$A = \max_{1 \leq i \leq k} \{a_i\}, \quad B = \max_{1 \leq j \leq k} \{b_j\}$$

(the parameters a_i , b_j , and k are defined in Section 2). We wish to prove that $S_{\mathbb{Z}}^{\lambda}$ is globally irreducible if and only if $\lambda \in \{(n), (1^n)\}$. It is well-known that $(S_F^{\lambda})^* \cong (S_F^{\lambda'}) \otimes \text{sgn}$. Therefore, $S_{\mathbb{Z}}^{\lambda}$ is globally irreducible if and only if $S_{\mathbb{Z}}^{\lambda'}$ is. So no generality is lost by assuming that $B \geq A$. Let $f \in \{1, 2, \dots, k\}$ be such that $b_f = B$.

Lemma 3.1. *Suppose that $S_{\mathbb{Z}}^{\lambda}$ is globally irreducible, and $B \geq A$. Then any integer in the interval $[1, A_{1,f} + B_{1,f} - 1]$ is a hook length of λ .*

Proof. We prove by the downward induction on $d = f, f - 1, \dots, 1$ that any integer in the interval $[1, A_{d,f} + B_{d,f} - 1]$ is a hook length of λ .

If $d = f$, we have $A_{d,f} + B_{d,f} - 1 = a_f + b_f - 1$. In this case, it suffices to note that $h(A_{1,f}, B_{f,k} - v) = v + 1$, $v = 0, 1, \dots, b_f - 1$, and $h(A_{1,f} - w, B_{f+1,k} + 1) = b_f + w$, $w = 0, 1, \dots, a_f - 1$.

Now suppose that $1 \leq d < f$. By the inductive hypothesis, the hook lengths of λ cover the interval $[1, A_{d+1,f} + B_{d+1,f} - 1]$. Note that

$$h(A_{1,d}, B_{f,k} - v) = A_{d+1,f} + B_{d,f-1} + 1 + v, \quad v = 0, 1, \dots, b_f - 1,$$

and

$$h(A_{1,d} - w, B_{f+1,k} + 1) = A_{d+1,f} + B_{d,f} + w, \quad w = 0, 1, \dots, a_d - 1.$$

Thus each number in the union

$$(3) \quad [1, A_{d+1,f} + B_{d+1,f} - 1] \cup [A_{d+1,f} + B_{d,f-1} + 1, A_{d,f} + B_{d,f} - 1]$$

is a hook length of λ . If $A_{d+1,f} + B_{d+1,f} > A_{d+1,f} + B_{d,f-1}$, the union (3) equals the interval $[1, A_{d,f} + B_{d,f} - 1]$, as desired. Otherwise we have

$$A_{d+1,f} + B_{d+1,f} \leq A_{d+1,f} + B_{d,f-1}$$

which is equivalent to $b_f \leq b_d$. Since $b_f = B$, we conclude that $b_d = b_f = B$ yielding

$$A_{d+1,f} + B_{d+1,f} = A_{d+1,f} + B_{d,f-1} =: x,$$

that is, the union (3) equals $[1, A_{d,f} + B_{d,f} - 1] \setminus \{x\}$. Note that

$$A_{d,f} + B_{d,f} \geq a_d + a_f + b_d + b_f \geq 2 + 2B.$$

Hence $[1, A_{d,f} + B_{d,f} - 1]$ contains $(B, 2B + 1]$. By Lemma 2.9, there is a prime $p > B$ which is a hook length of λ . Now Lemma 2.2 implies that $S_{\mathbb{Z}}^\lambda$ is not globally irreducible. This accomplishes the induction step. \square

Lemma 3.2. *Suppose $S_{\mathbb{Z}}^\lambda$ is globally irreducible. Then either $n = 1$ or $B \geq 2$.*

Proof. Suppose $B = 1$ and $n \geq 2$. Then λ is the “staircase partition”, and $k \geq 2$. Hence 3 is a hook length for λ . Now, by Lemma 2.2, S_F^λ is reducible in characteristic 3. \square

In view of Corollary 3.2 we may (and will) assume that our globally irreducible Specht module $S_{\mathbb{Z}}^\lambda$ satisfies the condition $B \geq \max\{2, A\}$. Let q denote the maximal prime in the interval $[2, B]$.

Lemma 3.3. *Assume that $S_{\mathbb{Z}}^\lambda$ is globally irreducible, and $B \geq \max\{2, A\}$. If $f = k$, then $\lambda = (n)$.*

Proof. By Lemma 3.1, the hook lengths of λ cover the interval $[1, A_{1,k} + B_{1,k} - 1]$. We claim that $a_i < q$ for all $i \leq k$. Indeed, suppose $a_i \geq q$ for some i . Then

$$A_{1,k} + B_{1,k} - 1 \geq b_k + a_i - 1 \geq 2q - 1.$$

So the hook lengths of λ cover $(q, 2q)$. By Lemma 2.10, there is a prime $p > q$ such that p is a hook length of λ . Now Lemma 2.2 leads us to a contradiction (it applies as $p > B$). Thus $a_i < q$ for all i , that is, λ is q -regular. But then λ should satisfy Carter’s condition (see Proposition 2.1).

Observe that

$$h(A_{1,k}, b_k - (q - 1)) = q.$$

Therefore, $a_k = 1$ since otherwise

$$h(A_{1,k} - 1, b_k - (q - 1)) = q + 1$$

and λ does not satisfy Carter’s condition. If $k = 1$, this implies that $\lambda = (n)$. So suppose $k > 1$. Then

$$h(A_{1,k} - 1, b_k - (q - 1)) = q + 1 + b_{k-1}.$$

Now Carter’s condition forces $b_{k-1} \equiv -1 \pmod{q}$. In particular, $b_{k-1} \geq q - 1$. This gives $A_{1,k} + B_{1,k} - 1 \geq 2 + b_k + b_{k-1} - 1 \geq 2q$, so that $[1, A_{1,k} + B_{1,k} - 1] \supseteq (q, 2q)$. Again Lemmas 2.10 and 2.2 lead us to a contradiction completing the proof. \square

Now we are in a position to finish the proof of Theorem B.

Theorem 3.4. *Suppose $S_{\mathbb{Z}}^{\lambda}$ is globally irreducible and $B \geq \max\{2, A\}$. Then $\lambda = (n)$.*

Proof. By Lemma 3.3, we may assume that $f < k$. Applying Lemma 2.8(i) with $r = f, s = f + 1$ shows that

$$(4) \quad a_f + a_{f+1} \geq (p - 1)b_f + p$$

for any prime p dividing $a_f + a_{f+1} + b_f$.

If an odd prime divides $a_f + a_{f+1} + b_f$, then (4) implies that $a_f + a_{f+1} \geq 2b_f + 3$. In this case, either $a_f > b_f$ or $a_{f+1} > b_f$ contrary to the choice of b_f . Thus we must have

$$(5) \quad a_f + a_{f+1} + b_f = 2^m.$$

Combining this with (4) we get

$$(6) \quad a_f + a_{f+1} \geq b_f + 2.$$

Assume $f > 1$. Applying Lemma 2.8(i) with $r = f - 1, s = f + 1$ gives that

$$(7) \quad a_{f-1} + a_f + a_{f+1} \geq (p - 1)(b_{f-1} + b_f) + p$$

for any prime p dividing $a_{f-1} + a_f + a_{f+1} + b_f + b_{f-1}$. If an odd prime divides this number, then (7) yields

$$(8) \quad a_{f-1} + a_f + a_{f+1} \geq 2b_{f-1} + 2b_f + 3.$$

By Lemma 3.1, the hook lengths of λ cover $[1, a_f + a_{f-1} + b_f + b_{f-1} - 1]$. If

$$a_f + a_{f-1} + b_f + b_{f-1} - 1 \geq 2b_f - 1,$$

then $[1, a_f + a_{f-1} + b_f + b_{f-1} - 1] \supseteq (q, 2q)$. So again Lemma 2.10 ensures that there exists a prime $p_1 > q$ which is a hook length of λ , and again Lemma 2.2 shows that this contradicts the global irreducibility of $S_{\mathbb{Z}}^{\lambda}$. Thus

$$a_f + a_{f-1} + b_f + b_{f-1} - 1 < 2b_f - 1$$

which is equivalent to

$$2b_f - b_{f-1} > a_f + a_{f-1} + b_f.$$

However, this violates (8) as $b_f \geq a_{f+1}$. So we conclude that

$$(9) \quad a_{f-1} + a_f + a_{f+1} + b_f + b_{f-1} = 2^M$$

for some $M > m$. From (5) and (9) we have

$$a_{f-1} + b_{f-1} = 2^M - 2^m = 2^m(2^{M-m} - 1) \geq 2^m.$$

Therefore, either $a_{f-1} \geq 2^{m-1}$ or $b_{f-1} \geq 2^{m-1}$. Now (6) says that

$$b_f < \frac{1}{2}(a_f + a_{f+1} + b_f) = 2^{m-1}.$$

So $b_f < a_{f-1}$ or $b_f < b_{f-1}$ contrary to the choice of f .

Thus we may assume that $f = 1$. If $b_2 \geq 2$, then applying Lemma 2.8(ii) gives that there exists an odd prime p for which

$$(10) \quad a_1 + a_2 \geq (p - 1)(b_1 + 2) + 1$$

(indeed, $a_1 + a_2 + b_1 + 1$ is odd by (5)). As $p \geq 3$, (10) implies that $a_1 + a_2 \geq 2b_1 + 5$, whence either $a_1 > b_1$ or $a_2 > b_1$. Since this is impossible, we must have $b_2 = 1$.

Applying Lemma 2.8(i) to λ' instead of λ we obtain that

$$1 + b_1 \geq (p - 1)a_2 + p$$

for any prime p dividing $a_2 + b_1 + 1$. If there is an odd prime with this property, then $1 + b_1 \geq 2a_2 + 3$. Combining this with (6) (which is $a_1 + a_2 \geq b_1 + 2$ as $f = 1$) we get $a_1 \geq a_2 + 4$. Therefore, $a_1 + a_1 - 4 \geq a_1 + a_2 \geq b_1 + 2$ yielding

$$a_1 \geq \frac{b_1}{2} + 3.$$

So $[1, a_1 + b_1 - 1] \supset (b_1, \frac{3}{2}b_1 + 2]$. Due to Lemmas 2.11 and 3.1 one of the hooks of the Young diagram of λ has length equal to a prime greater than q . By Lemma 2.2, this is impossible. Thus $a_2 + b_1 + 1 = 2^s$. In view of (5) we also have $a_1 + a_2 + b_1 = 2^m$. So

$$a_1 - 1 = 2^s(2^{m-s} - 1),$$

and $m \geq s$. If $m > s$, we get $b_1 < 2^s \leq a_1 - 1$ which is false as $1 = f$. If $m = s$, we conclude that $a_1 = 1$. But then $a_1 + a_2 \geq b_1 + 2$ forces $a_2 > b_1$ leading us to a contradiction with the choice of b_f . This contradiction proves the theorem. \square

REFERENCES

- [1] R. W. Carter and M. T. J. Payne, On homomorphisms between Weyl modules and Specht modules, *Math. Proc. Camb. Phil. Soc.* **87** (1980), 419–425. MR **81h**:20048
- [2] C. W. Curtis and I. Reiner, *Representation theory of finite groups and associative algebras*, John Wiley and Sons (Interscience), New York, 1962. MR **26**:2519
- [3] W. and F. Ellison, *Prime numbers*, John Wiley and Sons, Hermann, 1985. MR **87a**:11082
- [4] G. D. James, On the decomposition matrices of the symmetric groups, II, *J. Algebra* **43** (1976), 45–54. MR **55**:3057b
- [5] G. D. James, On a conjecture of Carter concerning irreducible Specht modules. *Math. Proc. Camb. Phil. Soc.* **83** (1978), 11–17. MR **57**:3234
- [6] G. D. James, *The representation theory of the symmetric groups*, Springer Lecture Notes 682, Berlin, Heidelberg, New York, 1978. MR **80g**:20019
- [7] G. D. James and G. E. Murphy, The determinant of the Gram matrix for a Specht module. *J. Algebra* **59** (1979), 222–235. MR **82j**:20025
- [8] P. Ribbenboim, *The book of prime number records*, Springer–Verlag, New York, Berlin, Heidelberg, 1988.
- [9] J. G. Thompson, Finite groups and even lattices. *J. Algebra* **38** (1976), 523–524. MR **53**:3108
- [10] J. G. Thompson, A simple subgroup of $E_8(3)$. In: N. Iwahori (ed.), *Finite groups symposium*, Japan Soc. for promotion of Science, 1976, pp. 113–116.
- [11] Pham Huu Tiep, Globally irreducible representations of finite groups and integral lattices, *Geometriae Dedicata* **64** (1997), 85–123. MR **98e**:20011

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OREGON, EUGENE, OREGON 97403
 E-mail address: klesh@math.uoregon.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MANCHESTER, OXFORD ROAD, MANCHESTER, M13 9PL, UNITED KINGDOM
 E-mail address: sashap@ma.man.ac.uk