

ON CONDENSATIONS OF C_p -SPACES ONTO COMPACTA

A. V. ARHANGEL'SKII

(Communicated by Alan Dow)

ABSTRACT. A condensation is a one-to-one onto mapping. It is established that, for each σ -compact metrizable space X , the space $C_p(X)$ of real-valued continuous functions on X in the topology of pointwise convergence condenses onto a metrizable compactum. Note that not every Tychonoff space condenses onto a compactum.

The space $C_p(X)$ of all real-valued continuous functions on a Tychonoff space X in the topology of pointwise convergence is σ -compact only in the trivial case when X is finite (see Theorem 2 below). This makes natural the following question: *when can $C_p(X)$ be condensed onto a compact space or onto a σ -compact space?* A condensation is any one-to-one onto continuous mapping. A compactum is a compact Hausdorff space. We consider only Tychonoff spaces (observe that *every T_1 -space can be condensed onto a compact T_1 -space*). S. Banach was probably the first to ask when a (separable metrizable) space can be condensed onto a compact space (see [4]). E.G. Pytkeev solved one of Banach's problems, proving that every separable Banach space can be condensed onto a (metrizable) compactum [7]. The question above appears as Problem 35 in [2], accompanied by several versions of it. In particular, Problem 39 in [2] runs as follows: *Is it possible to condense $C_p(D^\omega)$ onto a compact space* (where D is the discrete two-point space, and D^ω is the Cantor set)? We answer this question below. The main result is:

1. Theorem. *For any σ -compact metrizable space X , the space $C_p(X)$ condenses onto a metrizable compactum.*

To prove Theorem 1, we need a result from [1]. If X is a space and Y is a subspace of X , then $C_p(Y, X)$ is the subspace of $C_p(Y)$ consisting of restrictions to Y of continuous real-valued functions on X . The next theorem was established in [1] (see Theorem 1.2.2):

2. Theorem. *If Y is dense in X , and $C_p(Y, X)$ is σ -countably compact, then X is pseudocompact, and Y is a P -space.*

Recall that a P -space is a space in which every G_δ -subset is open, and that a space is σ -countably compact, if it is the union of a countable family of countably compact subspaces.

Received by the editors May 24, 1997 and, in revised form, May 15, 1998.

2000 *Mathematics Subject Classification.* Primary 54A10, 54C35, 54C10.

Key words and phrases. Condensation, compactum, network, topology of pointwise convergence, σ -compact space, Borel space, P -space.

Proof of Theorem 1. It suffices to establish that $C_p(X)$ condenses onto some compactum. Indeed, any compactum with a countable network is metrizable [3], and any network for $C_p(X)$ is a network for any coarser topology. Since X has a countable base, $C_p(X)$ has a countable network [1].

If X is discrete, then $C_p(X)$ is just R^X . Clearly, the space R condenses onto a compactum, since, obviously, every locally compact space condenses onto a compactum [4]. Therefore, R^X condenses onto a compactum.

It remains to consider the case when X is not discrete. We can fix a countable subspace Y dense in X such that $C_p(Y, X)$ is not σ -compact. Indeed, assume the contrary, and take any countable Y dense in X . Then, by Theorem 2, Y is a P -space. Since Y is countable, it follows that Y is discrete. Therefore, X is discrete, since X is separable and every countable dense subspace of X is discrete, a contradiction.

On the other hand, from Theorem 6.2 in [6] it follows that $C_p(Y, X)$ is a Borel subset of the separable complete metric space R^Y (it is here that we use σ -compactness of X). Now, E.G. Pytkeev established in [7] that every non- σ -compact separable metrizable Borel space condenses onto a compactum. (Notice that the space Q of rational numbers does not condense onto any compactum, since each non-empty countable compactum has an isolated point.)

Therefore, $C_p(Y, X)$ condenses onto a compactum K . Finally, since the natural restriction mapping of functions on X to Y is a condensation of $C_p(X)$ onto $C_p(Y, X)$, we conclude that $C_p(X)$ condenses onto the compactum K . \square

3. Remarks. A result similar to Theorem 1 holds for the space $C_p^b(X)$ of bounded continuous real-valued functions on X in the topology of pointwise convergence: this space also condenses onto a compactum whenever X is a σ -compact metrizable space. A minor change is needed in the proof: we should refer to Proposition 9.2 in [3]. Theorem 1 also implies that, under the same restrictions on X as in Theorem 1, $C(X)$ condenses onto a compactum when $C(X)$ is endowed with any stronger topology than the topology of pointwise convergence (for example, with the compact-open topology).

4. Problem. Is it true that $C_p(X)$ condenses onto a compactum (onto a σ -compact space) for every separable metrizable space X ?

I conjecture that it might be impossible to condense $C_p(J)$ onto any compactum, where J is the space of irrational numbers.

5. Problem. Is it true that $C_p(X)$ condenses onto a σ -compact space for every compact space X ?

REFERENCES

- [1] A.V. Arhangel'skii, *Topological Function Spaces* (Kluwer Academic Publishers, Dordrecht, 1992), 205. MR **92i**:54022
- [2] A.V. Arhangel'skii, *C_p-theory*, p. 1-56 in: M. Hušek and J. van Mill, Eds., *Recent Progress in General Topology*, North-Holland, Amsterdam-London-New-York, 1992, 796. MR **95g**:54004
- [3] A.V. Arhangel'skii, *Function spaces in the topology of pointwise convergence and compact sets*, *Russian Math. Surveys* 39:5 (1984), 9-56.
- [4] A.V. Arhangel'skii and V.I. Ponomarev, *Fundamentals of General Topology: Problems and Exercises*, Reidel, 1984. MR **87i**:54001
- [5] S. Banach, *Livre Ecossais*, Problem 1, 17:8, 1935; *Colloq. Math.* 1 (1947), p. 150.

- [6] T. Dobrowolski and W. Marciszewski, *Classification of function spaces with the topology determined by a countable dense set*, *Fundamenta Mathematicae* 148 (1995), 35-62. MR **96k**:54017
- [7] E.G. Pytkeev, *Upper bounds of Topologies*, *Matem. Notes* 20:4 (1976), 831-837. MR **55**:1262

DEPARTMENT OF MATHEMATICS, 321 MORTON HALL, OHIO UNIVERSITY, ATHENS, OHIO 45701
E-mail address, January 1–June 15: arhangel@bing.math.ohiou.edu

CHAIR OF GENERAL TOPOLOGY AND GEOMETRY, MECH.-MATH. FACULTY, MOSCOW STATE
UNIVERSITY, MOSCOW 119899, RUSSIA
E-mail address, June 15–December 31: arhala@arhala.mccme.ru