

ON NON-ORIENTABLE SURFACES IN 4-SPACE WHICH ARE PROJECTED WITH AT MOST ONE TRIPLE POINT

SHIN SATOH

(Communicated by Ronald A. Fintushel)

ABSTRACT. We show that if a non-orientable surface embedded in 4-space has a projection into 3-space with at most one triple point, then it is ambient isotopic to a connected sum of some unknotted projective planes and an embedded surface in 4-space with vanishing normal Euler number.

1. INTRODUCTION

In [4] Kinoshita constructed infinitely many projective planes embedded in \mathbf{R}^4 which are not ambient isotopic to each other. Each of his projective planes is a connected sum of an unknotted projective plane and a 2-knot. Price and Roseman gave a method to construct a projective plane in \mathbf{R}^4 from an invertible 1-knot and proved that every spun projective plane is ambient isotopic to a connected sum of an unknotted projective plane and a 2-twist spun 2-knot [6]. It is unknown whether there is an example of a projective plane in \mathbf{R}^4 which is not a connected sum of an unknotted projective plane and a 2-knot.

On the other hand, Yoshikawa gave infinitely many Klein bottles in \mathbf{R}^4 each of which is not ambient isotopic to a connected sum of an unknotted projective plane and a (possibly knotted) projective plane [7].

In this paper we discuss the problem of when a non-orientable surface in \mathbf{R}^4 is ambient isotopic to a connected sum of unknotted projective planes and a surface in \mathbf{R}^4 from the viewpoint of *broken surface diagrams* due to Carter and Saito [2].

A *surface in \mathbf{R}^4* means a closed and connected PL 2-manifold without boundary embedded in \mathbf{R}^4 locally flatly. We call a surface in \mathbf{R}^4 a *2-knot* if the embedded surface is homeomorphic to a 2-sphere. Moreover we can assume that the image $p(F)$ under the projection $p : \mathbf{R}^4 \rightarrow \mathbf{R}^3$ is a *generic* surface; that is, the singular set $\Gamma(p(F))$ on $p(F)$ consists of (possibly empty) isolated branch points, double point curves, and isolated triple points. Then we prove the following theorem.

Theorem 1.1. *Let F be a non-orientable surface in \mathbf{R}^4 and $e(F)$ its normal Euler number. If $\Gamma(p(F))$ contains at most one triple point, then F is ambient isotopic to a connected sum of $|e(F)|/2$ unknotted projective planes and a surface F' in \mathbf{R}^4 with $e(F') = 0$.*

Received by the editors July 20, 1998 and, in revised form, October 7, 1998.

1991 *Mathematics Subject Classification.* Primary 57Q45.

Key words and phrases. Non-orientable surface, connected sum, projective plane, triple point, branch point, normal Euler number.

We notice that for Yoshikawa’s Klein bottle Kb , $\Gamma(p(Kb))$ contains no triple point and $e(Kb)$ is equal to zero.

Let χ be the Euler characteristic of a non-orientable surface F in \mathbf{R}^4 . Then the normal Euler number $e(F)$ is equal to $2\chi - 4, 2\chi, \dots, -2\chi$ or $4 - 2\chi$ (cf. [5]). Since the normal Euler number of every projective plane is ± 2 , we have the following.

Corollary 1.2. *Let P be a projective plane in \mathbf{R}^4 . If $\Gamma(p(P))$ contains at most one triple point, then P is ambient isotopic to a connected sum of an unknotted projective plane and a 2-knot.*

2. PROJECTION WITH NO TRIPLE POINT

Let F be a surface in \mathbf{R}^4 and $p(F)$ its generic projection, where $p : \mathbf{R}^4 \rightarrow \mathbf{R}^3$ is the projection map.

The regular neighborhood in \mathbf{R}^3 of a branch point on $p(F)$ is as depicted in Figure 1(A). We define signs of branch points in such a way that a branch point with the crossing information depicted in Figure 1(B) receives the sign $+1$, and one with the crossing information in Figure 1(C) receives the sign -1 . Then the sum of signs taken over all the branch points in $\Gamma(p(F))$ is equal to the normal Euler number $e(F)$ of F (cf. [1]).




FIGURE 1.

A projective plane P in \mathbf{R}^4 is *unknotted* if it is ambient isotopic to the surface whose projection into \mathbf{R}^3 is as depicted in Figure 2(A). There are two distinct unknotted projective planes, one of which has the normal Euler number $e(P) = +2$ and the other has $e(P) = -2$; see Figure 2(B) and (C). We denote these P by P_+ (positive type) and P_- (negative type) respectively.




FIGURE 2.

Let C_1, \dots, C_n be the connected components of the singular set $\Gamma(p(F))$. Each C_i is regarded as a union of immersed loops and immersed arcs in \mathbf{R}^3 such that the endpoints of the immersed arcs are branch points. Suppose that C_i is one simple arc. Such a simple arc is called an m -arc (resp. a -arc) if the two branch points of

its ends have the same sign (resp. opposite signs). Notice that the neighborhood of an m -arc (resp. a -arc) is homeomorphic to a Möbius band (resp. an annulus). Call an m -arc *positive* if it terminates with both positive type, or *negative* if it terminates with negative type. These terms are used in [3].

Lemma 2.1. *Let F be a non-orientable surface in \mathbf{R}^4 . If $\Gamma(p(F))$ contains no triple point, then F is ambient isotopic to a connected sum of $|e(F)|/2$ unknotted projective planes and a surface F' in \mathbf{R}^4 with $e(F') = 0$.*

Proof. It is sufficient to prove the lemma in the case that $e(F) > 0$. Since $\Gamma(P(F))$ contains no triple point, $\Gamma(P(F))$ is a disjoint union of simple loops, a -arcs and m -arcs. Notice that $e(F) = 2(p - n)$, where p is the number of positive m -arcs and n is the number of negative m -arcs in $\Gamma(p(F))$. Then $\Gamma(p(F))$ contains at least $e(F)/2$ positive m -arcs, and so F is ambient isotopic to a connected sum of $e(F)/2$ P_+ and a surface F' in \mathbf{R}^4 . Since $\Gamma(p(F'))$ is obtained from $\Gamma(p(F))$ by deleting the $e(F)/2$ positive m -arcs, we have $e(F') = 0$. □

3. PROJECTION WITH ONE TRIPLE POINT

Let F be a surface in \mathbf{R}^4 and $p(F)$ its generic projection where $p : \mathbf{R}^4 \rightarrow \mathbf{R}^3$ is a projection map. The neighborhood of a triple point on $p(F)$ consists of three sheets. These sheets can be labeled *top*, *middle* and *bottom*, and these indicate the relative position of the sheets with respect to the projection direction.

The following is the key lemma in this paper.

Lemma 3.1. (1) *Let S be a generic closed surface in \mathbf{R}^3 and c a simple closed curve in \mathbf{R}^3 such that c is transverse to S with $c \cap \Gamma(S) = \emptyset$. Then the number of $c \cap S$ is even.*

(2) *Let t be a triple point in $\Gamma(p(F))$ and H one of the three sheets around t . Suppose that there exists a subarc $\alpha : [0, 1] \rightarrow \Gamma(p(F))$ such that $\alpha(0) = \alpha(1) = t$, such that $\alpha([0, \varepsilon]) \subset H$, $\alpha([1-\varepsilon, 1]) \subset H$ for a small $\varepsilon > 0$, and such that $\alpha(\text{int}[0, 1])$ contains no triple point. Then H is a top sheet or a bottom sheet.*

Proof. (1) Since the \mathbf{Z}_2 -intersection number of c and S in \mathbf{R}^3 is zero, the number of $c \cap S$ is even.

(2) There are two cases as depicted in Figure 3, where the shaded sheet represents the sheet H . Suppose that H is a middle sheet in each case. Then its broken surface diagram is as depicted in Figure 4. Therefore the pasting map in the figure must be a rotation by 0° or 180° in Figure 4(A) and $\pm 90^\circ$ in Figure 4(B). This contradicts (1). □



FIGURE 3.




FIGURE 4.

We notice that in Lemma 3.1(2), even if H is a top sheet or a bottom sheet, the case in Figure 4(A) does not occur.

The following lemma is proved by performing a sequence of two Roseman moves — “a sheet moving through a saddle point” and “passing a branch point through a third sheet”, where the *Roseman moves* are generalizations of the Reidemeister moves and are sufficient to move embedded surfaces in \mathbf{R}^4 . The list of the Roseman moves appears in [2].

Lemma 3.2. *Let t be a triple point in $\Gamma(p(F))$ and H_i ($i = 1, 2, 3$) three sheets around t . Suppose that the small arc $H_2 \cap H_3$ is extended to an immersed arc γ in $\Gamma(p(F))$ which ends in two branch points b and b' and that the subarc γ' of γ connecting between t and b contains no triple point inside. If H_1 is a top sheet or a bottom sheet, then the triple point t can be removed from $p(F)$ by deforming F by an ambient isotopy of \mathbf{R}^4 without introducing new triple points.*

Proof. Let D be a small neighborhood of t in H_1 . Push D along γ' towards b . Since H_1 is a top sheet or a bottom sheet, we can push out D over b . This deformation can be realized by an ambient isotopy of \mathbf{R}^4 and does not introduce new triple points in the projection. See Figure 5 (the figure is in the case that H_1 is a top sheet). \square




FIGURE 5.

Proof of Theorem 1.1. By Lemma 2.1, it is sufficient to consider the case that $\Gamma(p(F))$ contains just one triple point t . Let C_1, \dots, C_n be the connected components of $\Gamma(p(F))$ with $t \in C_1$.

If C_1 is a union of immersed loops, then all the branch points in $\Gamma(p(F))$ belong to $C_2 \cup \dots \cup C_n$, where each of C_2, \dots, C_n is a simple loop or a simple arc. Then we can apply an argument similar to the argument in the proof of Lemma 2.1.

Suppose that C_1 contains at least one immersed arc, say γ . We will prove that t can be removed from C_1 without introducing new triple points. Let $c : [0, 1] \rightarrow \mathbf{R}^3$




FIGURE 6.

be an immersion representing γ . First consider the case that $c^{-1}(t)$ consists of two or three points; see Figure 6(B) and (C). Then we can perform the deformation in Lemma 3.2 so that t can be removed from C_1 . Consider the case that $c^{-1}(t)$ is one; see Figure 6(A). If the perpendicular sheet to γ is top or bottom, then we can also perform the deformation in Lemma 3.2. If the perpendicular sheet is middle, then C_1 has the only form which consists of three arcs intersecting at t by Lemma 3.1. Therefore we can perform the deformation in Lemma 3.2. \square

REFERENCES

- [1] Carter, J. Scott, and Saito, Masahico, *Canceling branch points on projections of surfaces in 4-space*, Proc. of the AMS. **116**, No 1 (Sept 1992), pp. 229-237. MR **93i**:57029
- [2] ———, *Reidemeister moves for surface isotopies and their interpretation as moves to movies*, J. of Knot Theory and its Ramifications **2** (1993), pp. 251-284. MR **94i**:57007
- [3] ———, *Normal Euler classes of knotted surfaces and triple points on projections*, Proc. of the AMS. **125**, No 2 (Feb 1997), pp. 617-623. MR **97d**:57030
- [4] S. Kinoshita, *On the Alexander polynomials of 2-spheres in a 4-sphere*, Ann. of Math. **74**, No 3 (Nov 1961), pp. 518-531. MR **24**:A2960
- [5] W. S. Massey, *Proof of a conjecture of Whitney*, Pacific J. Math. **31** (1969), pp. 143-156. MR **40**:3570
- [6] T. M. Price and D. Roseman, *Embeddings of the projective plane in four space*, preprint.
- [7] K. Yoshikawa, *The order of a meridian of a knotted Klein bottle*, preprint.

DEPARTMENT OF MATHEMATICS, OSAKA CITY UNIVERSITY, SUGIMOTO, SUMIYOSHI-KU, OSAKA, 558-5858, JAPAN

E-mail address: susato@sci.osaka-cu.ac.jp