

ON THE REGULARITY OF p -BOREL IDEALS

JÜRGEN HERZOG AND DORIN POPESCU

(Communicated by Wolmer V. Vasconcelos)

ABSTRACT. In this paper we prove Pardue's conjecture on the regularity of principal p -Borel ideals. As a consequence we obtain an upper bound for the regularity of general p -Borel ideals.

INTRODUCTION

Let K be field and $I \subset S$ a graded ideal in the polynomial ring $S = K[x_1, \dots, x_n]$. Recall that the generic initial ideal $\text{Gin}(I)$ of I with respect to the reverse lexicographical order is Borel-fixed. This means that $\text{Gin}(I)$ is fixed under the action of the Borel group of the upper triangular invertible matrices acting linearly on the polynomial ring. By a theorem of Bayer and Stillman [2], the regularity of I and $\text{Gin}(I)$ coincide. This is one of the reasons why one is interested in computing the regularity of $\text{Gin}(I)$. In characteristic zero a Borel-fixed ideal is strongly stable, and so its regularity is simply the highest degree of a minimal generator. In positive characteristic however, Borel-fixed ideals are p -Borel (see 1.1 for the definition), and these are monomial ideals with a quite difficult combinatorial structure.

Monomials $u_1, \dots, u_m \in I$ of a p -Borel are called Borel generators of I if I is the smallest p -Borel ideal containing u_1, \dots, u_m . In this case we write $I = \langle u_1, \dots, u_m \rangle$. The ideal I is called principal p -Borel if I has only one Borel generator. Pardue conjectured a formula for the regularity of a principal p -Borel ideal which only depends on the exponents of the Borel generator (see 1.4). In a paper by Aramova and Herzog [1] it was shown that Pardue's formula indeed gives a lower bound for the regularity. Some of the results in that paper were later extended by Ene, Pfister and Popescu [5] to more general ideals. In the present paper we will show that Pardue's formula also yields an upper bound. Our method in proving this uses a criterion of Eisenbud, Reeves and Totaro [4] for determining the regularity of p -Borel ideals.

The authors are grateful to the Alexander von Humboldt Foundation for supporting their collaboration.

1. p -BOREL IDEALS

Throughout this paper we fix a field K and let $S = K[x_1, \dots, x_n]$ be the polynomial in n indeterminates over K .

Received by the editors October 12, 1999 and, in revised form, January 24, 2000.
1991 *Mathematics Subject Classification*. Primary 13P10; Secondary 13D02, 13C13.
Key words and phrases. Regularity, p -Borel ideals.

Let p be a prime number, and k and l be non-negative integers with p -adic expansion $k = \sum_i k_i p^i$ and $l = \sum_i l_i p^i$. We set $k \leq_p l$ if $k_i \leq l_i$ for all i .

Definition 1.1. A monomial ideal $I \subset S$ is p -Borel if the following condition holds: for each monomial $u \in I$, $u = \prod_i x_i^{\mu_i}$, one has $(x_i/x_j)^\nu u \in I$ for all i, j with $1 \leq i < j \leq n$ and all $\nu \leq_p \mu_j$.

The significance of p -Borel principal ideals is given by

Proposition 1.2 (Pardue). *Suppose $\text{char } K = p$, and let $I \subset S$ be a monomial ideal. Then I is Borel-fixed if and only if I is p -Borel.*

We denote by $G(I)$ the unique minimal set of monomial generators of a monomial ideal I . It is easy to see (cf. [1]) that I is p -Borel if the conditions of 1.1 are satisfied for all $u \in G(I)$.

A principal p -Borel ideal can be explicitly described. We use the following standard notation: If J is a monomial ideal we let $J^{[p^i]}$ be the ideal generated by all monomials u^{p^i} with $u \in G(J)$. The ideal $J^{[p^i]}$ is called the p^i th Frobenius power of J . Note that we define the p^i th Frobenius power of monomial ideals in any characteristic.

Proposition 1.3 (Pardue). *Let $u = \prod_i x_i^{\mu_i}$, and let $\mu_i = \sum_j \mu_{ij} p^j$ be the p -adic expansion of μ_i for $i = 1, \dots, n$. Then*

$$\langle u \rangle = \prod_{i=1}^n \prod_j ((x_1, \dots, x_i)^{\mu_{ij}})^{[p^j]}.$$

In particular, $\langle u \rangle = \prod_{i=1}^n \langle x_i^{\mu_i} \rangle$.

It follows from 1.3 that $\langle x^\mu \rangle = (x_1^{\mu_1}) \langle x^\mu / x_1^{\mu_1} \rangle$, so that $\text{reg} \langle x^\mu \rangle = \mu_1 + \text{reg} \langle x^\mu / x_1^{\mu_1} \rangle$. Therefore, if we are interested in the regularity of the p -Borel principal ideal $\langle x^\mu \rangle$ we may assume that x_1 does not divide x^μ .

Denote by $[*]$ the greatest integer function, and for $1 \leq k \leq n$ and $j \geq 0$ define

$$d_{kj}(\mu) = \sum_{i=1}^k \lfloor \frac{\mu_i}{p^j} \rfloor.$$

For each k such that $\mu_k \neq 0$, let $s_k = \lfloor \log_p \mu_k \rfloor$, and set

$$D_k = d_{ks_k}(\mu) p^{s_k} + (k - 1)(p^{s_k} - 1).$$

Conjecture 1.4 (Pardue). *If x_1 does not divide x^μ , then*

$$\text{reg} \langle x^\mu \rangle = \max_{k: \mu_k \neq 0} \{D_k\}.$$

In the following we will express the right-hand side of Conjecture 1.4 in different ways. The following easy fact can be found in [1].

Proposition 1.5. *Let $\mathcal{S} = \{s_k: \mu_k \neq 0\}$, let $q_j = \max\{k: s_k = j\}$ for each $j \in \mathcal{S}$, and set $E_j = D_{q_j}$. Then:*

- (i) $E_j = \sum_{i=j}^s (\sum_{k=2}^{q_j} \mu_{ki}) p^i + (q_j - 1)(p^j - 1)$ for all $j \in \mathcal{S}$;
- (ii) $\max\{D_k: \mu_k \neq 0\} = \max\{E_j: j \in \mathcal{S}\}$.

We shall need still another reformulation of Pardue’s formula for the regularity of a principal p -Borel ideal. Set $s = \max\{s_k : \mu_k \neq 0\}$, and for each t with $1 \leq t \leq s$ let $m_t = \max\{k : \mu_{kt} \neq 0\}$. Finally set

$$F_t = \sum_{i=t}^s \left(\sum_{k=2}^n \mu_{ki} \right) p^i + (m_t - 1)(p^t - 1) \quad \text{for all } t = 1, \dots, s.$$

Proposition 1.6. *With the notation introduced we have*

$$\max_{1 \leq t \leq s} F_t = \max_{j \in \mathcal{S}} E_j.$$

Proof. It is clear that $m_j \geq q_j$ for all $j \in \mathcal{S}$, so that $\max_{1 \leq t \leq s} F_t \geq \max_{j \in \mathcal{S}} E_j$.

In order to show the opposite inequality we first prove the following claim: Let $\mathcal{S}_t = \{j \in \mathcal{S} : j \geq t\}$ and $Q_t = \{q_j : j \in \mathcal{S}_t\}$. (Note that $\mathcal{S}_t \neq \emptyset$, since $s \in \mathcal{S}_t$.) Let $e \in \mathcal{S}_t$ such that $q_e = \max\{q_j \in Q_t\}$. Then we claim that $F_e \geq F_t$.

Indeed, we have

$$F_e - F_t = - \sum_{i=t}^{e-1} \left(\sum_{k=2}^n \mu_{ki} \right) p^i + (m_e - 1)(p^e - 1) - (m_t - 1)(p^t - 1).$$

Since we assume that q_e is maximal in Q_t it follows that $\mu_{ki} = 0$ for $k > q_e$ and $i \geq t$. Thus $m_e = q_e \geq \max\{k : \mu_{kt} \neq 0\} = m_t$, because again $\mu_{kt} = 0$ for $k > q_e$.

Now it follows that

$$F_e - F_t = - \sum_{i=t}^{e-1} \left(\sum_{k=2}^{q_e} \mu_{ki} \right) p^i + (q_e - 1)(p^e - 1) - (m_t - 1)(p^t - 1).$$

Finally, since $\sum_{i=t}^{e-1} \left(\sum_{k=2}^{q_e} \mu_{ki} \right) p^i = \sum_{k=2}^{q_e} \left(\sum_{i=t}^{e-1} \mu_{ki} p^i \right) \leq (q_e - 1)(p^e - p^t)$, we get $F_e - F_t \geq (q_e - 1)(p^t - 1) - (m_t - 1)(p^t - 1) = (q_e - m_t)(p^t - 1) \geq 0$. This concludes the proof of the claim.

Continuing with the proof of the opposite inequality, we let $t \leq s$ be the maximal number for which $F_t = \max_{1 \leq r \leq s} F_r$. Let $e \in \mathcal{S}_t$ be chosen such that q_e is maximal in Q_t . Then, according to our claim, we have $F_e \geq F_t$. By the choice of t this implies that $e = t$, so that, in particular, $t \in \mathcal{S}_t$. Since q_t is maximal in Q_t it now follows that $\mu_{ki} = 0$ for $i \geq t$ and $k > q_t$. Consequently, $m_t = q_t$, and so $F_t = E_t$. □

Remark 1.7. Using the methods of [1] the following result was proved in [5]: Let $(I_t)_{1 \leq t \leq s}$ be some stable ideals and $I = \prod_{t=1}^s I_t^{[p^{r_t}]}$ for some integers $0 \leq r_1 < \dots < r_s$. If I_j contains $x_{m(I_{j+1})}^{p^{r_{j+1}-r_j}-1}$ for all $1 \leq j < s$ (we set $m(u) = \max\{j : x_j | u\}$ for a monomial u) and $m(I_{j+1}) = \max\{m(u) : u \in G(I_{j+1})\}$, then $\text{reg}(I) = \text{pa}(I)$, where

$$\text{pa}(I) = \max_{1 \leq t \leq s} \left\{ \sum_{i>t}^s p^{r_i} \max(I_i) + \max_{u \in G(I_t)} [p^{r_t} \deg(u) + (m(u) - 1)(p^{r_t} - 1)] \right\}.$$

Moreover if I_t has the form $I_t = \prod_{i=2}^n (x_1, \dots, x_i)^{\mu_{it}}$ with $0 \leq \mu_{it} < p$ for all $t < s$, the above result gives $\text{reg}(I) = \max_{1 \leq t \leq s} F_t$. Hence the Pardue Conjecture holds in a special case, which can also be obtained directly from [1]. Trying to extend the equality $\text{reg}(I) = \text{pa}(I)$ for general products of p^i th Frobenius powers of stable ideals one must consider first the following example which shows how tight Pardue’s Conjecture is: Let $n = 3$, $p = 2$, $I_1 = (x_1, x_2)^2$, $I_2 = (x_1, x_2, x_3)$ and $I = I_1 I_2^{[2]}$. Then $\text{pa}(I) = 4$, but $\text{reg}(I) > 4$, because I is not stable (see Proposition 2.1 below).

2. THE PROOF OF PARDUE’S CONJECTURE

In [1] it is shown that if $x^\mu \in S$ is a monomial which is not divisible by x_1 , then

$$\text{reg}(x^\mu) \geq \max_{k: \mu_k \neq 0} \{D_k\}.$$

In this section we will prove the opposite inequality. Our proof is based on the following result [4]:

Proposition 2.1 (Eisenbud, Reeves, Totaro). *Let I be a p -Borel ideal with $\max(I) = d$, and let $e \geq d$ be the smallest integer such that $I_{\geq e}$ is stable. Then $\text{reg}(I) = e$.*

Proposition 2.1 needs some explanation: $\max(I) = \max\{\deg u : u \in G(I)\}$, and $I_{\geq e}$ is the ideal generated by all monomials $u \in I$ with $\deg u \geq e$. Finally, recall that, according to Eliahou and Kervaire [3], a monomial ideal I is stable if for all monomials (or equivalently all generators) u of I one has $(x_i/x_{m(u)})u \in I$ for all $i \leq m(u)$, where $m(u) = \max\{j : x_j|u\}$.

Recall from Section 1 that $\langle x^\mu \rangle = \prod_t I_t^{[\mu_{it}]}$ where $I_t = \prod_{i=2}^n (x_1, \dots, x_i)^{\mu_{it}}$ with $0 \leq \mu_{it} < p$. Thus the desired inequality will follow from Proposition 2.1 and

Theorem 2.2. *For given integers $0 \leq r_1 < \dots < r_s$, and integers $0 \leq a_{tk} < p^{r_{t+1}-r_t}$ for $t = 1, \dots, s$ and $k = 1, \dots, m_t$ let $I_t = \prod_{k=2}^{m_t} (x_1, \dots, x_k)^{a_{tk}}$ and $I = \prod_{t=1}^s I_t^{[p^{r_t}]}$. Let $\delta_t = \sum_{i=t}^s p^{r_i} \max(I_i) + (m_t - 1)(p^{r_t} - 1)$ and $d = \max\{\delta_t | 1 \leq t \leq s\}$. Then $I_{\geq d}$ is stable.*

The proof needs some preparation.

Lemma 2.3. *Let $J = \prod_{k=2}^m (x_1, \dots, x_k)^{a_k}$ with $0 \leq a_k \leq p^r - 1$, and let $\eta \in J$ be a monomial such that $\deg \eta \geq 1 + \max\{m(\eta)(p^r - 1), \max(J) + p^r - 1\}$. Then there exists an integer t such that $\eta \in x_t^{p^r} J$.*

Proof. We reduce the problem to the case where $m(\eta) \leq m$. Since J is stable, η has the following Eliahou-Kervaire decomposition: $\eta = vw$ for monomials v and w with $v \in G(J)$ and $\min(w) \geq m(v)$. We may assume that $w \notin (x_t^{p^r})$ for all t . Then $w = w'x_{m+1}^{\beta_{m+1}} \dots x_{m(\eta)}^{\beta_{m(\eta)}}$ with $\beta_i \leq p^r - 1$. Thus the element $\eta' = vw'$ has degree $\geq \deg \eta - (m(\eta) - m)(p^r - 1) \geq m(p^r - 1) + 1$. Since $m(\eta') \leq m$ and $\max(J) \leq (m - 1)(p^r - 1)$, we may replace η by η' , and thus may as well suppose that $m(\eta) \leq m$.

Let $\eta = x_1^{\alpha_1} \dots x_{m(\eta)}^{\alpha_{m(\eta)}}$. We apply induction on $\max(J)$, and may assume that $a_m \neq 0$. If $\max(J) = 1$, then $J = (x_1, \dots, x_m)$. If $m(\eta) = 1$, then $\eta = x_1^{\alpha_1}$ with $\alpha_1 \geq \max(J) + p^r = p^r + 1$. In that case, $\eta = x_1^{p^r} \eta'$ with $\eta' = x_1^{\alpha_1 - p^r} \in J$. Suppose now that $m(\eta) \geq 2$. Then, since $\deg \eta \geq m(\eta)(p^r - 1) + 1$, it follows that $\eta \in (x_t^{p^r})$ for a certain $t \leq m(\eta)$, and so $\eta = x_t^{p^r} \eta'$ where η' is a monomial of degree $\geq (m(\eta) - 1)(p^r - 1) > 1$. Hence $\eta' \in J$, and so $\eta \in x_t^{p^r} J$.

Now suppose that $\max(J) > 1$. We will distinguish several cases. In the first case suppose that $\alpha_{m(\eta)} \geq p^r$. Let again $\eta = vw$ be the Eliahou-Kervaire decomposition of η . Then $\deg w = \deg \eta - \deg v \geq p^r > 0$. Hence, since $m(v) \leq \min(w)$, it follows that $x_{m(\eta)}^{p^r}$ divides w , and we are done.

Now we consider the case that $\alpha_{m(\eta)} \leq p^r - 1$, $m(\eta) \geq 3$ and $\alpha_{m(\eta)} \leq \sum_{i=m(\eta)}^m a_i$. We choose the maximal integer t , $m(\eta) \leq t \leq m$, such that $\alpha_{m(\eta)} \leq \sum_{i=t}^m a_i$, and

write $\alpha_{m(\eta)} = \sum_{i=t+1}^m a_i + b_t$ with $1 \leq b_t \leq a_t$. Now set $\varphi = \eta/x_{m(\eta)}^{\alpha_{m(\eta)}}$. Observe that for all monomials $\rho \in J$ with $m(\rho) \leq m$ one has

$$(1) \quad \rho/x_{m(\eta)} \in \prod_{k=2}^{m-1} (x_1, \dots, x_k)^{a_k} (x_1, \dots, x_m)^{a_m-1}.$$

Applying (1) successively we see that $\varphi \in J''$ where

$$J'' = \prod_{k=2}^{t-1} (x_1, \dots, x_k)^{a_k} (x_1, \dots, x_t)^{a_t-b_t}.$$

We have

$$\deg \varphi = \deg \eta - \alpha_{m(\eta)} \geq \deg \eta - (p^r - 1) \geq 1 + (m(\eta) - 1)(p^r - 1) \geq 1 + m(\varphi)(p^r - 1),$$

and $\deg \varphi = \deg \eta - \alpha_{m(\eta)} \geq \max(J) + p^r - \alpha_{m(\varphi)} = \max(J'') + p^r$. Hence we may apply our induction hypothesis and conclude that there exists an integer $q \leq m(\varphi)$ such that $\varphi \in x_q^{p^r} J''$. It follows that $\eta \in x_q^{p^r} J$.

Next we consider the case $\alpha_{m(\eta)} \leq p^r - 1$, $m(\eta) \geq 3$, and $\alpha_{m(\eta)} > \sum_{i=m(\eta)}^m a_i$. Using (1) again we see that $\eta = x_{m(\eta)}^{\sum_{i=m(\eta)}^m a_i} \eta'$ with $\eta' \in \tilde{J}$ where

$$\tilde{J} = \prod_{k=2}^{m(\eta)-1} (x_1, \dots, x_k)^{a_k}.$$

Note that for any monomial $\rho \in J$ with $m(\rho) > m$ it follows that $\rho/x_{m(\eta)} \in J$. Applying this successively to η' we see that $\varphi = \eta/x_{m(\eta)}^{\alpha_{m(\eta)}}$ belongs to \tilde{J} . As in the second case it follows that $\deg \varphi \geq 1 + m(\varphi)(p^r - 1)$. Since on the other hand $\max(\tilde{J}) \leq (m(\eta) - 2)(p^r - 1)$, it also follows that $\deg \varphi \geq 1 + (m(\eta) - 1)(p^r - 1) \geq \max \tilde{J} + p^r$. Applying the induction hypothesis to φ and \tilde{J} yields the desired conclusion for η .

It remains to consider the case $\alpha_{m(\eta)} \leq p^r - 1$ and $m(\eta) \leq 2$. If $m(\eta) = 1$, then $\alpha_1 \geq p^r$, a contradiction. Therefore $\eta = x_1^{\alpha_1} x_2^{\alpha_2}$ with $\alpha_2 \neq 0$ and $\alpha_1 + \alpha_2 \geq \max\{2p^r - 1, \max J + p^r\}$. It follows that $\alpha_1 \geq p^r$. Then the element $\eta' = x_1^{\alpha_1 - p^r} x_2^{\alpha_2}$ belongs to $(x_1, x_2)^{\max J}$ which is contained in J , and so $\eta = x_1^{p^r} \eta' \in x_1^{p^r} J$. \square

Corollary 2.4. *Let $J = \prod_{k=2}^m (x_1, \dots, x_k)^{a_k}$ where $0 \leq a_k \leq p^r - 1$ for $k = 2, \dots, m$, and let q be a positive integer and $\eta \in J$ a monomial with $m(\eta) < q$ and $\deg \eta \geq 1 + \max\{(q - 1)(p^r - 1), \max(J)\}$. Then there exists $t \leq m(\eta)$ such that $x_q^{p^r-1} \eta \in x_t^{p^r} J$.*

Proof. Let $\eta' = x_q^{p^r-1} \eta$. We have $\deg \eta' \geq 1 + \max\{m(\eta')(p^r - 1), \max(J) + p^r - 1\}$ since $m(\eta') = q$. Thus by Lemma 2.3 there exists an integer $t \leq m(\eta') = q$ such that $\eta' \in x_t^{p^r} J$, and hence $x_q^{p^r-1} \eta \in x_t^{p^r} J$. Since $p^r - 1$ is the maximal power of x_q which divides η' , we have $t \neq q$ and so $t \leq m(\eta)$. \square

Lemma 2.5. *Let $J = \prod_{k=2}^m (x_1, \dots, x_k)^{a_k}$ with $0 \leq a_k \leq p^{r-e} - 1$ for $k = 2, \dots, m$ and integers $0 \leq e < r$. Let $I = J^{[p^e]}$ and $\eta \in I$ be a monomial such that $\deg \eta \geq 1 + \max\{m(\eta)(p^r - 1), \max(I) + p^r - p^e + m(\eta)(p^e - 1)\}$. Then there exists $t \leq m(\eta)$, such that $\eta \in x_t^{p^r} I$.*

Proof. We may write $\eta = v^{p^e} w$, $v \in G(J)$ and $w = \sigma_1^{p^e} \sigma_0$, where σ_0 and σ_1 are monomials, and $\sigma_0 \notin (x_1^{p^e}, \dots, x_{m(\eta)}^{p^e})$. Thus $\deg \sigma_0 \leq m(\eta)(p^e - 1)$ and the monomial $\eta' = v\sigma_1$ belongs to J . Since $\eta = \eta'^{p^e} \sigma_0$ it follows that

$$\begin{aligned} p^e \deg \eta' &= \deg \eta - \deg \sigma_0 \geq \deg \eta - m(\eta)(p^e - 1) \\ &\geq 1 + \max\{m(\eta)(p^r - p^e), p^e \max(J) + p^r - p^e\}. \end{aligned}$$

Therefore $\deg \eta' \geq (1/p^e) + \max\{m(\eta)(p^{r-e} - 1), \max(J) + p^{r-e} - 1\}$. Since $\deg \eta'$ is an integer we get $\deg \eta' \geq 1 + \max\{m(\eta)(p^{r-e} - 1), \max(J) + p^{r-e} - 1\}$. Note that $m(\eta') \leq m(\eta)$. Therefore by Lemma 2.3 there exists an integer t , $t \leq m(\eta') \leq m(\eta)$, such that $\eta' \in x_t^{p^{r-e}} J$. Thus $\eta \in x_t^{p^r} I$. \square

Applying Lemma 2.5 recursively we get

Corollary 2.6. *With the hypotheses of Lemma 2.5 suppose in addition that $\deg \eta \geq cp^r + 1 + \max\{m(\eta)(p^r - 1), \max(I) + p^r - p^e + m(\eta)(p^e - 1)\}$ for some integer $c \geq 0$. Then there exists a monomial σ of degree $c + 1$ such that $m(\sigma) \leq m(\eta)$ and $\eta \in \sigma^{p^r} I$.*

Lemma 2.7. *Let $J = \prod_{k=2}^m (x_1, \dots, x_k)^{a_k}$ with $0 \leq a_k \leq p^{r-e} - 1$ for $k = 2, \dots, m$ and integers $0 \leq e < r$. Let $I = J^{[p^e]}$, and let q be a positive integer and $\eta \in J$ a monomial with $m(\eta) < q$ and $\deg \eta \geq cp^r + 1 + \max\{(q - 1)(p^r - 1), \max(I) + (q - 1)(p^e - 1)\}$ for some integer $c \geq 0$. Then there exists a monomial of degree $c + 1$ such that $x_q^{p^r - 1} \eta \in \sigma^{p^r} I$ and $m(\sigma) \leq m(\eta)$.*

Proof. Set $\eta' = x_q^{p^r - 1} \eta$. Then

$$\deg \eta' \geq cp^r + 1 + \max\{m(\eta')(p^r - 1), \max(I) + p^r - p^e + m(\eta')(p^e - 1)\}.$$

So by Corollary 2.6 there exists a monomial σ with $m(\sigma) \leq m(\eta')$ and $\deg \sigma = c + 1$ such that $\eta' \in \sigma^{p^r} I$. Finally we have $m(\sigma) \leq m(\eta)$ because x_q does not divide σ since it appears only with power $p^r - 1$ in σ . \square

Lemma 2.8. *Let $I_t = \prod_{k=2}^{m_t} (x_1, \dots, x_k)^{a_{tk}}$ with $0 \leq a_{tk} \leq p^{e_t+1-e_t} - 1$, $1 \leq t \leq s$, and integers $0 \leq e_1 < \dots < e_s < r = e_{s+1}$. Let $I = \prod_{t=1}^s I_t^{[p^{e_t}]}$, and let q be a positive integer and $\eta \in I$ a monomial with $m(\eta) < q$ and $\deg \eta \geq cp^r + 1 + \max_{1 \leq t \leq s+1} \{\sum_{i=t}^s p^{e_i} \max(I_i) + (q - 1)(p^{e_t} - 1)\}$ for some integer $c \geq 0$. (Here $\sum_{i=t}^s p^{e_i} \max(I_i) = 0$ for $t = s + 1$.) Then there exists a monomial σ with $\deg \sigma = c + 1$, $m(\sigma) \leq m(\eta)$ and $x_q^{p^r - 1} \eta \in \sigma^{p^r} I$.*

Proof. We apply induction on s . The case $s = 1$ is given in Lemma 2.7. Let $d_j = \sum_{i=j}^s p^{e_i} \max(I_i) + (q - 1)(p^{e_j} - 1)$, $1 \leq j \leq s + 1$, and let $t \leq s$ be a maximal integer such that $d_t = \max\{d_j : 1 \leq j \leq s\}$. Then we have $d_j < d_t$ for $t < j \leq s$ and $d_j \leq d_t$ for $j < t$.

We now distinguish two cases. In case $t > 1$, write $\eta = \eta' \prod_{i=t}^s v_i^{p^{e_i}}$ with $v_i \in G(I_i)$ and $\eta' \in I' = \prod_{i=1}^{t-1} I_i^{[p^{e_i}]}$. We have $\deg(\eta') = \deg(\eta) - \sum_{i=t}^s p^{e_i} \max(I_i) \geq cp^r + 1 + d_t - \sum_{i=t}^s p^{e_i} \max(I_i) = cp^r + 1 + (q - 1)(p^{e_t} - 1)$. Choose the maximal integer $\varepsilon \geq 0$ such that $\deg(\eta') \geq cp^r + 1 + \varepsilon p^{e_t} + (q - 1)(p^{e_t} - 1)$. As $d_t = \max\{d_i : 1 \leq i \leq s\}$, we see that η' satisfies the necessary inequalities and so by induction hypothesis there exists a monomial τ with $\deg(\tau) = cp^r - \varepsilon + \varepsilon + 1$, $m(\tau) \leq m(\eta') \leq m(\eta)$ and $x_q^{p^{e_t} - 1} \eta' = \tau^{p^{\varepsilon}} \eta''$ for some $\eta'' \in I'$. Note that by the choice of ε , we have $\deg(\eta') \leq cp^r + 1 + \varepsilon p^{e_t} + q(p^{e_t} - 1)$, and so $p^{e_t} \deg(\tau) + \deg(\eta'') \leq cp^r + 1 + \varepsilon p^{e_t} + (q + 1)(p^{e_t} - 1)$. Hence $\deg(\eta'') \leq 1 - p^{e_t} + (q + 1)(p^{e_t} - 1) = q(p^{e_t} - 1)$.

Set $\rho = \tau \prod_{i=t}^s v_i^{p^{e_i - e_t}}$; then $\rho \in \tilde{I} = \prod_{i=t}^s I_i^{[p^{e_i - e_t}]}$, and $p^{e_t} \deg(\rho) = \deg(x_q^{p^{e_t} - 1} \eta) - \deg(\eta'') \geq cp^r + p^{e_t} + \max\{d_t, d_{s+1}\} - q(p^{e_t} - 1) = cp^r + 1 + a$ for $a = p^{e_t} b$ and $b = \max\{\sum_{j=i}^s p^{e_j - e_t} \max(I_j) + (q - 1)(p^{e_i - e_t} - 1) : t \leq i \leq s + 1\}$. Therefore $\deg(\rho) \geq cp^{r - e_t} + 1/p^{e_t} + b$. Using that $\deg \rho$ is an integer we get $\deg(\rho) \geq cp^{r - e_t} + 1 + b$. Thus we may apply the induction hypothesis to ρ, \tilde{I} , and hence there exists a monomial σ with $m(\sigma) \leq m(\rho) \leq m(\eta)$, $\deg(\sigma) = c + 1$ such that $x_q^{p^{r - e_t} - 1} \rho \in \sigma^{p^{r - e_t}} \tilde{I}$. Hence $x_q^{p^r - 1} \eta = x_q^{p^r - p^{e_t}} \rho^{p^{e_t}} \eta'' \in \sigma^{p^r} I$.

Finally we consider the case $t = 1$, and write $\eta = \varphi \prod_{i=2}^s v_i^{p^{e_i}}$, $v_i \in G(I_i)$ and $\varphi \in I_1^{[p^{e_1}]}$. We have $\deg(\varphi) = \deg(\eta) - \sum_{i=2}^s p^{e_i} \max(I_i) \geq cp^r + 1 + d_1 - \sum_{i=2}^s p^{e_i} \max(I_i) = cp^r + 1 + p^{e_1} \max(I_1) + (q - 1)(p^{e_1} - 1)$. However, $t = 1$ implies $d_2 < d_1$, and so $p^{e_1} \max(I_1) + (q - 1)(p^{e_1} - 1) > (q - 1)(p^{e_2} - 1)$. Thus φ satisfies the condition of Lemma 2.7. As in the previous case we choose the maximal integer $\varepsilon \geq 0$ such that $\deg(\varphi) \geq cp^r + 1 + \varepsilon p^{e_2} + (q - 1)(p^{e_2} - 1)$. Then by Lemma 2.7 there exists a monomial γ with $\deg(\gamma) = cp^{r - e_2} + 1 + \varepsilon$, $m(\gamma) \leq m(\eta)$ and $x_q^{p^{e_2} - 1} \varphi = \gamma^{p^{e_2}} \varphi'$ for some $\varphi' \in I_1^{[p^{e_1}]}$, and we see as above that $\deg(\varphi') \leq q(p^{e_2} - 1)$.

Set $\psi = \gamma' \prod_{i=2}^s v_i^{p^{e_i - e_2}}$; then $\psi \in \hat{I} = \prod_{i=2}^s I_i^{[p^{e_i - e_2}]}$, $m(\psi) \leq m(\eta)$, and

$$\begin{aligned} p^{e_2} \deg(\psi) &= \deg(x_q^{p^{e_2} - 1} \eta) - \deg(\varphi') \\ &\geq cp^r + p^{e_2} + \max\{d_j : 2 \leq j \leq s + 1\} - q(p^{e_2} - 1) \\ &= cp^r + 1 + \max\left\{\sum_{i=j}^s p^{e_i} \max(I_i) + (q - 1)(p^{e_j} - p^{e_2}) : 2 \leq j \leq s + 1\right\}. \end{aligned}$$

Since $\deg \psi$ is an integer we get

$$\begin{aligned} \deg(\psi) &\geq cp^{r - e_2} + 1 \\ &\quad + \max\left\{\sum_{i=j}^s p^{e_i - e_2} \max(I_i) + (q - 1)(p^{e_j - e_2} - 1) : 2 \leq j \leq s + 1\right\}. \end{aligned}$$

Thus we may apply our induction hypothesis to ψ and \hat{I} and conclude that $x_q^{p^{r - e_2} - 1} \psi \in \nu^{p^{r - e_2}} \hat{I}$ for some monomial ν with $\deg(\nu) = c + 1$ and $m(\nu) \leq m(\psi) \leq m(\eta)$. This yields the desired conclusion. \square

We are now in the position to prove Theorem 2.2.

Proof of 2.2. Let $\rho = \prod_{t=1}^s u_t^{p^{r_t}} w$, $u_t \in G(I_t)$ and w is a monomial such that $\deg(\rho) = d$. Let $j < m(\rho)$. We must show that $x_j \rho / x_{m(\rho)} \in I_{\geq d}$. Apply induction on s , case $s = 0$ being trivial. If $m(\rho) = m(\eta)$, $\eta = \prod_{t=1}^{s-1} u_t^{p^{r_t}} w$, then we may apply the induction hypothesis because $\deg(\eta) = d - \deg(u_s^{p^{r_s}}) \geq \max\{\delta'_t : 1 \leq t \leq s - 1\}$ for $\delta'_t = \sum_{i=t}^{s-1} p^{r_i} \max(I_i) + (m_t - 1)(p^{r_t} - 1)$. By the induction hypothesis $I' = \prod_{t=1}^{s-1} I_t^{[p^{r_t}]}$ has $I'_{\geq \deg(\eta)}$ stable and so $x_j \eta / x_{m(\eta)} \in I'_{\geq \deg(\eta)}$. Hence $x_j \rho / x_{m(\rho)} = (x_j \eta / x_{m(\eta)}) u_s^{p^{r_s}} \in I_{\geq d}$.

We may suppose from now on that $m(\rho) = m(u_s) > m(\eta)$. Set $q = m(u_s)$, $d_t = \sum_{i=t}^{s-1} p^{r_i} \max(I_i) + (q - 1)(p^{r_t} - 1)$. We have $\delta'_t \geq d_t$ if and only if $m_t \geq q$. In particular $\delta'_s \geq d_s$ since $m_s \geq q$. If $m_t < q$, then $\max(I_t) \leq (q - 1)(p^{r_{t+1} - r_t} - 1)$ since $a_{tk} \leq p^{r_{t+1} - r_t} - 1$, and so $d_{t+1} - \delta'_t = (q - 1)(p^{r_{t+1} - 1} - 1) - (m_t - 1)(p^{r_t} - 1) - p^{r_t} \max(I_t) > (q - 1)(p^{r_{t+1} - 1} - 1) - p^{r_t} \max(I_t) \geq 0$. Thus $\delta'_t < d_{t+1}$ if $m_t < q$. The same argument shows that $d_t \leq d_{t+1}$ if $m_t < q$.

By backwards induction on j we now show that

$$(2) \quad \max\{d_t : j \leq t \leq s\} \leq \max\{\delta'_t : j \leq t \leq s\}.$$

We have already seen that (2) holds for $j = s$. Suppose the inequality holds for $j + 1$. If $m_j \geq q$, then $\delta'_j \geq d_j$, and so (2) is implied by the induction hypothesis. If $m_j < q$, then $\delta'_j < d_{j+1}$, $d_j \leq d_{j+1}$, and hence $\delta'_j < d_{j+1} \leq \max\{d_t : j + 1 \leq t \leq s\} \leq \max\{\delta'_t : j + 1 \leq t \leq s\}$. Hence $\max\{\delta'_t : j \leq t \leq s\} = \max\{\delta'_t : j + 1 \leq t \leq s\} \geq \max\{d_t : j + 1 \leq t \leq s\} = \max\{d_t : j \leq t \leq s\}$, as desired.

Now since by (2) we have $\deg(x_j \eta) = 1 + \deg(\eta) = 1 + \deg(\rho) - p^{r_s} \max(I_s) = 1 + \max\{\delta'_t : 1 \leq t \leq s\} \geq 1 + \max\{d_t : 1 \leq t \leq s\}$, we may apply Lemma 2.8 for $s - 1$, $x_j \eta \in I'$ and q . Then there exists $e \leq m(\eta)$ such that $x_q^{p^{r_s}-1}(x_j \eta) = x_e^{p^{r_s}} \eta'$ for some $\eta' \in I'$. Thus $x_j \rho / x_{m(\rho)} = (x_e u_s / x_{m(u_s)})^{p^{r_s}} \eta'$ belongs to I , because I_s is stable. \square

For a monomial u , we set $\text{pa}(u) = \max_k: \mu_k \neq 0 \{D_k\}$ if u is not a multiple of x_1 . Otherwise $u = x_1^{\mu_1} v$ such that $v \notin (x_1)$, and we set $\text{pa}(u) = \mu_1 + \text{pa}(v)$ (cf. Section 1). By our main theorem we have $\text{reg}(u) = \text{pa}(u)$. More generally we get

Corollary 2.9. *Let I be a p -Borel ideal with Borel generators u_1, \dots, u_m . Then*

$$\text{reg}(I) \leq \max\{\text{pa}(u_1), \dots, \text{pa}(u_m)\},$$

and equality holds if I is principal p -Borel.

Proof. For each $i = 1, \dots, m$, the ideal $\langle u_i \rangle_{\geq d}$ is stable for $d \geq \text{pa}(u_i)$. Thus $I_{\geq d}$ is stable for $d \geq \max\{\text{pa}(u_1), \dots, \text{pa}(u_m)\}$. Therefore the assertion follows from Proposition 2.1. \square

REFERENCES

- [1] A. Aramova and J. Herzog, p -Borel principal ideals, Illinois J. Math. 41, no. 1 (1997), 103–121. MR **98i**:13020
- [2] D. Bayer and M. Stillman, A criterion for detecting m -regularity, Invent. Math. 87 (1987), 1–11. MR **87k**:13019
- [3] S. Eliahou and M. Kervaire, Minimal resolutions of some monomial ideals, J. of Algebra 129 (1990) 1–25. MR **91b**:13019
- [4] D. Eisenbud, A. Reeves and B. Totaro, Initial ideals, Veronese subrings, and rates of algebras, Adv Math. 109 (1994), 168–187. MR **96d**:13030
- [5] V. Ene, G. Pfister and D. Popescu, Betti numbers for p -stable ideals, Comm. Algebra 28 (2000), 1515–1531. CMP 2000:08
- [6] K. Pardue, Nonstandard Borel-fixed ideals, Dissertation, Brandeis University, 1994.

FB6 MATHEMATIK UND INFORMATIK, UNIVERSITÄT – GHS – ESSEN, POSTFACH 103764, 45117 ESSEN, GERMANY

E-mail address: juergen.herzog@uni-essen.de

INSTITUTE OF MATHEMATICS, UNIVERSITY OF BUCHAREST, P.O. BOX 1-764, BUCHAREST 70700, ROMANIA

E-mail address: dorin@stoilow.imar.ro