

VICTORIS CONTINUOUS SELECTIONS AND DISCONNECTEDNESS-LIKE PROPERTIES

VALENTIN GUTEV AND TSUGUNORI NOGURA

(Communicated by Alan Dow)

ABSTRACT. Suppose that X is a Hausdorff space such that its Vietoris hyperspace $(\mathcal{F}(X), \tau_V)$ has a continuous selection. Do disconnectedness-like properties of X depend on the variety of continuous selections for $(\mathcal{F}(X), \tau_V)$ and vice versa? In general, the answer is “yes” and, in some particular situations, we were also able to set proper characterizations.

1. INTRODUCTION

Let X be a Hausdorff space, and let $\mathcal{F}(X)$ be the family of the non-empty closed subsets of X . Also, let $\mathcal{D} \subset \mathcal{F}(X)$. A map $f : \mathcal{D} \rightarrow X$ is a *selection* for \mathcal{D} if $f(S) \in S$ for every $S \in \mathcal{D}$. A map $f : \mathcal{D} \rightarrow X$ is a *continuous* selection for \mathcal{D} if it is a selection which is continuous with respect to the relative Vietoris topology τ_V on \mathcal{D} . Let us recall that the *Vietoris topology* τ_V on $\mathcal{F}(X)$ is generated by all collections of the form

$$\langle \mathcal{V} \rangle = \left\{ S \in \mathcal{F}(X) : S \cap V \neq \emptyset, V \in \mathcal{V}, \text{ and } S \subset \bigcup \mathcal{V} \right\},$$

where \mathcal{V} runs over the finite families of open subsets of X .

In what follows, we use $\text{Sel}(X)$ to denote the set of all continuous selections for $\mathcal{F}(X)$, and $\dim(X)$ to denote the *covering dimension* of X . Also, we denote by $\text{ind}(X)$ the *small inductive dimension* of X . Finally, we shall say that X is *zero-dimensional* if it has a base of clopen sets (i.e., if $\text{ind}(X) = 0$), and that X is *strongly zero-dimensional* if $\dim(X) = 0$. Note that every strongly zero-dimensional space is zero-dimensional but the converse fails ([12], [13], see also [11]).

The dimension-type of restrictions play an important role in the selection theory for hyperspaces. For instance, $\text{Sel}(X) \neq \emptyset$ for every strongly zero-dimensional completely metrizable space X (see [3], [5]). On the other hand, we have the following two results in the opposite direction.

Theorem 1.1 ([8]). *If X is a compact Hausdorff space with $\text{Sel}(X) \neq \emptyset$, then it is a linear ordered topological space. In particular, $\dim(X) \leq 1$.*

Theorem 1.2 ([10]). *If X is a compact Hausdorff space with $\text{Sel}(X) \neq \emptyset$, then it has finitely many connected components if and only if $\text{Sel}(X)$ is finite. In particular, $\dim(X) = 1$ provided X is infinite and $\text{Sel}(X)$ is finite.*

Received by the editors November 17, 1999 and, in revised form, January 17, 2000.

2000 *Mathematics Subject Classification*. Primary 54C65, 54B20, 54F45.

Key words and phrases. Selections, hyperspaces, zero-dimensionality.

In the present paper we are interested in relations between the set $\text{Sel}(X)$ and zero-dimensionality of X . As Theorems 1.1 and 1.2 suggest, we may expect “sufficiently many” continuous selections for $\mathcal{F}(X)$ provided $\dim(X) = 0$ and $\text{Sel}(X) \neq \emptyset$. In fact, this is the first result of the paper. The following theorem will be proved in the next section.

Theorem 1.3. *If X is a zero-dimensional Hausdorff space such that $\text{Sel}(X) \neq \emptyset$, then the set $\{f(X) : f \in \text{Sel}(X)\}$ is dense in X .*

The proof of Theorem 1.3 does not involve such complicated arguments. However, in general, the conclusion “ $\{f(X) : f \in \text{Sel}(X)\}$ is dense in X ” cannot be strengthened to “ $\{f(X) : f \in \text{Sel}(X)\} = X$ ” (see Example 2.1). On the other hand, this becomes possible provided X is a first countable space which allows us to obtain also the converse.

Theorem 1.4. *Let X be a first countable Hausdorff space such that $\text{Sel}(X) \neq \emptyset$. Then it is zero-dimensional if and only if for every point $x \in X$ there exists $f_x \in \text{Sel}(X)$ such that $f_x^{-1}(x) = \{S \in \mathcal{F}(X) : x \in S\}$.*

In view of Theorem 1.4, it would be interesting to know if the converse of Theorem 1.3 holds as well. Here we have only the following partial result.

Theorem 1.5. *If X is a Hausdorff space such that $\{f(X) : f \in \text{Sel}(X)\}$ is dense in X , then it is totally disconnected.*

Since every totally disconnected locally compact space is zero-dimensional, by Theorems 1.3 and 1.5, we get the following consequence.

Corollary 1.6. *Let X be a locally compact Hausdorff space such that $\text{Sel}(X) \neq \emptyset$. Then $\text{ind}(X) = 0$ if and only if $\{f(X) : f \in \text{Sel}(X)\}$ is dense in X .*

A word should be said about the proofs of Theorems 1.4 and 1.5. A preparation for that is done in Sections 3 and 4. A proof of Theorem 1.5 is obtained in Section 5. Since the proof of Theorem 1.4 involves that of Theorem 1.5, it will be finally accomplished in Section 6 of the paper.

2. SELECTIONS AND CLOPEN SETS

Throughout this section, and in the sequel, X is always a Hausdorff space. We first prove Theorem 1.3. In fact, this theorem is a consequence of the following lemma.

Lemma 2.1. *Let X be a space such $\text{Sel}(X) \neq \emptyset$, and let G be a non-empty clopen subset of X . Then there exists a selection $g \in \text{Sel}(X)$ with $g(X) \in G$.*

Proof. Note that the sets

$$\mathcal{G}_0 = \{S \in \mathcal{F}(X) : S \cap G = \emptyset\} \quad \text{and} \quad \mathcal{G}_1 = \{S \in \mathcal{F}(X) : S \cap G \neq \emptyset\}$$

constitute a disjoint and τ_V -open cover of $\mathcal{F}(X)$. Define another subset \mathcal{G}_2 of $\mathcal{F}(X)$ by $\mathcal{G}_2 = \{S \in \mathcal{F}(X) : S \subset G\}$. Take a selection $f \in \text{Sel}(X)$. Note that each $f_i = f|_{\mathcal{G}_i}$, $i = 0, 1, 2$, is a continuous selection for \mathcal{G}_i . We now consider the map $\varphi : \mathcal{G}_1 \rightarrow \mathcal{G}_2$ defined by $\varphi(S) = S \cap G$ for every $S \in \mathcal{G}_1$. This map is continuous with respect to the relative Vietoris topologies on \mathcal{G}_1 and \mathcal{G}_2 . Finally, define a map $g : \mathcal{F}(X) \rightarrow X$ by $g|_{\mathcal{G}_0} = f_0$ and $g|_{\mathcal{G}_1} = f_2 \circ \varphi$. Clearly, g is a selection for $\mathcal{F}(X)$. That $g \in \text{Sel}(X)$ follows from the fact that φ is continuous and \mathcal{G}_0 and \mathcal{G}_1

are τ_V -open in $\mathcal{F}(X)$. This g is the required one because $X \in \mathcal{G}_1$ and, therefore, $g(X) = f_2(\varphi(X)) = f_2(X \cap G) = f(G) \in G$. \square

Example 2.2. There exists a space X_p with only one non-isolated point $p \in X_p$ such that $\text{Sel}(X_p) \neq \emptyset$ and $f(X_p) \neq p$ for every $f \in \text{Sel}(X_p)$.

Proof. Let $p = \omega_1$ be the first uncountable ordinal, and let $X_p = \{\alpha + 1 : \alpha < \omega_1\} \cup \{p\}$. We endow X_p with the relative topology as a subspace of the ordinal space $\omega_1 + 1$. Then, X_p is a space with only one non-isolated point p such that every countable subset of $X_p \setminus \{p\}$ is closed in X_p . Hence, by a result of [1], $f(X_p) \neq p$ for every $f \in \text{Sel}(X_p)$. On the other hand, X_p is a linear ordered space and $f(S) = \min S$, $S \in \mathcal{F}(X_p)$, defines a continuous selection for $\mathcal{F}(X_p)$. \square

In contrast to Example 2.2, we have the following lemma demonstrating the relationship of the problem with the convergent structure of X_p in p .

In what follows, we use ω to denote the first infinite ordinal. As a space, every ordinal has the usual order topology.

Lemma 2.3. *Let X be a space, $p \in X$, and let Y_q be the quotient space on the disjoint union $X \sqcup (\omega + 1)$ obtained by identifying the points p and ω to a single point $q \in Y_q$. If $\text{Sel}(Y_q) \neq \emptyset$, then there exists $g \in \text{Sel}(Y_q)$ with $g(Y_q) = q$.*

Proof. Let

$$\mathcal{V}_0 = \{F \in \mathcal{F}(Y_q) : 0 \notin F\} \quad \text{and} \quad \mathcal{V}_1 = \{F \in \mathcal{F}(Y_q) : 0 \in F\}.$$

Since $0 \in Y_q$ is an isolated point, this defines a disjoint τ_V -open cover $\{\mathcal{V}_0, \mathcal{V}_1\}$ of $\mathcal{F}(Y_q)$. Consider the sets

$$\mathcal{V}_1^0 = \{F \in \mathcal{V}_1 : \omega \setminus F \neq \emptyset\} \quad \text{and} \quad \mathcal{V}_1^1 = \{F \in \mathcal{V}_1 : \omega \subset F\}.$$

Next, for every $F \in \mathcal{V}_1^0$, define $n(F) = \min\{n \in \omega \cap F : n + 1 \notin F\}$. Now, take $f \in \text{Sel}(Y_q)$ and then define another selection $g : \mathcal{F}(Y_q) \rightarrow Y_q$ for $\mathcal{F}(Y_q)$ by $g|_{\mathcal{V}_0} = f|_{\mathcal{V}_0}$ while $g(F) = n(F)$ if $F \in \mathcal{V}_1^0$ and $g(F) = q$ otherwise. Since $Y_q \in \mathcal{V}_1^1$, we have $g(Y_q) = q$. Thus to finish the proof it only remains to show that g is continuous. In fact, it suffices to show that $g|_{\mathcal{V}_1}$ is continuous because $g|_{\mathcal{V}_0} = f|_{\mathcal{V}_0}$. Take an $F \in \mathcal{V}_1$. We distinguish the following two cases. If $F \in \mathcal{V}_1^0$, then $g(F) = n(F) < \omega$ and $\{n < \omega : n \leq n(F)\} \subset F$. Set $\mathcal{U} = \{\{n\} : n \leq n(F)\} \cup \{Y_q \setminus \{n(F) + 1\}\}$. Then $\langle \mathcal{U} \rangle \subset \mathcal{V}_1^0$ is a τ_V -neighbourhood of F such that $n(S) = n(F)$ for every $S \in \langle \mathcal{U} \rangle$. Hence, $g(\langle \mathcal{U} \rangle) = \{g(F)\}$. Suppose now that $F \in \mathcal{V}_1^1$, and let V be a neighbourhood of $q = g(F)$. Take an $m < \omega$ such that $\{n < \omega : n \geq m\} \subset V$. Then, let $\mathcal{U} = \{\{n\} : n \leq m\} \cup \{Y_q\}$. In this way, we get a τ_V -neighbourhood $\langle \mathcal{U} \rangle$ of F such that $g(\langle \mathcal{U} \rangle) \subset V$. Indeed, take an $S \in \langle \mathcal{U} \rangle$. In case $\omega \subset S$, by definition, we have $g(S) = q \in V$. Otherwise, $\{n \in \omega \cap S : n \leq m\} \subset S$ implies that $n(S) \geq m$. Therefore, $g(S) = n(S) \in V$. \square

3. SELECTIONS AND ORDER-LIKE RELATIONS

In this section we collect some known facts we need for the proof of Theorem 1.5. Suppose that X is a space with $\text{Sel}(X) \neq \emptyset$. Following [7], to every selection $f \in \text{Sel}(X)$ we associate an order-like relation \preceq_f on X defined for $x, y \in X$ by

$$x \preceq_f y \quad \text{if and only if} \quad f(\{x, y\}) = x.$$

In what follows, we shall refer to “ \preceq_f ” as an f -order on X . Also, let us agree to write $x \prec_f y$ provided $x \preceq_f y$ and $x \neq y$.

Finally, for every $f \in \text{Sel}(X)$ and $x \in X$, we consider the following special subsets of X :

$$\begin{aligned} (-\infty, x]_f &= \{z \in X : z \preceq_f x\} \quad \text{and} \quad [x, +\infty)_f = \{z \in X : x \preceq_f z\}, \\ (-\infty, x)_f &= \{z \in X : z \prec_f x\} \quad \text{and} \quad (x, +\infty)_f = \{z \in X : x \prec_f z\}. \end{aligned}$$

The following observation is an immediate consequence of the continuity of f (see [7, Lemma 7.2]) and we left the corresponding arguments to the interested reader.

Lemma 3.1. *Let X be a space, and let $f \in \text{Sel}(X)$. Then, for every $x \in X$, the sets $(-\infty, x)_f$ and $(x, +\infty)_f$ are open in X . In particular, $(-\infty, x]_f$ and $[x, +\infty)_f$ are closed in X .*

It should be mentioned that, in contrast to usual linear orders on X , the f -orders are not engaged to be *transitive*. However, in case of connected spaces, this is so. The following result is a partial case of [7, Lemmas 7.2 and 7.3] (see also [9, Lemma 10]).

Lemma 3.2. *Let X be a connected space such that $\text{Sel}(X) \neq \emptyset$. Then, $|\text{Sel}(X)| \leq 2$ and, for every $f \in \text{Sel}(X)$, the following hold.*

- (1) *The f -order on X is transitive.*
- (2) *$f(X) = \min_{\preceq_f} X$.*
- (3) *$g(X) = \max_{\preceq_f} X$ provided $g \in \text{Sel}(X) \setminus \{f\}$.*

In case of connected subsets of X we also have the following property of f -orders.

Lemma 3.3. *Let X be a space, $f \in \text{Sel}(X)$, and let $A \subset X$ be connected. Also, let $x, y \in A$ be such that $x \preceq_f y$. Then,*

$$(x, y)_f = \{z \in X : x \prec_f z \prec_f y\} \subset A.$$

Proof. Suppose that there exists $z \in (x, y)_f \setminus A$. Then, according to Lemma 3.1, $U = (-\infty, z]_f \cap A$ is a clopen (in A) subset of A because $U = (-\infty, z)_f \cap A$. Note that $x \in U$ while $y \in A \setminus U$. Since A is connected, this is impossible. \square

4. SELECTIONS AND COMPONENTS

Let X be a space. For every $x \in X$, we shall use $\mathcal{C}[x]$ to denote the *component* of the point x and $\mathcal{C}^*[x]$ the corresponding *quasi-component*. Let us recall that

$$\mathcal{C}[x] = \bigcup \{C \subset X : x \in C \text{ and } C \text{ is connected}\}$$

and, respectively,

$$\mathcal{C}^*[x] = \bigcap \{C \subset X : x \in C \text{ and } C \text{ is clopen}\}.$$

In this section we establish the following result which may have some independent interest.

Theorem 4.1. *Let X be a space with $\text{Sel}(X) \neq \emptyset$. Then, $\mathcal{C}^*[x] = \mathcal{C}[x]$ for every point $x \in X$.*

To prepare for the proof of Theorem 4.1, we need the following lemma.

Lemma 4.2. *Let X be such that $\text{Sel}(X) \neq \emptyset$, and let $f \in \text{Sel}(X)$. Then for every $x \in X$ and $y, z \in \mathcal{C}^*[x]$, with $y \preceq_f z$, we have*

$$[y, z]_f = \{t \in X : y \preceq_f t \preceq_f z\} \subset \mathcal{C}^*[x].$$

Proof. Suppose that there are points $y, z \in \mathcal{C}^*[x]$ and $t \in X \setminus \mathcal{C}^*[x]$ such that $y \prec_f t \prec_f z$. Since $t \notin \mathcal{C}^*[x]$, there exists a clopen subset $V \subset X$ such that $\mathcal{C}^*[x] \subset V$ and $t \notin V$. Then, by Lemma 3.1, $U = (-\infty, t]_f \cap V$ defines a clopen (in X) neighbourhood of y because $U = (-\infty, t]_f \cap V$. Note that $z \notin U$. However, this is impossible because $z \in \mathcal{C}^*[x] = \mathcal{C}^*[y] \subset U$. A contradiction. \square

Proof of Theorem 4.1. Let $x \in X$. It suffices to show that $\mathcal{C}^*[x]$ is connected. Towards this end, note that, by Lemma 4.2,

$$\mathcal{C}^*[x] = \bigcup \{[y, z]_f : y, z \in \mathcal{C}^*[x], y \preceq_f z, \text{ and } y \preceq_f x \preceq_f z\}.$$

Hence, it will be sufficient to show that for every $y, z \in \mathcal{C}^*[x]$, with $y \preceq_f z$, the set $[y, z]_f$ is connected. Suppose to the contrary that $[y, z]_f$ is not connected for some points $y, z \in \mathcal{C}^*[x]$ with $y \preceq_f z$. Then, there exists a clopen (in $[y, z]_f$) neighbourhood $W \subset [y, z]_f$ of z such that $[y, z]_f \setminus W \neq \emptyset$. Take a point $t \in [y, z]_f \setminus W$ and then set $T = W \cap [t, z]_f$. Thus, we get a clopen (in $[t, z]_f$) neighbourhood T of z such that $t \notin T$. Then, the set $G = T \cup [z, +\infty)_f$ is clopen in X . Indeed, according to Lemma 3.1, G is closed in X as a union of two closed subsets. To show that it is also open in X , note that there exists an open subset $E \subset (t, +\infty)_f$ such that $E \cap [t, z]_f = T$ because $t \notin T \subset [t, z]_f \subset [t, +\infty)_f$. Hence, by Lemma 3.1, the set G is open in X because $G = E \cup (z, +\infty)_f$. Thus, G is clopen in X . This however is impossible because $t \notin G$ and $z \in G$, while $t, z \in \mathcal{C}^*[x]$. A contradiction. \square

5. PROOF OF THEOREM 1.5

Let X be such that the set $D = \{f(X) : f \in \text{Sel}(X)\}$ is dense in X . Suppose if possible that X is not totally disconnected. Then, by Theorem 4.1, X must contain an infinite closed connected set A . Let $B = \{g(A) : g \in \text{Sel}(A)\}$, and let $f \in \text{Sel}(X)$. Note that, by Lemmas 3.2 and 3.3, there exists $a = \min_{\preceq_f} A$. According to the same lemmas, we consider the subset $H = \bigcup \{(a, x)_f : x \in A\}$ of $A \setminus B$. Note that it is non-empty because A is infinite while, by Lemma 3.1, it is open in X . Then, by hypothesis, there exists a selection $h \in \text{Sel}(X)$ with $h(X) \in H$. Since h is continuous, there now exists a finite open cover \mathcal{U} of X such that $h(\langle \mathcal{U} \rangle) \subset H$. Take a finite $F \subset X \setminus A$ so that $Z = F \cup A \in \langle \mathcal{U} \rangle$. Then, we have that $h(Z) \in H \subset A$. Set $k = h|_{\mathcal{F}(Z)}$ and $\mathcal{A} = \{S \cup F : S \in \mathcal{F}(A)\}$. Note that $\psi : \mathcal{F}(A) \rightarrow \mathcal{A}$, defined by $\psi(S) = S \cup F$, $S \in \mathcal{F}(A)$, becomes a continuous onto map. Since $\mathcal{F}(A)$ is connected (because so is A ; see [7, Theorem 4.10]), this implies that \mathcal{A} is also connected. On the other hand, $k^{-1}(A)$ is a τ_V -clopen subset of $\mathcal{F}(Z)$ because A is clopen in Z . Also, $A \cup F = Z \in k^{-1}(A)$ which finally implies that $\mathcal{A} \cap k^{-1}(A) \neq \emptyset$. Thus, $\mathcal{A} \subset k^{-1}(A)$. We may now define a continuous selection $g : \mathcal{F}(A) \rightarrow A$ by letting $g = k \circ \psi$. However, this selection has the property that $g(A) = k(\psi(A)) = k(A \cup F) = h(A \cup F) \in H \subset A \setminus B$. A contradiction.

6. PROOF OF THEOREM 1.4

Let X be as in Theorem 1.4, and let $\text{ind}(X) = 0$. Take an $f \in \text{Sel}(X)$, a point $x \in X$, and a decreasing clopen base $\mathcal{U} = \{U_n : n < \omega\}$ of x in X such that $U_0 = X$. For reasons of convenience, let $U_\omega = \{x\}$. Now, for every $n \leq \omega$ define a subset $\mathcal{F}_n = \{S \in \mathcal{F}(X) : S \cap U_n \neq \emptyset\}$. Note that $\mathcal{F}_\omega = \bigcap \{\mathcal{F}_n : n < \omega\}$ because \mathcal{U} is a base at x , while $\mathcal{F}_0 = \mathcal{F}(X)$ because $U_0 = X$. Then, define a map $\theta : \mathcal{F}(X) \rightarrow \omega + 1$ by setting $\theta(S) = \max\{n \leq \omega : S \in \mathcal{F}_n\}$ for every $S \in \mathcal{F}(X)$. Also, for every $n \leq \omega$, define a map $\varphi_n : \mathcal{F}_n \rightarrow \mathcal{F}(X)$ by $\varphi_n(S) = S \cap U_n$, $S \in \mathcal{F}_n$. Finally, define

a map $g : \mathcal{F}(X) \rightarrow X$ by $g(S) = f(\varphi_{\theta(S)}(S))$ for every $S \in \mathcal{F}(X)$. Thus, we get a selection g for $\mathcal{F}(X)$ such that $g^{-1}(x) = \mathcal{F}_\omega$. It only remains to show that g is continuous. Clearly, g is continuous at the singleton $\{x\}$. So, take an $F \in \mathcal{F}(X)$ with $F \neq \{x\}$. We distinguish the following two cases. If $g(F) = x$, then for every $n < \omega$ let $\mathcal{V}_n = \{U_n, X \setminus U_n\}$. In this case, $g(\langle \mathcal{V}_n \rangle) \subset U_n$. Indeed, let $S \in \langle \mathcal{V}_n \rangle$. Then, $S \cap U_n \neq \emptyset$ implies $\theta(S) \geq n$. Hence, $g(S) = f(\varphi_{\theta(S)}(S)) \in U_{\theta(S)} \subset U_n$ because \mathcal{U} is decreasing. Therefore, g is continuous at F because \mathcal{U} is a local base at $g(F)$ while the set $\{n < \omega : F \notin \langle \mathcal{V}_n \rangle\}$ is finite. Suppose now that $g(F) \neq x$. The definition of g implies that $m = \theta(F) < \omega$. Then, set $\mathcal{W} = \{U_m \setminus U_{m+1}, X \setminus U_{m+1}\}$. We have that $F \in \langle \mathcal{W} \rangle$ because $F \cap U_{m+1} = \emptyset$. On the other hand, $S \in \langle \mathcal{W} \rangle$ implies that $\theta(S) = m$ because \mathcal{U} is decreasing, $S \cap U_m \neq \emptyset$ and $S \cap U_{m+1} = \emptyset$. This finally implies that g is continuous at F because $\langle \mathcal{W} \rangle \subset \mathcal{F}_m$ and $\varphi_m|_{\langle \mathcal{W} \rangle}$ is continuous.

Suppose now that, for every $x \in X$, there exists $f_x \in \text{Sel}(X)$ such that $f_x^{-1}(x) = \{S \in \mathcal{F}(X) : x \in S\}$. Next, take a point $x \in X$ and then set $g = f_x$. Also, let V be a neighbourhood of x such that $F = X \setminus V$ is non-empty, i.e. $F \in \mathcal{F}(X)$. Then, by hypothesis, $g(F \cup \{x\}) = x$ and $g(F) \neq x$. Hence, by Theorem 1.5, there exists a clopen set U such that $g(F) \in U$ and $x \notin U$. In this way, we get a τ_V -clopen neighbourhood $g^{-1}(U)$ of F in $\mathcal{F}(X)$. Let $\mathcal{M} \subset g^{-1}(U)$ be a chain which is maximal with respect to the usual set-theoretical inclusion and $F \in \mathcal{M}$. Then, there exists $M = \max \mathcal{M}$ because $g^{-1}(U)$ is τ_V -closed (see [2], [4], [6]). Indeed, it suffices to show that $M = \bigcup \mathcal{M} \in g^{-1}(U)$. Take a basic τ_V -neighbourhood $\langle \mathcal{W} \rangle$ of M . Then, for every $W \in \mathcal{W}$ there exists $M_W \in \mathcal{M}$ with $M_W \cap W \neq \emptyset$ because $(\bigcup \mathcal{M}) \cap W \neq \emptyset$. Then, $M_W = \bigcup \{M_W : W \in \mathcal{W}\} \in \langle \mathcal{W} \rangle \cap \mathcal{M}$ because \mathcal{M} is a chain. Hence, in particular, $\langle \mathcal{W} \rangle \cap g^{-1}(U) \neq \emptyset$ which finally implies that $M \in g^{-1}(U)$ because $g^{-1}(U)$ is τ_V -closed. Having established this, let us also observe that M is open because $g^{-1}(U)$ is τ_V -open. Namely, $M \in \langle \mathcal{U} \rangle \subset g^{-1}(U)$ for some finite family \mathcal{U} of open subsets of X . Then, $M = \bigcup \mathcal{U}$ because \mathcal{M} is maximal with respect to the inclusion. Thus, M is a clopen subset of X which contains F because $F \in \mathcal{M}$. Finally note that $g(M) \in U$ which implies $x \notin M$. Hence, $G = X \setminus M$ is a clopen neighbourhood of x with $G \subset V$. This completes the proof of Theorem 1.4.

REFERENCES

1. G. Artico, U. Marconi, R. Moresco and J. Pelant, *Selectors and Scattered Spaces*, Topology Appl., to appear.
2. D. Bertacchi and C. Costantini, *Existence of selections and disconnectedness properties for the hyperspace of an ultrametric space*, Topology Appl. **88** (1998), 179–197. MR **99g**:54003
3. M. Choban, *Many-valued mappings and Borel sets. I*, Trans. Moscow Math. Soc. **22** (1970), 258–280.
4. C. Costantini and V. Gutev, *Recognizing special metrics by topological properties of the “metric”-Proximal hyperspace*, preprint.
5. R. Engelking, R. W. Heath, and E. Michael, *Topological well-ordering and continuous selections*, Invent. Math. **6** (1968), 150–158. MR **39**:6272
6. V. Gutev and T. Nogura, *Selections for Vietoris-like hyperspace topologies*, Proc. London Math. Soc. **80** (1) (2000), 235–256. CMP 2000:04
7. E. Michael, *Topologies on spaces of subsets*, Trans. Amer. Math. Soc. **71** (1951), 152–182. MR **13**:54f
8. J. van Mill and E. Wattel, *Selections and orderability*, Proc. Amer. Math. Soc. **83** (1981), 601–605. MR **82i**:54038
9. T. Nogura and D. Shakhmatov, *Characterizations of intervals via continuous selections*, Rendiconti del Circolo Matematico di Palermo, Serie II **56** (1997), 317–328. MR **99d**:54012

10. T. Nogura and D. Shakhmatov, *Spaces which have finitely many continuous selections*, Bollettino U. M. I. (7) **11-A** (1997), 723–729. MR **99a**:54011
11. T. C. Przymusiński, *On the dimension of product spaces and an example of M. Wage*, Proc. Amer. Math. Soc. **76** (1979), 315–321. MR **80f**:54033
12. M. L. Wage, *The dimension of product spaces*, preprint (1977).
13. M. L. Wage, *The dimension of product spaces*, Proc. Mat. Acad. Sci. USA **75** (1978), 4671–4672. MR **80a**:54064

SCHOOL OF MATHEMATICAL AND STATISTICAL SCIENCES, FACULTY OF SCIENCE, UNIVERSITY OF NATAL, KING GEORGE V AVENUE, DURBAN 4041, SOUTH AFRICA
E-mail address: `gutev@sci.und.ac.za`

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, EHIME UNIVERSITY, MATSUYAMA, 790 JAPAN
E-mail address: `nogura@ehimegw.dpc.ehime-u.ac.jp`