

## NILPOTENCY DEGREE OF COHOMOLOGY RINGS IN CHARACTERISTIC 3

PHAM ANH MINH

(Communicated by Stephen D. Smith)

*To Phuong and Nin*

ABSTRACT. The purpose of this note is to provide a 3-group  $\mathcal{G}$  whose mod-3 cohomology ring has a nilpotent element  $\xi \in H^*(\mathcal{G})$  satisfying  $\xi^3 \neq 0$ .

### 1. STATEMENT OF THE MAIN RESULT

For every  $p$ -group  $G$ , denote by  $H^*(G)$  the mod- $p$  cohomology algebra of  $G$ . We are now interested in the nilpotency degrees of elements of  $H^*(G)$ . For the case  $p = 2$ , in [1], [2], it was shown that, given any positive integer  $n$ , there exists a 2-group whose cohomology ring has elements of nilpotency degree  $n + 1$ . However, the methods given there do not generalize to the case of odd characteristic  $p$ . In this case, it seems that, until now, we do not have any example of elements of  $H^*(G)$  having nilpotency degrees greater than  $p$ .

The purpose of this note is to provide a 3-group  $\mathcal{G}$  whose cohomology ring has a nilpotent element  $\xi \in H^*(\mathcal{G})$  satisfying  $\xi^3 \neq 0$ . The group  $\mathcal{G}$  is obtained as follows. Let

$$\mathbb{E} = \langle a_1, a_2 | a_1^3 = a_2^3 = [a_1, a_2]^3 = [a_1, [a_1, a_2]] = [a_2, [a_1, a_2]] = 1 \rangle$$

be the extraspecial 3-group of order  $3^3$  and of exponent 3, and let

$$K = C_3^4 = \langle a_3, \dots, a_6 | a_i^3 = [a_i, a_j] = 1 \rangle$$

be the elementary abelian 3-group of rank 4. Set  $G = \mathbb{E} \times K$ . Define  $u_i \in H^1(G) = \text{Hom}(G, \mathbb{Z}/3)$ ,  $v_i \in H^2(G)$ ,  $1 \leq i \leq 6$ , by

$$\begin{aligned} u_i(a_j) &= \delta_{ij}, \\ v_i &= \beta u_i, \end{aligned}$$

with  $\delta_{ij}$  the Kronecker symbol and  $\beta$  the Bockstein homomorphism. For  $3 \leq i \leq 6$ , consider  $u_i, v_i$  as elements of  $H^*(K)$  via the restriction map. So, by the Künneth formula,

$$H^*(K) = \Lambda[u_3, \dots, u_6] \otimes \mathbb{F}_3[v_3, \dots, v_6],$$

and  $H^*(G)$  may be identified with

$$H^*(\mathbb{E}) \otimes H^*(K).$$

---

Received by the editors February 21, 2000 and, in revised form, June 12, 2000.  
2000 *Mathematics Subject Classification*. Primary 20J06; Secondary 20D15, 55R40.

Consider the central extension

$$(\mathcal{G}) \quad 0 \rightarrow \mathbb{Z}/3 \xrightarrow{i} \mathcal{G} \rightarrow G \rightarrow 1$$

corresponding to the cohomology class  $z = v_1 + u_3u_4 + u_5u_6 \in H^2(G)$ . We will prove

**Theorem.** *There exists a nilpotent element  $\xi \in H^2(\mathcal{G})$  satisfying  $\xi^3 \neq 0$ .*

However, for  $p > 3$ , our method could not be applied. So, the existence of a  $p$ -group  $G$  having a cohomology class of nilpotency degree greater than  $p > 3$  still remains an open problem. Besides, we could not provide any example of cohomology classes of a 3-group having nilpotency degrees greater than a given  $n$ , although it is known that, for any  $p \geq 2$  and for any  $p$ -group  $G$  of order  $p^m$ , nilpotency degrees of nilpotent elements of  $H^*(G)$  are bounded above by  $p^{m-1}$  (see [5]).

## 2. PROOF OF THE MAIN RESULT

Via the restriction map, for  $1 \leq i \leq 2$ ,  $u_i$  and  $v_i$  can be considered as elements of  $H^*(\mathbb{E})$ . Set  $c = [a_1, a_2]$  and let  $A_0 = \langle a_1, c \rangle$ ,  $A_1 = \langle a_1a_2, c \rangle$ ,  $A_2 = \langle a_1^2a_2, c \rangle$ ,  $A_3 = \langle a_2, c \rangle$  be elementary abelian subgroups of  $\mathbb{E}$ . Set  $s_0 = \text{Res}_{A_0}^{\mathbb{E}}(u_1)$ ,  $s_i = \text{Res}_{A_i}^{\mathbb{E}}(u_2)$ ,  $1 \leq i \leq 3$ , and let  $t_j = \beta s_j$ ,  $0 \leq j \leq 3$ . Let  $u$  be a generator of  $H^1(\langle c \rangle)$  and set  $v = \beta u \in H^2(\langle c \rangle)$ . It follows that

$$H^*(\langle c \rangle) = \Lambda[u] \otimes \mathbb{F}_3[v].$$

So, by Künneth formula, we have

$$H^*(A_i) = \Lambda[s_i, u] \otimes \mathbb{F}_3[t_i, v],$$

$0 \leq i \leq 3$ .

It follows from [4] that  $H^*(\mathbb{E})$  is detected by the  $A_i$ 's. According to [3], we have

**Lemma 1.** (i) *There exist elements  $U_1, U_2 \in H^2(\mathbb{E})$  such that:*

(ia)  *$v_1, v_2, U_1, U_2$  is a basis of  $H^2(\mathbb{E})$ . Furthermore, we have*

$$\begin{aligned} u_1|_{A_0} &= s_0, & u_1|_{A_1} &= s_1, & u_1|_{A_2} &= 2s_2, & u_1|_{A_3} &= 0, \\ v_1|_{A_0} &= t_0, & v_1|_{A_1} &= t_1, & v_1|_{A_2} &= 2t_2, & v_1|_{A_3} &= 0, \\ u_2|_{A_0} &= 0, & u_2|_{A_1} &= s_1, & u_2|_{A_2} &= s_2, & u_2|_{A_3} &= s_3, \\ v_2|_{A_0} &= 0, & v_2|_{A_1} &= t_1, & v_2|_{A_2} &= t_2, & v_2|_{A_3} &= t_3, \\ U_1|_{A_0} &= s_0u, & U_1|_{A_1} &= s_1u + t_1, & U_1|_{A_2} &= 2s_2u + t_2, & U_1|_{A_3} &= 0, \\ U_2|_{A_0} &= 0, & U_2|_{A_1} &= 2s_1u + t_1, & U_2|_{A_2} &= 2s_2u + 2t_2, & U_2|_{A_3} &= s_3u; \end{aligned}$$

(ib)  *$v_1^2, v_1v_2, v_2^2, v_1U_1, v_1U_2, v_2U_1, v_2U_2$  is a basis of  $H^4(\mathbb{E})$  and  $U_1^2 = v_1U_2$ ,  $U_2^2 = v_2U_1$ ,  $U_1U_2 = v_1v_2$ .*

(ii)  *$v_1u_1, v_1u_2$  is linearly independent in  $H^3(\mathbb{E})$ .*

**Lemma 2.** *Let  $X$  be an element of  $H^*(G)$  of degree  $n \leq 2$ . Then  $X = 0$  provided that one of the following conditions is satisfied:*

- (i)  $Xv_1 = 0$ ;
- (ii)  $Xz = 0$ .

*Proof.* The case  $n = 1$  is trivial. For  $n = 2$ , write  $X = a_{2,0} + a_{1,1} + a_{0,2}$  with  $a_{i,j} \in H^i(\mathbb{E}) \otimes H^j(K)$ . If  $Xv_1 = 0$ , then  $a_{2,0}v_1 = a_{1,1}v_1 = a_{0,2}v_1 = 0$ , hence  $a_{0,2} = 0$ , and, by Lemma 1,  $a_{2,0} = a_{1,1} = 0$ ; so  $X = 0$ . If  $Xz = 0$ , then

$$\begin{aligned} a_{2,0}v_1 &= 0, \\ a_{2,0}(u_3u_4 + u_5u_6) + a_{0,2}v_1 &= 0, \\ a_{1,1}v_1 = a_{0,2}(u_3u_4 + u_5u_6) &= a_{1,1}(u_3u_4 + u_5u_6) = 0, \end{aligned}$$

hence, it follows from what we just proved that  $a_{2,0} = a_{1,1} = a_{0,2} = 0$ , so  $X = 0$ . The lemma follows.

**Lemma 3.** *The cup-product with  $v_1^2$  is an injective map from  $H^2(\mathbb{E})$  to  $H^6(\mathbb{E})$ .*

*Proof.* If  $v_1^2(\lambda_1v_1 + \lambda_2v_2 + \mu_1U_1 + \mu_2U_2) = 0$  with  $\lambda_i, \mu_i \in \mathbb{Z}/3$ , then restricting to  $A_j, 0 \leq j \leq 3$  yields  $\lambda_1 = \lambda_2 = \mu_1 = \mu_2 = 0$ . The lemma follows.

For elements  $X, Y, \dots$  of  $H^*(G)$ , denote by  $(X, Y, \dots)$  the ideal of  $H^*(G)$  generated by  $X, Y, \dots$ . We have

**Lemma 4.** *Let  $X$  be an element of  $H^1(G)$  with  $X\beta z \in (z)$ . Then  $X = 0$ .*

*Proof.* Write  $X\beta z = Yz$  with  $Y \in H^2(G)$ . Since  $X$  and  $\beta z$  are free of  $v_1$ , it follows that  $Yv_1 = 0$ . By Lemma 2,  $Y = 0$ , so  $X\beta z = 0$ . Write  $X = \sum_{i=1}^6 \lambda_i u_i$ . A direct verification shows that  $\lambda_1 = \dots = \lambda_6 = 0$ . The lemma follows.

**Lemma 5.**  $v_1^2v_2 \notin (z, \beta z)$ .

*Proof.* Assume that  $v_1^2v_2 \in (z, \beta z)$ . Then there exist elements  $a_{i,j}, b_{i,j} \in H^i(\mathbb{E}) \otimes H^j(K)$  satisfying

$$\begin{aligned} (1) \quad v_1^2v_2 &= \sum a_{i,j}z + \sum b_{i,j}\beta z \\ &= \sum a_{i,j}(v_1 + u_3u_4 + u_5u_6) + \sum b_{i,j}(v_3u_4 - v_4u_3 + v_5u_6 - v_6u_5). \end{aligned}$$

By decomposing (1), we get

$$\begin{aligned} (2) \quad v_1^2v_2 &= a_{4,0}v_1, \\ (3) \quad 0 &= a_{4,0}(u_3u_4 + u_5u_6) + a_{2,2}v_1, \\ (4) \quad 0 &= a_{2,2}(u_3u_4 + u_5u_6) + a_{0,4}v_1 + b_{2,1}(v_3u_4 - v_4u_3 + v_5u_6 - v_6u_5). \end{aligned}$$

It follows from (3) and Lemma 2 that  $a_{4,0}$  contains  $v_1$  as a factor. By Lemma 3,  $a_{4,0} = v_2v_1$ . Hence, by (3) and Lemma 2,  $a_{2,2} = -v_2(u_3u_4 + u_5u_6)$ . So, from (4), we get

$$2v_2u_3u_4u_5u_6 = a_{0,4}v_1 + b_{2,1}(v_3u_4 - v_4u_3 + v_5u_6 - v_6u_5),$$

a contradiction. The lemma follows.

With some abuse of notation, we consider elements of  $H^*(G)$  as elements of  $H^*(\mathcal{G})$  via the inflation map. Set  $\xi = U_1 \in H^2(\mathcal{G})$ . The first part of the theorem follows from

**Lemma 6.**  $\xi$  is nilpotent.

*Proof.* It follows from Lemma 1 that  $\xi^2 = v_1U_2$ ; hence

$$\xi^3 = v_1U_1U_2 = v_1^2v_2.$$

So  $\xi^9 = (v_1^3)^2v_2^3$ . Since, in  $H^*(\mathcal{G})$ ,  $z = 0$ , we have  $v_1^3 = (-u_3u_4 - u_5u_6)^3 = 0$ . It follows that  $\xi^9 = 0$ . Therefore  $\xi$  is nilpotent. The lemma is proved.

The proof of the theorem is completed by the following.

**Lemma 7.**  $\xi^3 \neq 0$  in  $H^*(\mathcal{G})$ .

*Proof.* Assume that  $\zeta = \xi^3 = v_1^2 v_2 = 0$  in  $H^*(\mathcal{G})$ . So  $\zeta \in \text{ImInf}_{\mathcal{G}}^G$ . Set  $Z = i(\mathbb{Z}/3) \subset \mathcal{G}$ . Denote by  $\{E_r, d_r\}$  the Hochschild-Serre spectral sequence corresponding to the extension  $(\mathcal{G})$ . So

$$E_2 = H^*(G) \otimes H^*(Z).$$

Let  $s$  be a generator of  $H^1(Z)$  and set  $t = \beta s \in H^2(Z)$ .  $s$  can be chosen such that  $d_2(s) = z$ .  $t$  is then transgressive and  $d_3(t) = \beta z$ . So, by Lemma 5,  $\zeta \neq 0$  in  $E_4^{6,0}$ . Since  $\zeta \in \text{ImInf}_{\mathcal{G}}^G$ , it must be hit by  $d_4$  or  $d_5$ . If  $\zeta = d_4(a \otimes st)$  with  $a \otimes st \in E_4^{2,3}$ , then

$$\begin{aligned} d_2(a \otimes st) &= (a \otimes t)d_2(s) \\ &= az \otimes t \\ &= 0 \end{aligned}$$

in  $E_2^{4,2}$ ; so  $az = 0$  in  $H^*(G)$ , hence  $a = 0$  by Lemma 2, a contradiction. If  $\zeta = d_5(b \otimes t^2)$  with  $b \otimes t^2 \in E_5^{1,4}$ , then

$$\begin{aligned} d_3(b \otimes t^2) &= (b \otimes 1)d_3(t^2) \\ &= (b \otimes 1)(2\beta z \otimes t) \\ &= 2b\beta z \otimes t \\ &= 0 \end{aligned}$$

in  $E_3^{4,2}$ ; so  $b\beta z \in (z)$ , hence  $b = 0$  by Lemma 4, a contradiction. Therefore  $\zeta \neq 0$ . The lemma follows.

#### REFERENCES

1. G.S. Arvunin, J.F. Carlson, *Nilpotency degree of cohomology rings in characteristic two*, Proc. Amer. Math. Soc. **118** (1993), 339-443. MR **93g**:20098
2. K. Inoue, A. Kono, *Nilpotency of a kernel of the Quillen map*, J. Math. Kyoto Univ. **33** (1993), 1047-1055. MR **95a**:20053
3. I.J. Leary, *The mod- $p$  cohomology rings of some  $p$ -groups*, Math. Proc. Camb. Phil. Soc. **112** (1992), 63-75. MR **94a**:20087
4. R.J. Milgram, M. Tezuka, *The geometry and cohomology of  $M_{12}$ . II*, Bol. Soc. Mat. Mexicana **1** (1995), 91-108. MR **97f**:20064
5. P.A. Minh, *Essential mod- $p$  cohomology classes of  $p$ -groups: an upper bound for the nilpotency degrees*, Bull. London Math. Soc. **32** (2000), 285-291. MR **2001c**:20106

DEPARTMENT OF MATHEMATICS, COLLEGE OF SCIENCES, UNIVERSITY OF HUE, DAI HOC KHOA HOC, HUE, VIETNAM

*E-mail address:* paminh@dong.vnn.vn

*Current address:* 53 Craig Road, Stockport SK4 2AP, England