

THE MAGNUS REPRESENTATION OF THE TORELLI GROUP $\mathcal{I}_{g,1}$ IS NOT FAITHFUL FOR $g \geq 2$

MASAAKI SUZUKI

(Communicated by Ronald A. Fintushel)

ABSTRACT. In this paper we consider the Magnus representation of the Torelli group. We prove that it is not faithful by showing a non-trivial element in the kernel of this representation.

1. INTRODUCTION

Let Σ_g be a closed oriented surface of genus g and let \mathcal{M}_g be its mapping class group. Namely it is the group of path components of $\text{Diff}_+\Sigma_g$ which is the group of orientation preserving diffeomorphisms of Σ_g . We write $\Sigma_{g,1}$ for an oriented surface obtained from Σ_g by removing an open disk D^2 . We denote by $\mathcal{M}_{g,1}$ the mapping class group of $\Sigma_{g,1}$ relative to the boundary, that is, the group of path components of the group of orientation preserving diffeomorphisms of $\Sigma_{g,1}$ which restrict to the identity on the boundary. Let \mathcal{I}_g (resp. $\mathcal{I}_{g,1}$) be the Torelli group of Σ_g (resp. $\Sigma_{g,1}$), namely the normal subgroup of \mathcal{M}_g (resp. $\mathcal{M}_{g,1}$) consisting of all the elements which act on the homology of Σ_g (resp. $\Sigma_{g,1}$) trivially. Johnson obtained fundamental results concerning the structure of \mathcal{I}_g and $\mathcal{I}_{g,1}$ (e.g. [J]).

As is shown in Birman's book [Bi], Fox's free differential calculus [F] can be used to define a number of interesting matrix representations of free groups of finite ranks and also of various subgroups of the automorphism group of a free group. The first such representation was introduced by Magnus so they are called Magnus representations. For example, the classical Burau and the Gassner representation of the Artin braid group can be obtained in this way. The Magnus representation for the mapping class group $\mathcal{M}_{g,1}$,

$$r : \mathcal{M}_{g,1} \rightarrow GL(2g; \mathbf{Z}[\Gamma_0]),$$

has been studied in [Mo1], where $\Gamma_0 = \pi_1(\Sigma_{g,1})$. We restrict this mapping to the Torelli group $\mathcal{I}_{g,1}$ and reduce the coefficients to $\mathbf{Z}[H]$ which is induced by the abelianization $\mathfrak{a} : \Gamma_0 \rightarrow H$, where $H = H_1(\Sigma_{g,1}; \mathbf{Z})$. In this way, we obtain a homomorphism

$$\bar{r} : \mathcal{I}_{g,1} \rightarrow GL(2g; \mathbf{Z}[H]).$$

We call this mapping \bar{r} the Magnus representation of the Torelli group.

Moody has shown in [M] that the Burau representation τ_n of the braid group B_n is not faithful for $n \geq 9$. His techniques were improved by Long and Paton in [LP],

Received by the editors May 16, 2000 and, in revised form, August 30, 2000.
2000 *Mathematics Subject Classification*. Primary 57M05; Secondary 20F34, 57N05.
Key words and phrases. Magnus representation, mapping class group, Torelli group.




FIGURE 1. Generators of Γ_0

who showed that τ_n is not faithful for $n \geq 6$. Moreover, Bigelow recently proved in [B] that τ_5 is not faithful. On the other hand it seems to still be unknown whether the Gassner representation of the pure braid group is faithful or not. Similarly it has been an open problem to determine whether \bar{r} is injective or not (see [Mo3, Problem 15], [Mo4, Problem 6.23]). In this paper we settle this problem by showing a non-trivial element in the kernel of \bar{r} .

2. DEFINITION OF THE MAGNUS REPRESENTATION

In this section, we recall the definition of Magnus representation for the mapping class group $\mathcal{M}_{g,1}$ from [Mo1].

Let $\mathbf{Z}[\Gamma_0]$ be the integral group ring of $\Gamma_0 = \pi_1(\Sigma_{g,1})$. Then the Magnus representation for the mapping class group is a mapping

$$r : \mathcal{M}_{g,1} \rightarrow GL(2g; \mathbf{Z}[\Gamma_0])$$

defined as follows. We fix a system of generators $\alpha_1, \beta_1, \alpha_2, \beta_2, \dots, \alpha_g, \beta_g$ of the free group Γ_0 as shown in Figure 1. Let us simply write $\gamma_1, \dots, \gamma_{2g}$ for them. Then we have the free differential calculus $\frac{\partial}{\partial \gamma_i} : \mathbf{Z}[\Gamma_0] \rightarrow \mathbf{Z}[\Gamma_0]$. Let $\bar{\cdot} : \mathbf{Z}[\Gamma_0] \rightarrow \mathbf{Z}[\Gamma_0]$ be the antiautomorphism induced by the mapping $\gamma \mapsto \gamma^{-1}$.

Definition 2.1. We call the mapping

$$r : \mathcal{M}_{g,1} \longrightarrow GL(2g; \mathbf{Z}[\Gamma_0])$$

$$\varphi \longmapsto \left(\frac{\partial \varphi(\gamma_j)}{\partial \gamma_i} \right)_{i,j}$$

the Magnus representation for the mapping class group $\mathcal{M}_{g,1}$.

This mapping is not a homomorphism in the usual sense but is rather a crossed homomorphism. That is to say,

Proposition 2.2 ([Mo1]). *For any two elements $\varphi, \psi \in \mathcal{M}_{g,1}$, we have*

$$r(\varphi\psi) = r(\varphi) \cdot {}^\varphi r(\psi)$$

where ${}^\varphi r(\psi)$ denotes the matrix obtained from $r(\psi)$ by applying the automorphism $\varphi : \mathbf{Z}[\Gamma_0] \rightarrow \mathbf{Z}[\Gamma_0]$ on each entry.

To obtain a genuine homomorphism, we have to restrict this mapping to the Torelli group $\mathcal{I}_{g,1}$ and reduce the coefficients to $\mathbf{Z}[H]$ which is induced by the abelianization $\mathfrak{a} : \Gamma_0 \rightarrow H$, where $H = H_1(\Sigma_{g,1}; \mathbf{Z})$. Since the Torelli group $\mathcal{I}_{g,1}$ acts trivially on H , we obtain a homomorphism

$$\bar{r} : \mathcal{I}_{g,1} \rightarrow GL(2g; \mathbf{Z}[H]).$$




FIGURE 2. Simple closed curve c_1




FIGURE 3. Simple closed curve c_2




FIGURE 4. Lickorish generators

Thus we have the following commutative diagram:

$$\begin{array}{ccc}
 \mathcal{M}_{g,1} & \xrightarrow{r} & GL(2g; \mathbf{Z}[\Gamma_0]) \\
 \nabla & & \downarrow \alpha \\
 \mathcal{I}_{g,1} & \xrightarrow{\bar{r}} & GL(2g; \mathbf{Z}[H])
 \end{array}$$

3. MAIN THEOREM

We can easily show that the Magnus representation of the Torelli group is faithful for $g = 1$. However, for higher genera, we obtain the following.

Theorem 3.1. *The Magnus representation of the Torelli group $\mathcal{I}_{g,1}$ is not faithful for $g \geq 2$.*

Proof. We consider two elements $\varphi_1, \varphi_2\varphi_1\varphi_2^{-1}$ of $\mathcal{I}_{g,1}$, where φ_1 and φ_2 are the Dehn twists about simple closed curves c_1 and c_2 as depicted in Figure 2 and Figure 3 respectively. We can write $\varphi_1 = (\mu_2\lambda_2)^6$, $\varphi_2 = \delta\mu_2\delta^{-1}$, $\delta = \lambda_2\nu_1\lambda_1\mu_1^2\lambda_1\nu_1\lambda_2$, where λ_i, μ_i, ν_i are the Lickorish generators which are the Dehn twists along simple closed curves l_i, m_i, n_i as shown in Figure 4. We can see the formulas for the action of λ_i, μ_i and ν_i on the free group generators α_i and β_j in [Mo2, Lemma 4.4].

Since

$$\begin{aligned} \varphi_1(\gamma_j) &= \begin{cases} \beta_2\alpha_2\beta_2^{-1}\alpha_2\beta_2\alpha_2^{-1}\beta_2^{-1}, & j = 3, \\ \beta_2\alpha_2\beta_2^{-1}\alpha_2^{-1}\beta_2\alpha_2\beta_2\alpha_2^{-1}\beta_2^{-1}, & j = 4, \\ \gamma_j, & \text{otherwise,} \end{cases} \\ \varphi_2(\gamma_j) &= \begin{cases} \alpha_2\beta_2^{-1}\alpha_2^{-1}\beta_1\alpha_1\beta_1^{-1}\alpha_1\beta_1\alpha_1^{-1}\beta_1^{-1}\alpha_2\beta_2\alpha_2^{-1}, & j = 1, \\ \alpha_2\beta_2^{-1}\alpha_2^{-1}\beta_1\alpha_1\beta_1^{-1}\alpha_1^{-1}\beta_1\alpha_1\beta_1\alpha_1^{-1}\beta_1^{-1}\alpha_2\beta_2\alpha_2^{-1}, & j = 2, \\ \alpha_2\beta_2^{-1}\alpha_2^{-1}\beta_1\alpha_1\beta_1^{-1}\alpha_1^{-1}\alpha_2, & j = 3, \\ \gamma_j, & \text{otherwise,} \end{cases} \\ \varphi_2^{-1}(\gamma_j) &= \begin{cases} \alpha_1\beta_1\alpha_1^{-1}\beta_1^{-1}\alpha_2\beta_2\alpha_2^{-1}\alpha_1\alpha_2\beta_2^{-1}\alpha_2^{-1}\beta_1\alpha_1\beta_1^{-1}\alpha_1^{-1}, & j = 1, \\ \alpha_1\beta_1\alpha_1^{-1}\beta_1^{-1}\alpha_2\beta_2\alpha_2^{-1}\beta_1\alpha_2\beta_2^{-1}\alpha_2^{-1}\beta_1\alpha_1\beta_1^{-1}\alpha_1^{-1}, & j = 2, \\ \alpha_1\beta_1\alpha_1^{-1}\beta_1^{-1}\alpha_2\beta_2, & j = 3, \\ \gamma_j, & \text{otherwise,} \end{cases} \end{aligned}$$

we obtain

$$\begin{aligned} \frac{\partial\varphi_1(\gamma_3)}{\partial\gamma_i} &= \begin{cases} \beta_2 + \beta_2\alpha_2\beta_2^{-1} - \beta_2\alpha_2\beta_2^{-1}\alpha_2\beta_2\alpha_2^{-1}, & i = 3, \\ 1 - \beta_2\alpha_2\beta_2^{-1} + \beta_2\alpha_2\beta_2^{-1}\alpha_2 \\ -\beta_2\alpha_2\beta_2^{-1}\alpha_2\beta_2\alpha_2^{-1}\beta_2^{-1}, & i = 4, \\ 0, & \text{otherwise,} \end{cases} \\ \mathbf{a}\left(\frac{\partial\varphi_1(\gamma_3)}{\partial\gamma_i}\right) &= \begin{cases} y_2 + x_2 - x_2y_2, & i = 3, \\ 1 - x_2 + x_2^2 - x_2, & i = 4, \\ 0, & \text{otherwise.} \end{cases} \end{aligned}$$

Here we denote by x_i, y_i the abelianization of α_i, β_i respectively. By similar calculation, we have $\bar{\mathbf{r}}(\varphi_1) = I_{2g} + M_1$ and $\bar{\mathbf{r}}(\varphi_2\varphi_1\varphi_2^{-1}) = I_{2g} + M_2$, where

$$\begin{aligned} M_1 &= \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 0 & 0 & 0 & \cdots & 0 \\ 0 & 0 & -a_2b_2 & -b_2^2 & 0 & \cdots & 0 \\ 0 & 0 & a_2^2 & a_2b_2 & 0 & \cdots & 0 \\ 0 & 0 & 0 & 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & 0 & \cdots & 0 \end{pmatrix}, \\ M_2 &= \begin{pmatrix} -a_1b_1b_2b'_2 & -b_1^2b_2b'_2 & -a_2b_1b'_2 & -b_1b_2b'_2 & 0 & \cdots & 0 \\ a_1^2b_2b'_2 & a_1b_1b_2b'_2 & a_1a_2b'_2 & a_1b_2b'_2 & 0 & \cdots & 0 \\ -a_1b_2^2 & -b_1b_2^2 & -a_2b_2 & -b_2^2 & 0 & \cdots & 0 \\ a_1a_2b_2 & a_2b_1b_2 & a_2^2 & a_2b_2 & 0 & \cdots & 0 \\ 0 & 0 & 0 & 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & 0 & \cdots & 0 \end{pmatrix}. \end{aligned}$$

Here $a_i = 1 - x_i^{-1}$, $b_i = 1 - y_i^{-1}$, $b'_2 = 1 - y_2$. Because of the fact that $M_1M_2 = M_2M_1$ is the zero matrix, we have

$$\bar{\mathbf{r}}(\varphi_1)\bar{\mathbf{r}}(\varphi_2\varphi_1\varphi_2^{-1}) = \bar{\mathbf{r}}(\varphi_2\varphi_1\varphi_2^{-1})\bar{\mathbf{r}}(\varphi_1).$$




FIGURE 5. Simple closed curve c_3




FIGURE 6. Simple closed curve c_4

It follows that

$$\bar{r}([\varphi_1, \varphi_2 \varphi_1 \varphi_2^{-1}]) = I_{2g}.$$

It is easy to see that $[\varphi_1, \varphi_2 \varphi_1 \varphi_2^{-1}]$ is a non-trivial element. Hence \bar{r} is not injective and this completes the proof. \square

Remark 3.2. We set $\psi = [\varphi_1, \varphi_2 \varphi_1 \varphi_2^{-1}]$. Then ψ can be expressed as a word of length 116 in the Lickorish generators. The words $\psi(\gamma_i)$ ($1 \leq i \leq 4$) are so large that we cannot write them down here. However, it can be written as

$$\psi(\gamma_i) = l_i \gamma_i l'_i$$

for some $l_i, l'_i \in [[\Gamma_0, \Gamma_0], [\Gamma_0, \Gamma_0]]$. This equation suggests $\psi \in \text{Ker } \bar{r}$ (cf. [Bi, Theorem 3.5, Theorem 3.16]).

Remark 3.3. Here is another element in the kernel of \bar{r} . Let φ' and φ'' be the Dehn twists about simple closed curves c_3 and c_4 as depicted in Figure 5 and Figure 6 respectively. Here we can take i, j, k arbitrarily. Again explicit computation shows that

$$\bar{r}(\varphi') \bar{r}(\varphi'') = \bar{r}(\varphi'') \bar{r}(\varphi').$$

Hence $[\varphi', \varphi'']$ is also an element in the kernel of \bar{r} .

ACKNOWLEDGEMENTS

The author would like to express his gratitude to Prof. Shigeyuki Morita for encouragement and helpful suggestions and to thank Prof. Takayuki Morifuji for valuable discussions and advice. He also would like to thank Mrs. Miyamoto.

REFERENCES

[B] S. Bigelow *The Burau representation is not faithful for $n = 5$* , *Geom. Topol.* **3** (1999), 397–404 CMP 2000:05
 [Bi] J. Birman, *Braids, Links and Mapping Class Groups*, *Ann. of Math. Stud. No. 82*, Princeton University Press, 1975

- [F] R. H. Fox, *Free Differential Calculus I*, Ann. of Math. **57** (1953), 547–560 MR **14**:843d
- [J] D. Johnson, *A survey of the Torelli group*, Contemp. Math. **20** (1983), 165–179 MR **85d**:57009
- [LP] D. D. Long, M. Paton, *The Burau representation is not faithful for $n \geq 6$* , Topology **32** (1993), 439–447 MR **94c**:20071
- [M] J. Moody, *The Burau representation is unfaithful for large n* , Bull. Amer. Math. Soc. **25** (1991), 379–384 MR **92b**:20041
- [Mo1] S. Morita, *Abelian quotients of subgroups of the mapping class group of surfaces*, Duke Math. J. **70** (1993), 197–224 MR **97d**:57003
- [Mo2] S. Morita, *The extension of Johnson’s homomorphism from the Torelli group to the mapping class group*, Invent. Math. **111** (1993), no. 1, 197–224 MR **93j**:57001
- [Mo3] S. Morita, *Problems on the structure of the mapping class group of surfaces and topology of the moduli space of curves*, “Proceedings of the Taniguchi Symposium on Topology, Geometry and Field Theory”, held in Sanda, January 1993, World Scientific (1994) 101–110 MR **95m**:57028
- [Mo4] S. Morita, *Structure of the mapping class groups of surfaces: a survey and a prospect*, Geom. Topol. Monogr. Vol.2: Proceedings of the Kirbyfest (1999) 349–406 MR **2000j**:57039

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF TOKYO, 3-8-1 KOMABA,
MEGURO, TOKYO 153-8914, JAPAN

E-mail address: macky@ms.u-tokyo.ac.jp