

## ON PERFECTLY MEAGER SETS

TOMEK BARTOSZYNSKI

(Communicated by Alan Dow)

ABSTRACT. We show that it is consistent that the product of perfectly meager sets is perfectly meager.

### 1. INTRODUCTION

Suppose that  $X$  is a subset of a Polish space  $\mathbf{X}$ . We say that  $X$  is *perfectly meager* if for every perfect set  $P \subseteq \mathbf{X}$ ,  $X \cap P$  is meager in the relative topology of  $P$ . Let  $\mathbf{PM}$  denote the collection of perfectly meager sets.

Clearly all countable sets are perfectly meager but there are various examples of uncountable perfectly meager sets that can be constructed in  $\mathbf{ZFC}$  (see [5]).

In [4], Marczewski asked whether the product of perfectly meager sets is perfectly meager. This question was partially answered by Reclaw who showed that:

**Theorem 1** (Reclaw [7]). *Assume CH. Then there are two perfectly meager sets whose product is not perfectly meager.*

The proof relies on the existence of a Borel set having certain properties and the existence of a Luzin set (i.e. an uncountable set whose intersection with every meager set is countable).

The purpose of this note is to show that it is also consistent with  $\mathbf{ZFC}$  that the product of any two perfectly meager sets is perfectly meager. Thus, Marczewski's question is undecidable in  $\mathbf{ZFC}$ .

**Definition 2.** Suppose that  $X$  is a subset of a Polish space  $\mathbf{X}$ . We say that  $X$  is *universally meager* if every Borel isomorphic image of  $X$  in  $\mathbf{X}$  is meager. Let  $\mathbf{UM}$  denote the collection of universally meager sets.

It is clear that  $\mathbf{UM} \subseteq \mathbf{PM}$ , but the other inclusion may fail.

**Theorem 3** (Sierpiński [8]). *Assume CH. Then  $\mathbf{PM} \neq \mathbf{UM}$ .*

Unlike  $\mathbf{PM}$ , the class of universally meager sets is closed under products.

**Theorem 4** (Zakrzewski [10]). *The product of universally meager sets is universally meager.*

This is a consequence of the following characterization of the class  $\mathbf{UM}$ . Let  $\mathbb{C}$  denote the Cohen algebra.

---

Received by the editors May 5, 2000 and, in revised form, October 24, 2000.

1991 *Mathematics Subject Classification.* Primary 03E17.

*Key words and phrases.* Perfectly meager, products, consistency.

The author was partially supported by NSF grant DMS 99271.

**Theorem 5** (Zakrzewski [10]). *For a subset  $X$  of a perfect Polish space  $\mathbf{X}$ , the following are equivalent:*

- (1)  $X \in \mathbf{UM}$ .
- (2)  $X$  does not contain a Borel one-to-one image of a non-meager set.
- (3) For every  $\sigma$ -ideal  $\mathcal{J}$  in  $\mathbf{Borel}(\mathbf{X})$  such that  $\mathbf{Borel}(\mathbf{X})/\mathcal{J} \cong \mathbb{C}$  there is a Borel set  $B \in \mathcal{J}$  such that  $X \subseteq B$ .
- (4)  $X$  is meager in every Polish topology  $\tau$  on  $\mathbf{X}$  such that  $\mathbf{X}$  has no isolated points and  $\mathbf{Borel}(\mathbf{X}, \tau) = \mathbf{Borel}(\mathbf{X})$ .
- (5)  $X$  is meager in every second countable Hausdorff topology  $\tau$  on  $\mathbf{X}$  such that  $\mathbf{X}$  has no isolated points and all Borel sets (in the original Polish topology) have Baire Property in the topology  $\tau$ .
- (6) There is no  $\sigma$ -ideal  $\mathcal{J}$  in  $\mathbf{Borel}(X)$  such that  $\mathbf{Borel}(X)/\mathcal{J} \cong \mathbb{C}$ .
- (7)  $X$  is meager in every second countable Hausdorff topology  $\tau$  on  $X$  such that  $X$  has no isolated points and all Borel subsets of  $X$  (in the topology inherited from the original Polish topology on  $\mathbf{X}$ ) have Baire Property in the topology  $\tau$ .
- (8)  $X$  is meager in every separable metrizable topology  $\tau$  on  $X$  such that  $X$  has no isolated points and  $\mathbf{Borel}(X) = \mathbf{Borel}(X, \tau)$ .

Now we can formulate our main result:

**Theorem 6.** *It is consistent that  $\mathbf{PM} = \mathbf{UM}$ . In particular, it is consistent that  $\mathbf{PM}$  is closed under products.*

## 2. IN ZFC

In this section we will identify a more general property that implies that  $\mathbf{PM} = \mathbf{UM}$ .

For a function  $f : X \rightarrow Y$  let  $f[X] = \{f(x) : x \in X\}$  denote the image of  $X$ .

We need the following observation:

**Lemma 7.** *Suppose that  $X \notin \mathbf{UM}$ . Then there exists  $X' \subseteq X$ , a Borel set  $A \supseteq X'$  and Borel isomorphism  $f : A \rightarrow \omega^\omega$  such that:*

- (1)  $f^{-1}$  is continuous,
- (2)  $f[X']$  is not meager in  $\omega^\omega$ .

*Proof.* Assume that  $X$  is a subset of a Polish space  $\mathbf{X}$ . Let  $h : X \rightarrow h[X]$  be a Borel isomorphism witnessing that  $X \notin \mathbf{UM}$ . Find Borel sets  $A', B \subseteq \mathbf{X}$ ,  $A' \supseteq X$  and  $B \supseteq h[X]$  and the Borel isomorphism  $\bar{h} : A' \rightarrow B$  extending  $h$  (exercise in [9]).

Apply Kuratowski's theorem ([3], 8.38) to get a dense  $G_\delta$  subset  $B' \subseteq B$  such that  $\bar{h}^{-1} \upharpoonright B'$  is continuous. Since  $B'$  is a nonmeager Borel set it contains a relative  $G_\delta$  dense subset  $B''$  which is homeomorphic to  $\omega^\omega$  via a homeomorphism  $g$ . Finally set  $A = \bar{h}^{-1}(B'')$  and  $f = g \circ \bar{h}$ , and notice that this works since  $\bar{h}[X] \cap B''$  is not meager in  $B''$ .  $\square$

Consider the following principle:

**Definition 8.** Axiom P: For every nonmeager subset  $X \subseteq \omega^\omega$  there exists a compact subset  $P \subseteq \omega^\omega$  such that  $P \cap X$  is nonmeager in  $P$ .

**Theorem 9.** *Assume P. Then  $\mathbf{UM} = \mathbf{PM}$ .*

*Proof.* Suppose that  $X \notin \mathbf{UM}$ . By Lemma 7 there exists  $X' \subseteq X$ , a Borel set  $A \supseteq X'$  and Borel isomorphism  $f : A \rightarrow \omega^\omega$  such that:

- (1)  $f^{-1}$  is continuous,
- (2)  $f[X']$  is not meager in  $\omega^\omega$ .

Apply Axiom P to to find a compact set  $P \subseteq \omega^\omega$  such that  $P \cap f[X']$  is not meager in  $P$ . Set  $Q = f^{-1}(P)$  and note that  $f|_Q$  is a homeomorphism between  $P$  and  $Q$ . Under this homeomorphism  $Q \cap X'$  is the image of  $f[X'] \cap P$ , thus it is not meager in  $Q$ . Since  $Q \cap X \supseteq Q \cap X'$  it follows that  $X \notin \mathbf{PM}$ . As the other inclusion is obvious, the theorem follows.  $\square$

### 3. FORCING

In this section we will show that Axiom P is consistent with ZFC, which will finish the proof.

For a tree  $p$  and  $t \in p$ , let  $\text{succ}_p(t)$  be the set of all immediate successors of  $t$  in  $p$ ,  $p_t = \{v \in p : t \subseteq v \text{ or } v \subseteq t\}$  the subtree of  $p$  determined by  $t$ , and let  $[p]$  be the set of branches of  $p$ . By identifying  $s \in \omega^{<\omega}$  with the full-branching tree having root  $s$ , we can also denote  $[s] = \{f \in \omega^\omega : s \subseteq f\}$ . Let  $\omega^{\uparrow\omega} = \{s \in \omega^{<\omega} : s \text{ is strictly increasing}\}$ .

The rational perfect forcing  $\mathbb{M}$  is the following forcing notion:

$$p \in \mathbb{M} \iff p \subseteq \omega^{\uparrow\omega} \text{ is a perfect tree \& } \forall s \in p \exists t \in p (s \subseteq t \ \& \ |\text{succ}_p(t)| = \aleph_0).$$

For  $p, q \in \mathbb{M}$ ,  $p \geq q$  if  $p \subseteq q$ . Without loss of generality we can assume that  $|\text{succ}_p(s)| = 1$  or  $|\text{succ}_p(s)| = \aleph_0$  for all  $p \in \mathbb{M}$  and  $s \in p$ . Conditions of this type form a dense subset of  $\mathbb{M}$ .

Let

$$\text{split}(p) = \{s \in p : |\text{succ}_p(s)| > 1\} = \bigcup_{n \in \omega} \text{split}_n(p),$$

where  $\text{split}_n(p) = \{s \in \text{split}(p) : |\{t \subsetneq s : t \in \text{split}(p)\}| = n\}$ .

For  $p, q \in \mathbb{M}$ ,  $n \in \omega$ , we let

$$p \geq_n q \iff p \geq q \ \& \ \text{split}_n(q) = \text{split}_n(p).$$

If  $v \in \text{split}(p)$  let  $U_v^p = \{n \in \omega : v \hat{\ } n \in p\}$  and for  $n \in U_v^p$  let  $v^n$  be the first splitting node below  $v \hat{\ } n$ .

If  $G \subseteq \mathbb{M}$  is a generic filter over  $\mathbf{V}$  let  $\mathbf{m} = \bigcap_{p \in G} [p]$  be the generic real.

Let  $\mathbb{M}_{\omega_2}$  be the countable support iteration of  $\mathbb{M}$  of length  $\aleph_2$ .

The following facts about  $\mathbb{M}$  are well-known.

**Theorem 10.** (1) *The sequence  $\langle \leq_n : n \in \omega \rangle$  witnesses that  $\mathbb{M}$  satisfies axiom A. In particular,  $\mathbb{M}$  is proper.*

(2)  *$\mathbb{M}$  preserves non-meager sets, i.e. if  $A \subseteq 2^\omega$ , where  $A \in \mathbf{V}$  is not meager, then  $\mathbf{V}^{\mathbb{M}} \models A$  is not meager ([1], Theorem 7.3.46).*

(3)  *$\mathbb{M}$  also satisfies the iterable condition for preserving non-meager sets. In particular, countable support iteration of Miller forcing preserves non-meager sets ([1], Theorems 6.3.19 and 6.3.20).*  $\square$

**Theorem 11.**  $\mathbf{V}^{\mathbb{M}_{\omega_2}} \models \text{Axiom } P$ .

The idea of the proof is as follows. Suppose that  $X \in \mathbf{V}^{\mathbb{M}_{\omega_2}}$  and  $X$  is not meager in  $\omega^\omega$ . First we find  $\alpha < \omega_2$  such that  $\mathbf{V}^{\mathbb{M}_\alpha} \models X \cap \mathbf{V}^{\mathbb{M}_\alpha}$  is not meager (Lemma 12). Next we will find a compact set  $P \subseteq \omega^\omega$  belonging to  $\mathbf{V}^{\mathbb{M}_{\alpha+1}}$  such that

$$\mathbf{V}^{\mathbb{M}_{\alpha+1}} \models P \cap X \cap \mathbf{V}^{\mathbb{M}_\alpha} \text{ is not meager in } P \text{ (Theorem 13).}$$

Finally, by 10(3),  $\mathbb{M}_{\omega_2}$  preserves non-meager sets. Thus

$$\mathbf{V}^{\mathbb{M}_{\omega_2}} \models X \cap \mathbf{V}^{\mathbb{M}_\alpha} \text{ is not meager in } P,$$

which implies that  $\mathbf{V}^{\mathbb{M}_{\omega_2}} \models X$  is not meager in  $P$ .

**Lemma 12.** *Suppose that  $X \in \mathbf{V}^{\mathbb{M}_{\omega_2}}$ ,  $X \subseteq \omega^\omega$  and  $X$  is not meager in  $\mathbf{V}^{\mathbb{M}_{\omega_2}}$ . Then there exists  $\alpha < \omega_2$  such that  $\mathbf{V}^{\mathbb{M}_\alpha} \models X \cap \mathbf{V}^{\mathbb{M}_\alpha}$  is not meager.*

*Proof.* Let  $\langle \alpha_\xi : \xi < \omega_1 \rangle$  be a continuous increasing sequence such that

$$\mathbf{V}^{\mathbb{M}_{\omega_2}} \models X \cap \mathbf{V}^{\mathbb{M}_{\alpha_\xi+1}} \text{ is not covered by any meager set from } \mathbf{V}^{\mathbb{M}_{\alpha_\xi}}.$$

By properness,  $\alpha = \sup_{\xi < \omega_1} \alpha_\xi$  has the required property, because every real in  $\mathbf{V}^{\mathbb{M}_\alpha}$  is in some  $\mathbf{V}^{\mathbb{M}_{\alpha_\xi}}$ . □

**Theorem 13.** *Suppose that  $X \in \mathbf{V}$ ,  $X \subseteq \omega^\omega$  is a non-meager set. There is a compact set  $P \subseteq \omega^\omega$ ,  $P \in \mathbf{V}^{\mathbb{M}}$  such that :  $\mathbf{V}^{\mathbb{M}} \models X$  is not meager in  $P$ .*

*Proof.* For the sake of clarity we will break the proof into three lemmas. The main idea of the proof is already present in [2].

Let  $\dot{\mathbf{m}}$  be the canonical name for an  $\mathbb{M}$ -generic real and let  $\mathbb{C}$  be the Cohen forcing represented as  $\omega^{<\omega}$  with  $\dot{\mathbf{c}}$  being the canonical name for the Cohen real.

For  $x \in \omega^\omega$  let  $P_x = \{z \in \omega^\omega : \forall n z(n) \leq x(n)\}$ . Note that  $P_x$  is a compact set in  $\omega^\omega$ .

For two sequences  $s \in \omega^{<\omega}$ ,  $t \in \omega^{\uparrow\omega}$  we say that  $(s, t)$  is *good* if  $|s| = |t|$  and  $s(i) \leq t(i)$  for  $i < |t|$ .

**Lemma 14.** *Suppose that  $p \in \mathbb{M}$ ,  $s \in \mathbb{C}$ , and  $(s \restriction |\text{stem}(p)|, \text{stem}(p))$  is good. Then there is a  $\mathbb{C}$ -name  $\dot{q}$  for an element of  $\mathbb{M}$  such that:*

- (1)  $s \Vdash_{\mathbb{C}} \dot{q} \geq_0 p$ ,
- (2)  $(s, \dot{q}) \Vdash_{\mathbb{C} * \mathbb{M}} \dot{\mathbf{c}} \in P_{\dot{\mathbf{m}}}$ .

*Proof.* Let  $\mathbf{c} \supseteq s$  be a Cohen real over  $\mathbf{V}$ . Working in  $\mathbf{V}[\mathbf{c}]$  define

$$q = \{v \in p : (\mathbf{c} \restriction |v|, v) \text{ is good}\}.$$

It is enough to check that  $q \in \mathbb{M}^{\mathbf{V}[\mathbf{c}]}$ . In fact, we will show that if  $v \in \text{split}(p)$  and  $v \in q$ , then  $v \in \text{split}(q)$ .

Suppose that  $v \in \text{split}(p)$  and  $v \in q$ . In particular,  $(\mathbf{c} \restriction |v|, v)$  is good. Let  $s' = \mathbf{c} \restriction |v|$ . For  $k \in \omega$  let

$$D_k = \{t \in \mathbb{C} : s' \subseteq t \ \& \ \exists n > k \ (t \restriction |v^n|, v^n) \text{ is good}\}.$$

We show that  $D_k$  is dense in  $\mathbb{C}$  below  $s'$ . Take any  $s'' \geq s'$  and let  $n \in U_v^p \setminus \max(\text{range}(s''), k)$ . If  $|s''| > |v^n|$ , then put  $t = s''$ . Otherwise, let  $t \geq s''$  be such that  $|t| = |v^n|$  and

$$t(i) = \begin{cases} s''(i) & \text{if } i < |s''|, \\ 0 & \text{if } |s''| \leq i < |v^n|. \end{cases}$$

It is clear that  $t \in D_k$ . By genericity, we conclude that the set

$$\{n : (\mathbf{c} \upharpoonright |v^n|, v^n) \text{ is good}\}$$

is infinite in  $\mathbf{V}[\mathbf{c}]$ . In particular,  $v \in \text{split}(q)$ . Since  $\mathbf{c}$  was arbitrary, it finishes the proof.  $\square$

**Lemma 15.** *Suppose that  $p \in \mathbb{M}$ ,  $s \in \mathbb{C}$ , and  $(s \upharpoonright |\text{stem}(p)|, \text{stem}(p))$  is good. Let  $\dot{F}$  be an  $\mathbb{M}$ -name for a closed nowhere dense subset of  $P_{\dot{\mathbf{m}}}$ . There exists a  $\mathbb{C}$ -name  $\dot{q}$  for an element of  $\mathbb{M}$  such that:*

- (1)  $s \Vdash_{\mathbb{C}} \dot{q} \geq_0 p$ ,
- (2)  $(s, \dot{q}) \Vdash_{\mathbb{C} * \mathbb{M}} \dot{\mathbf{c}} \in P_{\dot{\mathbf{m}}}$ ,
- (3)  $(s, \dot{q}) \Vdash_{\mathbb{C} * \mathbb{M}} \dot{\mathbf{c}} \notin \dot{F}$ .

*Proof.* Let  $\mathbf{m} \in [p]$  be an  $\mathbb{M}$ -generic real over  $\mathbf{V}$ , and let  $F$  be the interpretation of  $\dot{F}$  using  $\mathbf{m}$ . In  $\mathbf{V}[\mathbf{m}]$  define sequences  $\langle s_n : n \in \omega \rangle \in (\omega^{<\omega})^\omega$  such that:

- (1)  $\forall v \in \prod_{j < n} (\mathbf{m}(j) + 1) [v \frown s_n] \cap F = \emptyset$ ,
- (2)  $s_n \in \prod_{j=n}^k (\mathbf{m}(j) + 1)$  for some  $k > n$ .

Since  $F$  is nowhere dense this definition is correct. Going back to  $\mathbf{V}$  we conclude that there is an  $\mathbb{M}$ -name  $\langle \dot{s}_n : n \in \omega \rangle$  such that:

- (1)  $p \Vdash_{\mathbb{M}} \forall v \in \prod_{j < n} (\dot{\mathbf{m}}(j) + 1) [v \frown \dot{s}_n] \cap \dot{F} = \emptyset$ ,
- (2)  $p \Vdash_{\mathbb{M}} \forall n (\dot{s}_n, \dot{\mathbf{m}} \upharpoonright [n, n + |\dot{s}_n|])$  is good.

For each  $n \in U_{\text{stem}(p)}^p$  find a condition  $p_n \geq p$  such that:

- (1) there is a sequence  $s_n$  such that  $p_n \Vdash_{\mathbb{M}} \dot{s}_n = s_n$ ,
- (2)  $\text{stem}(p_n) \supseteq \text{stem}(p)^n$ ,
- (3)  $|\text{stem}(p_n)| \geq n + |s_n|$ .

Observe that by the choice of  $\langle \dot{s}_n : n \in \omega \rangle$  it follows that

$$(s_n, \text{stem}(p_n) \upharpoonright [n, n + |s_n|]) \text{ is good.}$$

Let  $\mathbf{c} \supseteq s$  be a Cohen real over  $\mathbf{V}$ . Working in  $\mathbf{V}[\mathbf{c}]$  define

$$A = \{n \in \omega : (\mathbf{c} \upharpoonright |\text{stem}(p_n)|, \text{stem}(p_n)) \text{ is good and } \mathbf{c} \upharpoonright [n, n + |s_n|] = s_n\}.$$

We will show that  $A$  is an infinite set in  $\mathbf{V}[\mathbf{c}]$ .

For  $k \in \omega$  let

$$D_k = \left\{ t \in \mathbb{C} : s \subseteq t \ \& \ \exists n > k \left( (t \upharpoonright |\text{stem}(p_n)|, \text{stem}(p_n)) \text{ is good} \ \& \right. \\ \left. t \upharpoonright [n, n + |s_n|] = s_n \right\}.$$

We show that  $D_k$  is dense in  $\mathbb{C}$  below  $s$ . Suppose that  $s' \geq s$  and let  $\ell = \max(\text{range}(s'), |s'|, k)$ . Pick  $n \in U_{\text{stem}(p)}^p \setminus \ell$  and define  $t \geq s'$  such that  $|t| = |\text{stem}(p_n)|$  and

$$t(i) = \begin{cases} s'(i) & \text{if } i < |s'|, \\ 0 & \text{if } |s'| \leq i < n, \\ s_n(i) & \text{if } n \leq i < n + |s_n|, \\ 0 & \text{if } n + |s_n| \leq i < |\text{stem}(p_n)|. \end{cases}$$

Note that by the properties of  $\langle \dot{s}_n : n \in \omega \rangle$  it follows that  $t \in D_k$ . By genericity, for every  $k \in \omega$  there is  $n \geq k$  such that  $\mathbf{c} \upharpoonright n \in D_k$ , which implies that  $A$  is infinite.

Let  $p^* = \bigcup_{n \in U_{\text{stem}(p)}^p} p_n$ . Define in  $\mathbf{V}[\mathbf{c}]$ ,

$$q_1 = \bigcup_{n \in A} p_n \quad \text{and} \quad q_2 = \{v \in p^* : (\mathbf{c} \upharpoonright |v|, v) \text{ is good}\},$$

and let  $q = q_1 \cap q_2$ . Since the nodes corresponding to the elements of  $A$  were good, by Lemma 14, it follows that  $q \in \mathbb{M}^{\mathbf{V}[\mathbf{c}]}$  and  $q \geq_0 p$ . In addition, in  $\mathbf{V}[\mathbf{c}]$ ,  $q \Vdash_{\mathbb{M}} \forall n \mathbf{c}(n) \leq \dot{\mathbf{m}}(n)$  (by the choice of  $q_2$ ) and  $q \Vdash_{\mathbb{M}} \mathbf{c} \notin \dot{F}$  (by the choice of  $q_1$ ). Since  $\mathbf{c}$  was arbitrary, the proof is finished.  $\square$

Finally we show:

**Lemma 16.** *Suppose that  $p \in \mathbb{M}$ ,  $s \in \mathbb{C}$ , and  $(s \upharpoonright |\text{stem}(p)|, \text{stem}(p))$  is good. Let  $\langle \dot{F}_n : n \in \omega \rangle$  be an  $\mathbb{M}$ -name for a sequence of closed nowhere dense subsets of  $P_{\dot{\mathbf{m}}}$ . There exists a  $\mathbb{C}$ -name  $\dot{q}$  for an element of  $\mathbb{M}$  such that:*

- (1)  $s \Vdash_{\mathbb{C}} \dot{q} \geq p$ ,
- (2)  $(s, \dot{q}) \Vdash_{\mathbb{C} * \mathbb{M}} \dot{\mathbf{c}} \in P_{\dot{\mathbf{m}}}$ ,
- (3)  $(s, \dot{q}) \Vdash_{\mathbb{C} * \mathbb{M}} \dot{\mathbf{c}} \notin \bigcup_n \dot{F}_n$ .

*Proof.* The proof is a refinement of the proof of the previous lemma. Suppose that  $\langle \dot{F}_n : n \in \omega \rangle$  is an  $\mathbb{M}$ -name for a sequence of closed nowhere dense subsets of  $P_{\dot{\mathbf{m}}}$ . Without loss of generality we can assume that  $\Vdash_{\mathbb{M}} \forall n \dot{F}_n \subseteq \dot{F}_{n+1}$ . Find an  $\mathbb{M}$ -name  $\langle \dot{s}_n : n \in \omega \rangle$  such that:

- (1)  $p \Vdash_{\mathbb{M}} \forall v \in \prod_{j < n} (\dot{\mathbf{m}}(j) + 1) [v \frown \dot{s}_n] \cap \dot{F}_n = \emptyset$ ,
- (2)  $p \Vdash_{\mathbb{M}} \forall n (\dot{s}_n, \dot{\mathbf{m}} \upharpoonright [n, n + |\dot{s}_n|])$  is good.

Build by induction a sequence of conditions  $\langle p_n : n \in \omega \rangle$  such that:

- (1)  $p_0 = p$ ,
- (2)  $p_{n+1} \geq_n p_n$ ,
- (3) if  $v \in \text{split}_n(p_{n+1})$  and  $k \in U_v^{p_{n+1}}$ , then there exists a sequence  $s_{v,k} \in \omega^{<\omega}$  such that:
  - (a)  $(p_{n+1})_{v^k} \Vdash_{\mathbb{M}} \dot{s}_k = s_{v,k}$ ,
  - (b)  $|v^k| \geq k + |s_{v,k}|$ .

As in the previous lemma, it follows that for  $v \in \text{split}_n(p_{n+1})$  and  $k \in U_v^{p_{n+1}}$ ,  $(s_{v,k}, v^k \upharpoonright [k, k + |s_{v,k}|])$  is good.

The construction is straightforward; the first step is essentially described in the previous lemma. Let  $p^* = \bigcap_n p_n$  and let  $\mathbf{c} \supseteq s$  be a Cohen real over  $\mathbf{V}$ . Working in  $\mathbf{V}[\mathbf{c}]$  define for each  $n \in \omega$  and  $v \in \text{split}_n(p^*)$ :

$$A^v = \left\{ k \in U_v^{p^*} \setminus n : (\mathbf{c} \upharpoonright |v^k|, v^k) \text{ is good and } \mathbf{c} \upharpoonright [k, k + |s_{v,k}|] = s_{v,k} \right\}.$$

As before, it follows that  $A^v$  is infinite in  $\mathbf{V}[\mathbf{c}]$  for every  $v \in \text{split}(p^*)$ . Finally, let  $q \geq p^* \geq p$  be defined so that for every  $v \in \text{split}(q)$ ,  $U_v^q = A^v$ .

It follows from the definition of  $q$  that  $\mathbf{V}[\mathbf{c}] \models q \Vdash_{\mathbb{M}} \mathbf{c} \in P_{\dot{\mathbf{m}}}$ . On the other hand, for every  $v \in \text{split}_n(q)$ ,  $\mathbf{V}[\mathbf{c}] \models q_v \Vdash_{\mathbb{M}} \exists k > n \mathbf{c} \notin \dot{F}_k$ . Thus

$$\mathbf{V}[\mathbf{c}] \models q \Vdash_{\mathbb{M}} \exists^\infty n \mathbf{c} \notin \dot{F}_n.$$

Since the sets  $\dot{F}_n$  are increasing, we conclude that  $\mathbf{V}[\mathbf{c}] \models q \Vdash_{\mathbb{M}} \mathbf{c} \notin \bigcup_n \dot{F}_n$ .  $\square$

Now we are ready to prove Theorem 13. Suppose that  $X \in \mathbf{V}$ ,  $X \subseteq \omega^\omega$  is not meager. We will show that  $\Vdash_{\mathbb{M}} X \cap P_{\dot{\mathbf{m}}}$  is not meager in  $P_{\dot{\mathbf{m}}}$ . Suppose otherwise

and let  $\langle \dot{F}_n : n \in \omega \rangle$  be an  $\mathbb{M}$ -name for a sequence of closed nowhere dense sets in  $P_{\dot{\mathbb{m}}}$  such that for some  $p \in \mathbb{M}$ ,

$$p \Vdash_{\mathbb{M}} X \cap P_{\dot{\mathbb{m}}} \subseteq \bigcup_{n \in \omega} \dot{F}_n.$$

Let  $N \prec \mathbf{H}(\chi)$  be a countable elementary submodel containing  $p$ ,  $X$ ,  $\langle \dot{F}_n : n \in \omega \rangle$ , etc. Since  $X$  is not meager there exists a real  $\mathbf{c} \in X$  which is Cohen over  $N$ . By Lemma 16 there exists a condition  $q \geq p$ ,  $q \in N[\mathbf{c}]$  such that

$$q \Vdash_{\mathbb{M}} \mathbf{c} \in P_{\dot{\mathbb{m}}} \setminus \bigcup_{n \in \omega} \dot{F}_n.$$

This contradicts the choice of  $p$  and finishes the proof.  $\square$

Note that in fact we have showed the following:

**Theorem 17.** *Suppose that  $\langle \mathcal{P}_\alpha, \dot{Q}_\alpha : \alpha < \omega_2 \rangle$  is a countable support iteration of proper forcing notions such that:*

- (1)  $\{\alpha < \omega_2 : \Vdash_\alpha \dot{Q}_\alpha \simeq \mathbb{M}\}$  is cofinal in  $\aleph_2$ ,
- (2)  $\Vdash_\alpha \dot{Q}_\alpha$  preserves non-meager sets for  $\alpha < \omega_2$ .

Then  $\mathbf{V}^{\mathcal{P}_{\omega_2}} \models$  Axiom P.

#### ACKNOWLEDGEMENTS

The work was done while I was spending a sabbatical year at Rutgers University and the College of Staten Island, CUNY, and I thank their mathematics departments for the support. I am also grateful to Andrzej Rosłanowski for carefully proofreading earlier versions of this paper and giving many valuable suggestions.

#### REFERENCES

- [1] Tomek Bartoszyński and Haim Judah. *Set Theory: on the structure of the real line*. A.K. Peters, 1995. MR **96k**:03002
- [2] Haim Judah and Saharon Shelah. Killing Luzin and Sierpiński sets. *Proceedings of the American Mathematical Society*, 120:917–920, 1994. MR **94e**:03046
- [3] Alexander Kechris. *Classical Descriptive Set Theory*, volume 156 of *Graduate Texts in Mathematics*. Springer Verlag, 1995. MR **96e**:03057
- [4] Edward Marczewski (Szpilrajn). Probleme 68. *Fundamenta Mathematicae*, 25:579, 1935.
- [5] Arnold W. Miller. Special subsets of the real line. In K. Kunen and J. E. Vaughan, editors, *Handbook of Set Theoretic Topology*, pages 201–235. North-Holland, Amsterdam, 1984. MR **86i**:54037
- [6] Janusz Pawlikowski. Products of perfectly meager sets and Luzin’s function. *Proceedings of the American Mathematical Society*, 107(3):811–815, 1989. MR **90b**:54025
- [7] Ireneusz Reclaw. Products of perfectly meagre sets. *Proceedings of the American Mathematical Society*, 112(4):1029–1031, 1991. MR **91j**:28001
- [8] Waclaw Sierpiński. Sur une probleme de M.Kuratowski concernant la propriete de Baire des ensembles. *Fundamenta Mathematicae*, 22:262–266, 1934.
- [9] S.M. Srivastava. *A Course on Borel sets*. Graduate Texts in Mathematics. Springer-Verlag, 1998. MR **99d**:04002
- [10] Piotr Zakrzewski. Universally meager sets. to appear in *Proceedings of the American Mathematical Society*.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, BOISE STATE UNIVERSITY, BOISE, IDAHO 83725

*E-mail address:* `tomek@math.boisestate.edu`

*URL:* `http://math.boisestate.edu/~tomek`