

JOINT SPECTRUM OF SUBNORMAL n -TUPLES OF COMPOSITION OPERATORS

JOSÉ GIMÉNEZ

(Communicated by Joseph A. Ball)

ABSTRACT. We compute the joint (Taylor) spectrum of an n -tuple of commuting composition operators acting on the Hardy space H^2 .

1. INTRODUCTION

Let $A(\Delta)$ denote the space of all analytic functions in the unit disk Δ with the topology of uniform convergence on compact subsets of Δ , and let \mathcal{H} be a linear subspace of $A(\Delta)$. If ψ is an analytic self map of Δ such that $f \circ \psi$ belongs to \mathcal{H} for all $f \in \mathcal{H}$, then ψ induces a linear operator $C_\psi : \mathcal{H} \rightarrow \mathcal{H}$ defined as $C_\psi(f) := f \circ \psi$. C_ψ is called the *composition operator* with *symbol* ψ .

In [12] we observed that, according to a result of C. C. Cowen (cf. [4]), if $\psi : \Delta \rightarrow \Delta$ is a linear fractional map, and the composition operator C_ψ is hyponormal, then C_ψ is unitarily equivalent to a composition operator with symbol ϕ of the form

$$(1.1) \quad \phi \equiv \phi_a := \frac{z}{az + (a+1)}, \text{ with } a > 0.$$

In fact, any such ψ induces a strongly continuous one-parameter semigroup of subnormal composition operators. More precisely, we may define non-negative powers of C_{ϕ_a} by the formula

$$(1.2) \quad C_{\phi_a}^\lambda := C_{\phi_{(1+a)\lambda - 1}}.$$

Then we proved:

Theorem 1.1 ([12]). *Let $\Phi = (\phi_1, \phi_2, \dots, \phi_n)$ be an n -tuple of commuting linear fractional self-maps of Δ , neither one of them having ∞ as a fixed point. Let $C_\Phi := (C_{\phi_1}, \dots, C_{\phi_n})$. The following statements are equivalent:*

- (i) C_{ϕ_k} is hyponormal for some k .
- (ii) C_Φ is (jointly) subnormal.
- (iii) C_Φ is (jointly) hyponormal.
- (iv) There exist positive real numbers $\lambda_1, \lambda_2, \dots, \lambda_n$, such that $C_\Phi \cong (C_\psi^{\lambda_1}, C_\psi^{\lambda_2}, \dots, C_\psi^{\lambda_n})$, where $\psi := \frac{z}{z+2}$.

Received by the editors January 24, 2001.

2000 *Mathematics Subject Classification*. Primary 47B33; Secondary 47A13, 47B20.

This paper is part of the author's doctoral dissertation at the University of Iowa, written under the supervision of Professor Raúl Curto.

Observe that for each $a > 0$, ϕ_a has Denjoy-Wolff fixed point 0, and a single fixed point, 1, on the boundary. It follows from [5, Theorem 7.30, page 289] that the spectrum of C_{ϕ_a} is

$$(1.3) \quad \sigma(C_{\phi_a}) = \left\{ z \in \mathbb{C} : |z| \leq \frac{1}{(1+a)^{1/2}} \right\} \cup \{1\}.$$

In this note we compute the Taylor spectrum of a commuting n -tuple $(C_{\phi_1}, C_{\phi_2}, \dots, C_{\phi_n})$, where the ϕ_i 's are as in (1.1).

2. JOINT SPECTRUM

As pointed out in [7], when dealing with n -tuples of elements in a Banach algebra, the (joint) spectral theory is more complicated. In this case, not only is there a difference between the study of commutative and non-commutative n -tuples, but also there is a distinction between algebraic and spatial joint spectra. The notion of *Taylor spectrum* belongs to the spatial theory.

Definition 2.1 (Taylor Spectrum). Let $\Lambda = \Lambda[e] = \Lambda_n[e]$ be the *exterior algebra* on n generators e_1, e_2, \dots, e_n , with identity $e_0 \equiv 1$. That is, Λ is the algebra of forms in e_1, e_2, \dots, e_n , such that $e_i \wedge e_j = -e_j \wedge e_i$ ($1 \leq i, j \leq n$). Λ is a graded algebra, for if Λ_n^k is the algebra of k -forms on e_1, e_2, \dots, e_n , then $\Lambda = \bigoplus_{k=-\infty}^{\infty} \Lambda_n^k$, with $\Lambda_n^k \wedge \Lambda_n^l \subset \Lambda_n^{k+l}$.

For each $i \in \{1, 2, \dots, n\}$ define $E_i : \Lambda \rightarrow \Lambda$ by

$$E_i \xi := e_i \wedge \xi.$$

E_1, E_2, \dots, E_n are called the *creation operators*, and they clearly satisfy the property $E_i E_j = -E_j E_i$ ($1 \leq i, j \leq n$).

Λ can be regarded as a Hilbert space if one declares the set

$$\{\xi = e_{i_1} \wedge e_{i_2} \wedge \dots \wedge e_{i_k} : i_1 < i_2 < \dots < i_k\}$$

as an orthonormal basis. Then $E_i^* \xi = \xi'$, where $\xi = e_i \xi' + \xi''$ is the unique decomposition of a form ξ as the sum of an element in the range of E_i and an element in the kernel of E_i^* (actually each E_i is a partial isometry, and $E_i^* E_j + E_j E_i^* = \delta_{ij}$).

For \mathcal{B} a Banach space and $A = (A_1, A_2, \dots, A_n)$ an n -tuple of commuting operators in $\mathcal{L}(\mathcal{B})$, define $D_A : \Lambda(\mathcal{B}) \rightarrow \Lambda(\mathcal{B})$ ($\Lambda(\mathcal{B}) := \mathcal{B} \otimes_{\mathbb{C}} \Lambda$) by

$$D_A := \sum_{i=1}^n A_i \otimes E_i.$$

Then

$$D_A^2(x \wedge \xi) = \sum_{i=1}^n A_j A_i x \otimes E_j E_i \xi = \sum_{i < j} A_i A_j x \otimes (E_i E_j + E_j E_i) \xi = 0.$$

Set $\Lambda^k(\mathcal{B}) := \mathcal{B} \otimes_{\mathbb{C}} \Lambda_n^k$ and $D_A^k := D_A|_{\Lambda^k(\mathcal{B})}$. Then we can construct a cochain complex $K(A, \mathcal{B})$, called the *Koszul complex* associated to A on \mathcal{B} , as follows:

$$K(A, \mathcal{B}) : \quad \mathbf{0} \longrightarrow \Lambda^0(\mathcal{B}) \xrightarrow{D_A^0} \Lambda^1(\mathcal{B}) \xrightarrow{D_A^1} \dots \xrightarrow{D_A^{n-1}} \Lambda^n(\mathcal{B}) \longrightarrow \mathbf{0}.$$

The Taylor spectrum of A on \mathcal{B} is then the set

$$\sigma_T(A, \mathcal{B}) := \{\lambda \in \mathbb{C}^n : K(A - \lambda, \mathcal{B}) \text{ is not exact}\}.$$

$\sigma_T(A, \mathcal{B})$ generalizes the one variable notion of spectrum (see Example 2.2(i) below); it is a nonempty compact subset of \mathbb{C}^n that carries an analytic functional calculus: If Ω is an open set in \mathbb{C}^n containing $\sigma(A, \mathcal{B})$, then $\sigma_T(f(A), \mathcal{B}) = f(\sigma_T(A, \mathcal{B})) \quad \forall f \in A(\Omega)$; see [20], [21]. The reader can find an excellent account of the Taylor spectrum and its relations with other multiparameter spectral theories in [7].

Example 2.2. Let \mathcal{H} be a Hilbert space.

(i) For $A = (A_1)$,

$$K(A, \mathcal{H}) : \quad \mathbf{0} \longrightarrow \mathcal{H} \xrightarrow{D_A^0} \mathcal{H} \longrightarrow \mathbf{0}$$

with $D_A^0 = A_1$. Clearly, $K(A, \mathcal{H})$ is exact if and only if A_1 is one-to-one and onto. Hence in this case, $\sigma_T(A, \mathcal{H}) = \sigma(A)$, the usual spectrum of A .

(ii) Suppose that $A_1, A_2 \in \mathcal{L}(\mathcal{H})$ commute. Then the *Koszul complex* associated to $A = (A_1, A_2)$ on \mathcal{H} is the cochain complex

$$K(A, \mathcal{H}) : \quad \mathbf{0} \longrightarrow \mathcal{H} \xrightarrow{D_A^0} \mathcal{H} \oplus \mathcal{H} \xrightarrow{D_A^1} \mathcal{H} \longrightarrow \mathbf{0},$$

where

$$D_A^0(f) := (A_1 f, A_2 f)$$

and

$$D_A^1(f, g) := -A_2 f + A_1 g.$$

□

Example 2.3. Let (C_ϕ, C_ψ) be a subnormal pair of (commuting operators). Then by Theorem 1.1

$$(C_\phi, C_\psi) \cong (C_{\phi_a}, C_{\phi_a}^\lambda),$$

where $\phi_a \equiv \frac{z}{az+1+a}$, $a > 0$, and $\lambda > 0$.

Assume now that $\lambda \in \mathbb{Q}$, say $\lambda = m/n$, with m, n positive integers. Choose b such that $(1+b)^n = 1+a$. Then

$$(C_{\phi_a}, C_{\phi_a}^\lambda) = (C_{\phi_b}^n, C_{\phi_b}^m)$$

and therefore, by the functional calculus,

$$\sigma_T((C_\phi, C_\psi), H^2) = F(\sigma(C_{\phi_b}))$$

where $F : \Delta \rightarrow \mathbb{C}^2$ is the polynomial mapping defined by $F(z) := (z^n, z^m)$.

By (1.3) it follows that

$$\sigma_T((C_{\phi_a}, C_{\phi_a}^\lambda), H^2) = \{(z^n, z^m) : |z| \leq \frac{1}{(1+b)^{1/2}}\} \cup \{(1, 1)\}.$$

Now by setting $w := z^n$ (for $z \neq 0$) we can write

$$\sigma_T((C_{\phi_a}, C_{\phi_a}^\lambda), H^2) = \{(w, w^\lambda) : 0 < |w| \leq \frac{1}{(1+a)^{1/2}}\} \cup \{(0, 0), (1, 1)\}.$$

Thus if $f_\lambda : \sigma(C_{\phi_a}) \rightarrow \mathbb{C}$ is the (multivalued) function defined by

$$(2.1) \quad f_\lambda(z) = \begin{cases} 0 & \text{if } z = 0, \\ 1 & \text{if } z = 1, \\ z^\lambda & \text{if } z \neq 0, z \neq 1, \end{cases}$$

then we have

$$\sigma_T((C_\phi, C_\psi), H^2) = \sigma_T((C_{\phi_a}, C_{\phi_a}^\lambda), H^2) = f_\lambda(\sigma(C_{\phi_a})).$$

We thus observe that the spectral mapping theorem holds in this particular situation. \square

It is interesting to notice that although the functional relation

$$\mathbb{R}^+ \ni \lambda \longrightarrow C_{\phi_a}^\lambda$$

was not achieved via a standard functional calculus, such as the Riesz analytic functional calculus, it was still possible in the example above to apply the spectral mapping theorem. The core of the problem is, of course, the fact that the function $z \rightarrow z^\lambda$, $\lambda > 0$, is not analytic on $\sigma(C_{\phi_a})$. To extend the above result to the general case we will need to combine results from general theory of strongly continuous semigroups of operators, theory of semigroups of composition operators and multiparameter spectral theory (of which Theorem 2.13 will be our main tool).

We now begin by recalling some facts about strongly continuous semigroups of contractions.

Let $\mathcal{T} = \{T(s)\}_{s \geq 0}$ be a strongly continuous semigroup of contractions in a Hilbert space \mathcal{H} . The *infinitesimal generator* A of \mathcal{T} is defined by

$$Ah := \lim_{s \rightarrow +0} \frac{1}{s}[T(s) - I]h$$

whenever this limit exists. A is a closed operator with domain $\mathfrak{D}(A)$ dense in \mathcal{H} ; also, A determines the semigroup \mathcal{T} uniquely (cf. [18]).

A proof of the next theorem can be found in [1].

Theorem 2.4. *Let A be the infinitesimal generator of the strongly continuous semigroup $\{T(s)\}_{s > 0} \subset \mathcal{L}(\mathcal{H})$. Then:*

- (i) $Re(Ah, h) \leq 0$ for $h \in \mathfrak{D}(A)$.
- (ii) $\lambda I - A$ is invertible whenever $Re(\lambda) > 0$ and, for such λ ,

$$\|(\lambda I - A)^{-1}\| \leq 1/Re(\lambda)$$

and

$$(\lambda I - A)^{-1}h = \int_0^\infty e^{-\lambda s} T(s)h ds, \quad h \in \mathcal{H}.$$

Conversely, if A is an operator satisfying $\|(\lambda I - A)^{-1}\| \leq 1/\lambda$ for $\lambda > 0$, then there exists a unique semigroup of contractions whose generator is A .

It follows from Theorem 2.4(ii) that $A - I$ is boundedly invertible and it is easy to verify (cf. [1]) that the operator T defined by

$$(2.2) \quad T := (A + I)(A - I)^{-1}$$

is a contraction. T is called the *co-generator* (cf. [18]) of the semigroup \mathcal{T} .

From (2.2) it follows that

$$T - I = (A + I)(A - I)^{-1} - I = 2(A - I)^{-1}.$$

Thus $(T - I)^{-1}$ exists and $A = (T + I)(T - I)^{-1}$. Note that the existence of $(T - I)^{-1}$ is equivalent to the fact that 1 is not an eigenvalue of T . The co-generator T can be used instead of A to study the semigroup \mathcal{T} . It has the advantage that T is bounded, while A is generally unbounded, and this actually occurs in the case of the semigroups $\{C_{\phi_a}^\lambda\}$; cf. [2], [17].

The following theorem ([19], Theorem 8.1) shows an explicit functional relation between T and \mathcal{T} .

Theorem 2.5. *Let T be a contraction in \mathcal{H} . In order that there exists a strongly continuous semigroup $\mathcal{T} = \{T(s)\}_{s \geq 0}$ whose co-generator is T it is necessary and sufficient that 1 not be an eigenvalue of T . In this case T and \mathcal{T} determine one another by the relations*

$$(2.3) \quad T(s) = e_s(T)$$

and

$$T = \lim_{s \rightarrow +0} \phi_s(T(s)) \quad (SOT)$$

where

$$(2.4) \quad e_s(\lambda) := \exp \left[s \frac{\lambda + 1}{\lambda - 1} \right] \quad \text{and} \quad \phi_s(T(s)) := \frac{\lambda - 1 + s}{\lambda - 1 - s}.$$

Corollary 2.6. *Let T be the co-generator of a strongly continuous semigroup $\mathcal{T} = \{T(s)\}_{s \geq 0}$ of contractions in $L(\mathcal{H})$. Then*

$$(2.5) \quad (I - T)f = 2 \int_0^\infty e^{-t} e_t(T)f \, dt, \quad f \in \mathcal{H}.$$

Proof. This is a direct consequence of Theorems 2.4 and 2.5. □

The next example (see [11]) shows the kind of hurdles one might expect when dealing with spectral functional calculus and semigroups.

Example 2.7 (Foiaş-Mlak). Let $\mathcal{H} = L_2(0, 1)$ and let $\{T_t\}_{t > 0}$ be a strongly continuous semigroup of contractions defined by

$$(2.6) \quad (T_t f)(x) := \begin{cases} 0 & \text{if } x < t \text{ and } 0 \leq t < 1, \\ f(x - t) & \text{if } x \geq t \text{ and } 0 \leq t < 1, \\ 0 & \text{if } t \geq 1. \end{cases}$$

Let T be the co-generator of $\{T_t\}$. Then $T_t = e_t(T)$.

Since $e_1(T) \equiv 0$, $\sigma(e_t(T)) = \{0\}$. It can be shown, on the other hand, that $\sigma(T) = \{1\}$. Therefore the expression

$$\sigma(u(T)) = u(\sigma(T)), \quad u \in H^\infty,$$

does not make sense in general. Moreover, since the set L of limit values of $e_1(z)$ at $z = 1$ equals $\bar{\Delta}$, $u(\sigma(T))$ may not even be replaced by the set of limit values of u at $\sigma(T)$. □

In general, weaker spectral mapping relations hold:

Proposition 2.8. (Spectral mapping theorems for semigroups of operators [14].) *Let A be the infinitesimal generator of the strongly continuous semigroup $\{T(s)\}_{s > 0} \subset \mathcal{L}(\mathcal{H})$. Then:*

- (a) $e^{t\sigma(A)} \subset \sigma(T_t)$ for $t \geq 0$.
- (b) $e^{t\sigma_p(A)} = \sigma_p(T_t) \setminus \{0\}$, $t \geq 0$, where σ_p denotes point spectrum.
- (c) If $\{T(s)\}_{s > 0}$ consists of normal operators, then

$$\sigma(T_t) = \overline{e^{t\sigma(A)}}, \quad \text{for } t \geq 0.$$

(d) If $\{T(s)\}_{s \in \mathbb{R}}$ is a bounded strongly continuous group, then

$$\sigma(T_t) = \overline{e^{t\sigma(A)}}, \text{ for } t \in \mathbb{R}.$$

Let $H_0^2 := H^2 - \langle \mathbf{1} \rangle$ (the orthogonal complement of the constant functions in H^2). Since for all $s \in \mathbb{C}$, $C_{\phi_a}(s\mathbf{1}) = s\mathbf{1} \circ \phi_a = s$ and $C_{\phi_a}^*(s\mathbf{1}) = sC_{\phi_a}^*(K_0) = sK_{\phi_a(0)} = s$ it follows that $\langle \mathbf{1} \rangle$ reduces C_{ϕ_a} and therefore that H_0^2 is a reducing subspace for C_{ϕ_a} .

Lemma 2.9. *The (relative) semigroup $\{C_{\phi_a}^\lambda|_{H_0^2}\}$ consists of completely non-unitary contractions (i.e. there are no non-zero reducing subspaces \mathcal{H}_λ of H_0^2 such that $C_{\phi_a}^\lambda|_{\mathcal{H}_\lambda}$ is a unitary operator).*

Proof. If there exist $\lambda > 0$ and a non-zero subspace $\mathcal{H}_\lambda \subset H_0^2$ such that $C_{\phi_a}^\lambda|_{\mathcal{H}_\lambda}$ is a unitary operator, then for all $0 \neq h \in \mathcal{H}_\lambda$ and $n = 0, 1, 2, \dots$

$$\|(C_{\phi_a}^\lambda)^n h\| = \|h\| = \|(C_{\phi_a}^{\lambda*})^n h\|.$$

Now

$$\|C_{\phi_a}^{n\lambda} h\| = \|h(\phi_a^{n\lambda})\| = \|h(\phi_{(1+a)^{n\lambda}-1})\|.$$

Since $\lim_{n \rightarrow \infty} \phi_{(1+a)^{n\lambda}-1} = 0$ (the Denjoy-Wolff fixed point) uniformly on any compact subset of Δ , and $h(0) = 0$, one must necessarily have $\|h\| = 0$, a contradiction. \square

Proposition 2.10. *Let $\{T_s\}_{s>0}$ be a (s.c.) semigroup of contractions with infinitesimal generator A and co-generator T . Then $\sigma(T) = \theta(\sigma(A))$ where θ is the Cayley transform $\theta(\mu) = \frac{\mu+1}{\mu-1}$.*

Proof. The result is a consequence of the identity

$$\mu - A = [(\mu - 1)T - (\mu + 1)](T - 1)^{-1} = \frac{1}{\mu - 1}(T - \theta(\mu))(T - 1)^{-1}.$$

Indeed, it is readily seen that if $(T - \lambda)^{-1}$ exists, then

$$S := \frac{1}{\mu - 1}(T - 1)(T - \lambda)^{-1} \quad (\mu := \theta^{-1}(\lambda))$$

is an inverse for $\mu - A$.

Conversely if $(\mu - A)^{-1}$ exists, then

$$\frac{1}{\theta(\mu) - 1}(I - A)((\mu - A)^{-1}$$

is an inverse for $\theta(\mu) - T$. \square

Proposition 2.11. *Let A be the infinitesimal generator of the normalized semigroup $\{e^{-sk}C_{\phi_a}^\lambda|_{H_0^2}\}$, and let $k := \ln\|\frac{1}{(1+a)^{1/2}}\|$. Then $\sigma(A) = \{z | \operatorname{Re}(z) \leq 0\}$.*

The proof is an adaptation of an argument given by A. G. Siskakis in [17, p. 238].

Proof. It is easy to see that $\frac{\partial}{\partial t}(\phi_a^t)|_{t=0}(z) = -lz(1+z)$, where $l := \log(1+a)$. Thus the infinitesimal generator, A , of $\{C_{\phi_a}^t|_{H_0^2}\}$, is given by the formula

$$Af(z) = -lz(1+z)f'(z),$$

whenever $f' \in H^2$.

Let $\lambda \in \mathbb{C}$ be such that $\operatorname{Re}(\lambda) \leq -1/2$, set

$$P_n(z) := 1 + \sum_{j=1}^n \frac{\lambda}{j} \binom{n}{j} z^j,$$

and let $y = f_{n,\lambda}(z) := (1+z)^\lambda e^{P_n(z)}$. Then it is readily seen that $y' = \lambda((1+z)^n - 1)$ and that

$$lz(1+z)y' = \lambda l(1+z)^{\lambda+1+n} e^{P_n(z)} = \lambda l f_{n,n+1+\lambda}.$$

That is,

$$(\lambda l - A)y = \lambda l f_{n,n+1+\lambda}.$$

Choose n such that $\operatorname{Re}(n+1+\lambda) > -1/2$. Then $f_{n,n+1+\lambda} \in H^2$.

If $\lambda l - A$ were invertible, then $\lambda l R(A, \lambda) f_{n,n+1+\lambda} = f_{n,\lambda} \in H^2$, a contradiction, since $\operatorname{Re}(\lambda) \leq -1/2$. Hence

$$(2.7) \quad \{\mu : \operatorname{Re}(\mu) \leq -l/2\} = \{\lambda l : \operatorname{Re}(\lambda) \leq -1/2\} \subseteq \sigma(A).$$

On the other hand, by (1.3)

$$(2.8) \quad \begin{aligned} \sigma(C_{\phi_a}^t) &= \sigma(C_{\phi_{(1+a)^t-1}}) = \{z \in \mathbb{C} : |z| \leq \frac{1}{(1+a)^{t/2}}\} \cup \{1\} \\ &= \{z \in \mathbb{C} : |z| \leq e^{-\frac{t}{2}l}\} \cup \{1\}. \end{aligned}$$

Therefore from (2.7) and Proposition 2.8(i) we have that

$$\sigma(A) = \{z : \operatorname{Re} z \leq -l/2\} \cup \{0\},$$

and consequently

$$\sigma(A|_{H_0^2}) = \{z : \operatorname{Re} z \leq -l/2\}.$$

Now consider the normalized semigroup $\{e^{-kt} C_{\phi_a}^t|_{H_0^2}\}$, where $k = \log \|C_{\phi_a}\|$. Its generator \tilde{A} is given by

$$\begin{aligned} \tilde{A} &= \frac{\partial}{\partial t} (e^{-kt} C_{\phi_a}^t)|_{t=0} = -k e^{-kt} C_{\phi_a}^t|_{t=0} + e^{-kt} \frac{\partial}{\partial t} (C_{\phi_a}^t)|_{t=0} \\ &= -k + A \\ &= \frac{1}{2}l + A. \end{aligned}$$

Consequently

$$\mu - \frac{1}{2}l - A \text{ is invertible} \iff \operatorname{Re}(\mu - \frac{1}{2}l) > -\frac{1}{2}l \iff \operatorname{Re}(\mu) > 0.$$

Therefore

$$\sigma(\tilde{A}) = \{z : \operatorname{Re} z \leq 0\}.$$

□

Corollary 2.12. *Let T be the co-generator of the semigroup $\{e^{-sk} C_{\phi_a}^s|_{H_0^2}\}$. Then $\sigma(T) = \overline{\Delta}$.*

Proof. This is an immediate consequence of Propositions 2.11 and 2.10. □

Next we cite a result, due to R. Curto and L. Fialkow [8], which is crucial to attaining our goal.

Theorem 2.13 (R. Curto - L. Fialkow). *Let T be a c.n.u. contraction and let u be such that $u(\Delta) \subset \overline{u(\Delta \cap \sigma(T))}$. Then $\sigma_T(u(T)) = \overline{u(\Delta)}$.*

By virtue of the preceding discussion, the following proposition is a corollary of the previous theorem.

Theorem 2.14. *Let $\psi \equiv z/(z+2)$, and let $C_\Psi := (C_\psi^{\lambda_1}, C_\psi^{\lambda_2}, \dots, C_\psi^{\lambda_n})$. Then*

$$\sigma_T(C_\Psi) = \overline{u(\Delta)} \cup \{(1, 1, \dots, 1)\}$$

where $u \in H^\infty$ is the function defined by

$$(2.9) \quad u(z) = (e^{k\lambda_1} e_{\lambda_1}(z), e^{k\lambda_2} e_{\lambda_2}(z), \dots, e^{k\lambda_n} e_{\lambda_n}(z)),$$

with $e_{\lambda_i}(z) = \exp\left[\lambda_i \frac{z+1}{z-1}\right]$ and $k := \ln\|C_\psi\|_{H_0^2}$ ($= \frac{-\ln 2}{2}$).

Remarks. • The function ψ in Theorem 2.14 is equal to ϕ_a with $a = 1$.

- By (1.3) $\sigma(C_\psi) = \{e^k e_1(z) : z \in \Delta\} \cup \{1\}$.
- Putting $w := e^k e_1(z)$, $z \in \Delta$, then

$$\sigma_T(C_\Psi) = \overline{\{(w^{\lambda_1}, w^{\lambda_2}, \dots, w^{\lambda_n})\}} \cup \{(1, 1, \dots, 1)\}.$$

I would like to thank my advisor Dr. Raúl Curto, who suggested this line of work, for his constant guidance throughout the preparation of this article.

REFERENCES

- [1] H. Bercovici, *Operator Theory and Arithmetic in H^∞* , Mathematical Surveys and Monographs, 26, A.M.S., 1988. MR **90e**:47001
- [2] E. Berkson and H. Porta, *Semigroups of analytic functions and composition operators*, Michigan Math. J. 25 (1978), 101-115. MR **58**:1112
- [3] C. C. Cowen, *Composition operators on H^2* , J. Operator Theory 9 (1983), 77-106. MR **84d**:47038
- [4] C. C. Cowen, *Linear fractional composition operators on H^2* , Integral Equations Operator Theory 11 (1988), 151-160. MR **89b**:47044
- [5] C. C. Cowen, and B. D. MacCluer, *Composition Operators on Spaces of Analytic Functions*, CRC Press, New York, 1995. MR **97i**:47056
- [6] C. C. Cowen, and T. L. Kriete, *Subnormality and composition operators on H^2* , J. Funct. Anal. 81 (1988), 298-319. MR **90c**:47055
- [7] R. Curto, *Applications of several complex variables to multiparameter spectral theory* (J. B. Conway and B. B. Morrel, eds.), Surveys of Some Recent Results in Operator Theory, Vol. II, Longman Publ. Co., London (1988), 25-90. MR **90d**:47007
- [8] R. Curto and L. Fialkow, *Elementary operators with H^∞ symbol*, Integral Equations Operator Theory 10 (1987), 707-720. MR **88j**:47016
- [9] R. Curto, P. Muhly and J. Xia, *Hyponormal pairs of commuting operators*, Oper. Theory: Adv. Appl., Vol. 35, Birkhäuser-Verlag Basel (1988), 1-22. MR **90m**:47037
- [10] F. Jafary, B. D. MacCluer, C. C. Cowen and D. Porter, *Studies on Composition Operators*, Contemp. Math., 213, RI, 1997. MR **98h**:47005
- [11] C. Foias and W. Mlak, *The extended spectrum of completely non-unitary contractions and the spectral mapping theorem*, Studia Math. XXVI (1966), 239-245. MR **34**:610
- [12] J. Giménez, *Joint hyponormality of composition operators with linear fractional symbols*, Preprint, 2000.
- [13] T. Itô, *On the commutative family of subnormal operators*, J. Fac Sci. Hokkaido Univ. 14 (1958), 1-15. MR **21**:5902
- [14] K-J. Engel and R. Nagel, *One-Parameter Semigroups for Linear Evolution Equations*, Grad. Texts in Math., 194, Springer-Verlag, New York, 2000. MR **2000i**:47075
- [15] A. I. Plesner, *Spectral Theory of Linear operators*, Vol. II, Frederick Ungar Publishing Co., New York, 1969. MR **39**:6106

- [16] J. H. Shapiro, *Composition Operators and Classical Function Theory*, Springer -Verlag, New York, 1993. MR **94k**:47049
- [17] A. G. Siskakis, *Semigroups of composition operators on spaces of analytic functions, a review*, Studies on Composition Operators, Contemporary Math, AMS, 213, 1988, 229-252. MR **98m**:47049
- [18] B. Sz.-Nagy, *Spektraldarstellung linearer transformationen des Hilbertschen raumes*, Ergebnisse der Mathematik, Springer-Verlag, Berlin, 1967. MR **35**:4744
- [19] B. Sz.-Nagy and C. Foiaş, *Harmonic Analysis of Operators on Hilbert Space*, North-Holland Publ. Co., 1970. MR **43**:947
- [20] J. L. Taylor, *A joint spectrum of several commuting operators*, J. Funct. Anal. 6 (1970), 172-191. MR **42**:3603
- [21] J. L. Taylor, *The analytic functional calculus for several commuting operators*, Acta Math. 125 (1970), 1-38. MR **42**:6622

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE LOS ANDES, MÉRIDA, VENEZUELA
E-mail address: jgimenez@ciens.ula.ve