

A PROOF OF A PARTITION THEOREM FOR $[\mathbb{Q}]^n$

VOJKAN VUKSANOVIC

(Communicated by Alan Dow)

ABSTRACT. In this note we give a proof of Devlin's theorem via Milliken's theorem about weakly embedded subtrees of the complete binary tree $2^{<\mathbb{N}}$. Unlike the original proof which is (still unpublished) long and uses the language of category theory, our proof is short and uses direct combinatorial reasoning.

A tree is any partially ordered set (T, \leq) such that for any $t \in T$ the set $\{s \in T : s \leq t\}$ is well ordered by the induced order. Given a tree T we say that S is a subtree of T if $S \subseteq T$. From now on we will suppose that every tree has a root, i.e. we will suppose that every tree has the unique minimal element. We will also suppose that for a given tree T , for every $t \in T$, $\{s \in T : s \leq t\}$ is finite. By $Succ(t, T)$ we will denote the set $\{s \in T : t \leq s\}$ and by $Pred(t, T)$ we will denote the set $\{s \in T : s \leq t\}$. Given a tree T and $t \in T$ by $IS(t, T)$ we will denote the set of immediate successors of t in T . By $T(n)$ we will denote the n -th level of T , i.e. the set of all $t \in T$ such that $|\{s \in T : s \leq t, s \neq t\}| = n$. For a tree T , $height(T)$ will denote $\sup\{n + 1 : T(n) \neq \emptyset\}$. We say that a tree (T, \leq) is *perfect* if for every $t \in T$ there is $s \in T$ such that $t \leq s$ and $|IS(s, T)| \geq 2$. Given a tree T and nodes s and t in T by $s \wedge t$ we will denote the maximal node in T which is below both s and t . If $A \subseteq T$, by $\wedge(A)$ we will denote the \wedge -closure of A , the smallest subset A' of T containing A such that $(\forall s, t \in A') s \wedge t$ belongs to A' .

Most of the time we shall be working inside the complete binary tree $2^{<\mathbb{N}}$ ordered by end-extension which we denote by \subseteq . We shall also need to consider the two orderings on $2^{<\mathbb{N}}$, $<_{lex}$ and $\leq_{\mathbb{Q}}$, which we introduce in the following two definitions.

Definition 0.1. Let $s, t \in 2^{<\mathbb{N}}$. We say that s is *lexicographically less* than t and write $s <_{lex} t$ provided the following hold: s and t are incomparable and if $i \in \mathbb{N}$ is the maximal integer such that $s \upharpoonright i + 1 = t \upharpoonright i + 1$, then $s(i + 1) < t(i + 1)$.

Definition 0.2. Given $s, t \in 2^{<\mathbb{N}}$ put $t \leq_{\mathbb{Q}} s$ iff $s = t$ or $((s \subseteq t \text{ and } s \hat{\ } 0 \subseteq t)$ or $(t \subseteq s \text{ and } t \hat{\ } 1 \subseteq s)$ or $(s \text{ and } t \text{ are incomparable and } t <_{lex} s)$.

Note that while $\leq_{\mathbb{Q}}$ is a total order $<_{lex}$ is not. But, $<_{lex}$ is a total order on any antichain of $2^{<\mathbb{N}}$ and in fact it agrees with $\leq_{\mathbb{Q}}$. It is also easy to check that $(2^{<\mathbb{N}}, \leq_{\mathbb{Q}})$ has the order type η .

Let $A \in [2^{<\mathbb{N}}]^n$ for some $n \in \mathbb{N}$. From now on by $\{a_1, \dots, a_n\}_{\leq_{\mathbb{Q}}}$ we will denote the enumeration of elements of A with respect to $\leq_{\mathbb{Q}}$.

Received by the editors March 29, 2001 and, in revised form, May 29, 2001.
2000 *Mathematics Subject Classification.* Primary 05A18.
Key words and phrases. Partitions of rationals.




FIGURE 1.

Definition 0.3. Suppose that S is a subtree of T . We say that S is *strongly embedded* in T (see Figure 1 for the case $T = 2^{<\mathbb{N}}$) if the following hold:

- 1) If s is a nonmaximal in S and $t \in IS(s, T)$, then $Succ(t, T) \cap IS(s, S)$ is a singleton,
- 2) If $height(T) = \omega$ and $height(S) \leq \omega$ there is strictly increasing function $f : height(S) \rightarrow \omega$ such that $S(n) \subseteq T(f(n))$ for each $n \in height(S)$.

Suppose in the previous definition we drop requirement 2), and instead of 1) we have:

- 1') If s is a nonmaximal in S and $t \in IS(s, T)$, then $Succ(t, T) \cap IS(s, S)$ is either a singleton or empty.

Then we say that S is *weakly embedded* in T .

By $WEm^{<\omega}(T)$ we will denote the set of all finite weakly embedded subtrees of T .

Definition 0.4. Given a strongly embedded subtree T of $2^{<\mathbb{N}}$ and trees A and B weakly embedded in T , we say that A and B have the same *embedding type* and we write $A \sim_{Em} B$ provided the following hold:

- 1) There is a bijection $f : A \rightarrow B$ satisfying $a \subseteq a'$ iff $f(a) \subseteq f(a')$.
- 2) If $a \in A \cap T(n)$, $a' \in A \cap T(n')$, $f(a) \in B \cap T(m)$, and $f(a') \in B \cap T(m')$, then $n < n'$ iff $m < m'$.
- 3) Suppose $n \in \omega$ and there is $d \in A \cap T(n)$. Suppose $a \in T(n)$ with $(Succ(a, T) \setminus \{a\}) \cap A \neq \emptyset$. Pick $c \in (Succ(a, T) \setminus \{a\}) \cap A$, and let b be the unique node with $b \in Pred(c, T) \cap T(n + 1)$. Suppose $f(d) \in T(m)$, and write a' and b' for the unique nodes $a' \in (Pred(f(c), T) \setminus \{f(c)\}) \cap T(m)$ and $b' \in Pred(f(c), T) \cap T(m + 1)$. Then we require that $a \hat{=} 0 \subseteq b$ iff $a' \hat{=} 0 \subseteq b'$ (see Figure 2).

Note that this is not the same as the original definition given in [3]. Originally, we have that in a finitely branching tree T , for each $t \in T$ there is a linear order \prec on $IS(t, T)$. In other words, for each $t \in T$, $IS(t, T)$ can be enumerated as

$$IS(t, T) = \{is(t, T)(j) : j \in |IS(t, T)|\}$$

so that

$$is(t, T)(i) \prec is(t, T)(j) \Leftrightarrow i \in j \in |IS(t, T)|.$$




FIGURE 2.

Also, in 3) of the previous definition of the embedding type of a weakly embedded subtree, it is required that for each $i \in |Is(a, T)|$, $b = is(a, T)(i)$ iff $b' = is(a', T)(i)$. In the case of strongly embedded subtrees of the complete binary tree we naturally take the ordering $<_{lex}$ to be the linear order $<$. Then it is easy to see that our definition is equivalent to the original one when working with strongly embedded subtrees of $2^{<\mathbb{N}}$. We write $Em^A(T)$ for the collection of all weakly embedded subtrees B of T with $A \sim_{Em} B$. The following theorem, which is due to Milliken [3], is crucial for our proof.

Theorem 0.5. *Suppose that T is a perfect strongly embedded subtree of $2^{<\mathbb{N}}$, that $A \in WEm^{<\omega}(T)$, and that $Em^A(T) = \bigcup_{i \in r} C_i$. Then there is a perfect strongly embedded subtree S of T and $k \in r$ such that $Em^A(T) \cap WEm^{<\omega}(S) \subseteq C_k$.*

We will now state the theorem which is due to Devlin [2].

Theorem 0.6. *We have:*

- 1) $\eta \rightarrow (\eta)_{<\omega/t_n}^n$,
- 2) $\eta \not\rightarrow (\eta)_{<\omega/t_{n-1}}^n$

$$\text{where } t_1 = 1 \text{ and } t_n = \sum_{i=1}^{n-1} \binom{2n-2}{2i-1} t_i t_{n-i} \text{ for } n \geq 2.$$

We will prove three lemmas which will prove Theorem 0.6. The first two prove 1), and the last proves 2).

From now on we identify the rationals with the set of all finite sequences of 0's and 1's.

Given a colouring $c : [2^{<\mathbb{N}}]^n \rightarrow r$, it naturally induces a colouring $c' : \mathcal{F} \rightarrow r$ where \mathcal{F} is a family of all subsets A of $2^{<\mathbb{N}}$ such that there is an $A' \in [2^{<\mathbb{N}}]^n$ and $A = \wedge(A')$. It is easy to see that we have only finitely many different embedding types which appear as a \wedge -closure of n -element subsets of $2^{<\mathbb{N}}$. For example the \wedge -closure of a pair in $2^{<\mathbb{N}}$ gives us seven different embedding types. These are (see Figure 3) $\{\{\emptyset, 0\}, \{\emptyset, 1\}, \{\emptyset, 0, 1\}, \{\emptyset, 0, 10\}, \{\emptyset, 0, 11\}, \{\emptyset, 1, 00\}, \{\emptyset, 1, 01\}\}$.

Hence, we can apply Theorem 0.5 successively finitely many times to get a perfect strongly embedded subtree T of the complete binary tree $2^{<\mathbb{N}}$ such that if $A =$

Lemma 0.8. *Let A and B be two n -element subsets of X . If $\wedge(A) \sim_{Em} \wedge(B)$, then $<_{W_A} = <_{W_B}$.*

Proof. We will prove this by induction on n . For $n = 1$ the lemma holds trivially. Suppose that the lemma is true for every $k \leq n$ and let us prove the lemma for $n+1$. Let $A, B \in [X]^{n+1}$ and let $A' = \{a_1, a_2, \dots, a_{2n+1}\}_{\leq \mathbb{Q}}$ and $B' = \{b_1, b_2, \dots, b_{2n+1}\}_{\leq \mathbb{Q}}$ where $A' = \wedge(A)$ and $B' = \wedge(B)$. Let a_k and b_l be the roots of A' and B' respectively. Set $A_1 = \{x \in A' : a_k \hat{0} \subseteq x\}$, $A_2 = \{x \in A' : a_k \hat{1} \subseteq x\}$ and similarly define B_1 and B_2 . Then, if $f : A' \rightarrow B'$ is a bijection witnessing $A' \sim_{Em} B'$ it is easy to see that $f(a_k) = f(b_l)$ and therefore $k = l$. Also, it is easy to see that $f \upharpoonright A_1$ is a bijection from A_1 onto B_1 witnessing $A_1 \sim_{Em} B_1$ and $f \upharpoonright A_2$ is a bijection from A_2 onto B_2 witnessing $A_2 \sim_{Em} B_2$. Therefore, we can apply inductive hypothesis to A_1, B_1 and to A_2 and B_2 and easily conclude the lemma. \square

Hence, from the lemma above one concludes that the embedding type of the $\wedge(A)$ of some $A \in [X]^n$ is determined completely by its well-ordering $<_{W_A}$. This will help us to count the number of possible colours for n -tuples of X .

Lemma 0.9. *For each n let t_n be the minimal number of embedding types which appear as the \wedge -closure of n -element subsets of X . Then*

$$t_1 = 1 \quad \text{and} \quad t_n = \sum_{i=1}^{n-1} \binom{2n-2}{2i-1} t_i t_{n-i}.$$

Proof. That $t_1 = 1$ is easy and well-known. So, suppose that $n > 1$ and let A' be the \wedge -closure of some n -element subset $A = \{a_1, a_2, \dots, a_n\}$ in X . Set $B' = \{x \in A' : a \hat{0} \subseteq x\}$ and $C' = \{x \in A' : a \hat{1} \subseteq x\}$ where a is the root of A' . Let $l = |\{a_i : a_i \in B'\}|$ and $r = |\{a_i : a_i \in C'\}|$. Note that $1 \leq l, r$ and $l + r = n$. Note also that $B' = \wedge(B)$ and $C' = \wedge(C)$ where $B = A \cap B'$ and $C = A \cap C'$. Note that for fixed $l, r \leq n$ we have by inductive hypothesis $t_l t_r$ different types of pairs $\langle B, C \rangle$. By the previous lemma structure A' is completely determined by its well-ordering $<_{W_A}$. The well-ordering $<_{W_A}$ interlaces the well-orderings $<_{W_B}$ and $<_{W_C}$ and adjoins a as the $<_{W_A}$ minimal element. (For $n = 2$ and $n = 3$ see Figure 6 and Figure 7 respectively.) The number of possible ways $<_{W_B}$ and $<_{W_C}$ can be interlaced to form $<_{W_A}$ is $\binom{|B'|+|C'|}{|B'|} = \binom{2n-2}{2l-1}$. Therefore, the total number of possibilities for A' is

$$t_n = \sum_{l=1}^{n-1} \binom{2n-2}{2l-1} t_l t_{n-l}. \quad \square$$

This proves 1) of Devlin's theorem. The next lemma will prove 2).

Lemma 0.10. *Let $S \subseteq X$ and let S have $\leq_{\mathbb{Q}}$ -order type η . Then for every $P \in [X]^n$ there is $A \in [S]^n$ such that $\wedge(P) \sim_{Em} \wedge(A)$.*

Proof. Set $S' = \{x \in W : \text{both } \{s \in S : x \hat{0} \subseteq s\} \text{ and } \{s \in S : x \hat{1} \subseteq s\} \text{ are densely ordered by } \leq_{\mathbb{Q}}\}$. We will prove that S' is a perfect subtree. Let s' be the root of the $\wedge(S)$. Note that $s' \in W$. Put $U = \{t \in S : s' \hat{0} \subseteq t\}$ and $V = \{t \in S : s' \hat{1} \subseteq t\}$. Then U and V are nonempty, $U \cup V = S$ and $\forall u \in U$ and $\forall v \in V$ we have that $u \leq_{\mathbb{Q}} v$. Since S has order type η , U and V are densely ordered by $\leq_{\mathbb{Q}}$ and $s' \in S'$. Given $s \in S'$ put $U = \{t \in S : s \hat{0} \subseteq t\}$ and let $U' = U \setminus \{u \in U : u \text{ is an}$




FIGURE 5.




FIGURE 6. Embedding types in the case $n = 2$

endpoint of the dense linear order $(U, \leq_{\mathbb{Q}})$. Let u be the \subseteq -least node of the $\wedge(U')$. As above, both $\{w \in U' : u \hat{0} \subseteq w\}$ and $\{w \in U' : u \hat{1} \subseteq w\}$ are densely ordered by $\leq_{\mathbb{Q}}$. Because the only points deleted from U to form U' were endpoints, $\{w \in U : u \hat{0} \subseteq w\}$ and $\{w \in U : u \hat{1} \subseteq w\}$ are also both densely ordered by $\leq_{\mathbb{Q}}$. Hence $u \in S'$. By the same argument, $v \in S'$ is found such that $s \hat{1} \subseteq v$. Note that $|IS(s, S')| = 2$ for every $s \in S'$. Let $f : S' \rightarrow S$ be a fixed one-to-one function such that

$$\text{a) } s \hat{0} \subseteq f(s)$$

for every $s \in S'$. It is easy to define such a function by induction because by the construction $\{s \in S : x \hat{0} \subseteq s\}$ is infinite for every $x \in S'$. Let Z be the perfect subtree of S' (see Figure 5) with the property that for every $x, y \in Z$:

- b) $|x| < |y|$ iff $|x| < |f(y)|$ iff $|f(x)| < |y|$ iff $|f(x)| < |f(y)|$,
- c) if $x \in Z(m), y \in Z(n)$, then $|x| < |y|$ iff $m < n$ or $(m = n \text{ and } x <_{lex} y)$.

Again, it is easy to construct such a subtree because S' is perfect. Define by induction a bijection $\psi : W \rightarrow Z$ as follows: put $\psi(\text{root}(W)) = \text{root}(Z)$. Having defined $\psi \upharpoonright \bigcup_{k \leq n} W(k)$, let $W(n+1) = \{w_1, \dots, w_{2^{n+1}}\}$ and $Z(n+1) = \{z_1, \dots, z_{2^{n+1}}\}$ be enumerations of $n+1$ -st levels of W and Z respectively such that for all $1 \leq i, j \leq 2^{n+1}$ we have $i < j$ iff $|w_i| < |w_j|$ iff $|z_i| < |z_j|$. Then put $\psi(w_i) = z_i$ for all $1 \leq i \leq 2^{n+1}$. Note that we must have necessarily:




FIGURE 7. Embedding types in the case $n = 3$

d) $w_1 \hat{0} \subseteq w_2$ iff $\psi(w_1) \hat{0} \subseteq \psi(w_2)$ for every $w_1, w_2 \in W$ and

e) $(w_1 \wedge w_2) \hat{0} \subseteq w_1$ iff $(\psi(w_1) \wedge \psi(w_2)) \hat{0} \subseteq \psi(w_1)$ for every $w_1, w_2 \in W$.

Let $P = \{p_1, \dots, p_n\}_{\leq \mathbb{Q}} \in [X]^n$. Let $P' = \{p : p \hat{0}1 \in P\}$, $A' = \psi''P'$ and $A = f''A'$. Let $f' : \wedge(P) \rightarrow \wedge(A)$ be an extension of f defined as follows: given $p_i, p_{i+1} \in P$ put $f'(p_i \wedge p_{i+1}) = f(\psi(p'_i)) \wedge f(\psi(p'_{i+1}))$ where $p'_i, p'_{i+1} \in P'$ are unique elements such that $p'_i \hat{0}1 = p_i$ and $p'_{i+1} \hat{0}1 = p_{i+1}$, and for every $p_i \in P$ put $f'(p_i) = f(\psi(p'_i))$ where $p'_i \in P'$ is the unique element such that $p'_i \hat{0}1 = p_i$. It is easy to see that f' is a bijection from $\wedge(P)$ onto $\wedge(A)$. Let us prove that f' satisfies 1), 2) and 3) of Definition 0.4. 1) follows from the definition of ψ and f and f' . By a), b) and c) we have that f' satisfies 2), and 3) follows from a), d) and e). Hence, $\wedge(P) \sim_{Em} \wedge(A)$. This finishes the proof of the lemma. \square

REFERENCES

1. D.Devlin, *Some partition theorems and ultrafilters on ω* , Ph.D. thesis, Dartmouth College (1979).
2. J.D.Halpern and H.Läuchli, *A partition theorem*, Trans. Amer. Math. Soc. 124 (1966), 360-367. MR **34**:71
3. K.Milliken, *A Ramsey Theorem for Trees*, J. Combinatorial Theory A 26 (1979), 215-237. MR **80j**:05090

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, TORONTO, ONTARIO, CANADA MS5
1A1

E-mail address: voja@math.toronto.edu