

**L^p BOUNDEDNESS OF LOCALIZATION OPERATORS
ASSOCIATED TO LEFT REGULAR REPRESENTATIONS**

M. W. WONG

(Communicated by David R. Larson)

ABSTRACT. We prove an L^p boundedness result for localization operators associated to left regular representations of locally compact and Hausdorff groups and give an application to wavelet multipliers.

1. INTRODUCTION

Let G be a locally compact and Hausdorff group on which the left Haar measure is denoted by μ . Let X be an infinite dimensional, separable and complex Hilbert space in which the inner product and the norm are denoted by (\cdot, \cdot) and $\|\cdot\|$ respectively. Let $B(X)$ be the C^* -algebra of all bounded linear operators on X . Let $\pi : G \rightarrow B(X)$ be an irreducible and unitary representation of G on X such that there exists a nonzero element φ in X for which

$$(1.1) \quad \int_G |(\varphi, \pi(g)\varphi)|^2 d\mu(g) < \infty.$$

Then we call the representation $\pi : G \rightarrow B(X)$ a square-integrable representation of G on X . We call any φ in X , such that $\|\varphi\| = 1$ and (1.1) is valid, an admissible wavelet for the square-integrable representation $\pi : G \rightarrow B(X)$ of G on X and we define the constant c_φ by

$$c_\varphi = \int_G |(\varphi, \pi(g)\varphi)|^2 d\mu(g).$$

The formula in the following theorem is known as the resolution of the identity formula.

Theorem 1.1. *Let φ be an admissible wavelet for the square-integrable representation $\pi : G \rightarrow B(X)$ of G on X . Then*

$$(x, y) = \frac{1}{c_\varphi} \int_G (x, \pi(g)\varphi)(\pi(g)\varphi, y) d\mu(g)$$

for all x and y in X .

Received by the editors February 21, 2001.

2000 *Mathematics Subject Classification.* Primary 47G10.

This research has been partially supported by the Natural Sciences and Engineering Research Council of Canada under Grant OGP0008562.

Theorem 1.1 as stated is an abridged version of Theorem 3.1 in the paper [8] by Grossmann, Morlet and Paul, where the original contributions due to Duflo and Moore in [4] are acknowledged. See also the paper [1] by Carey in this connection. A proof of Theorem 1.1 can also be found in the book [20] by Wong.

Let $F \in L^1(G) \cup L^\infty(G)$. Then it is easy to prove that the linear operator $L_{F,\varphi} : X \rightarrow X$ defined by

$$(1.2) \quad (L_{F,\varphi}x, y) = \frac{1}{c_\varphi} \int_G F(g)(x, \pi(g)\varphi)(\pi(g)\varphi, y) d\mu(g), \quad x, y \in X,$$

is a bounded linear operator on X . Using the Riesz-Thorin theorem, which we recall in Section 3, it is proved in the paper [9] by He and Wong, and in the book [20] by Wong that if $F \in L^p(G)$, $1 \leq p \leq \infty$, then there exists a unique bounded linear operator $L_{F,\varphi} : X \rightarrow X$ such that the formula (1.1) is valid for all F in $L^1(G) \cup L^\infty(G)$, and x and y in X . Furthermore,

$$\|L_{F,\varphi}\|_{B(X)} \leq c_\varphi^{-\frac{1}{p}} \|F\|_{L^p(G)},$$

where $\|\cdot\|_{B(X)}$ is the norm in $B(X)$.

Remark 1.2. The bounded linear operators $L_{F,\varphi} : X \rightarrow X$ are known as localization operators and have been studied in [5] by Du and Wong, in [9] by He and Wong, and in the book [20] by Wong. The origin of these operators can be traced back to the study of a class of bounded linear operators introduced by Daubechies in [2, 3] in signal analysis. The rationale for the terminology lies in the observation that by Theorem 1.1, the bounded linear operator $L_{F,\varphi} : X \rightarrow X$ is simply the identity operator I on X if $F(g) = 1$, $g \in G$. Thus, the role of the symbol F is to localize on the group G so as to produce a nontrivial bounded linear operator with various applications in signal analysis and quantization among others. In fact, it is proved in the paper [9] by He and Wong, and the book [20] by Wong that if $F \in L^p(G)$, $1 \leq p \leq \infty$, then the localization operator $L_{F,\varphi} : X \rightarrow X$ is in the Schatten-von Neumann class S_p .

In another direction, guided by the Landau-Pollak-Slepian operator in signal analysis, a theory of wavelet multipliers has been introduced and studied in [6] by Du and Wong, [10] by He and Wong and [20] by Wong. To recall, let $\sigma \in L^\infty(\mathbb{R}^n)$. Then we define the linear operator $T_\sigma : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$ by

$$T_\sigma u = \mathcal{F}^{-1} \sigma \mathcal{F} u, \quad u \in L^2(\mathbb{R}^n),$$

where \mathcal{F}^{-1} and \mathcal{F} are the inverse Fourier transform and the Fourier transform respectively. The Fourier transform $\mathcal{F}u$, sometimes denoted by \hat{u} , of a function u in $L^2(\mathbb{R}^n)$ is given by

$$\mathcal{F}u = \lim_{R \rightarrow \infty} \widehat{\chi_R u},$$

where χ_R is the characteristic function of the ball with center at the origin and radius R ,

$$\widehat{\chi_R u}(\xi) = (2\pi)^{-\frac{n}{2}} \int_{\mathbb{R}^n} e^{-ix \cdot \xi} \chi_R(x) u(x) dx, \quad \xi \in \mathbb{R}^n,$$

and the convergence of $\widehat{\chi_R u}$ to $\mathcal{F}u$ as $R \rightarrow \infty$ is understood to be in $L^2(\mathbb{R}^n)$. It follows from Plancherel's theorem that $T_\sigma : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$ is a bounded linear operator.

Let $B(L^2(\mathbb{R}^n))$ be the C^* -algebra of all bounded linear operators on $L^2(\mathbb{R}^n)$. Let $\pi : \mathbb{R}^n \rightarrow B(L^2(\mathbb{R}^n))$ be the unitary representation of the additive group \mathbb{R}^n on $L^2(\mathbb{R}^n)$ defined by

$$(\pi(\xi)u)(x) = e^{ix \cdot \xi}u(x), \quad x, \xi \in \mathbb{R}^n,$$

for all u in $L^2(\mathbb{R}^n)$. Let φ be any function in $L^2(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$ such that $\|\varphi\|_{L^2(\mathbb{R}^n)} = 1$. Then it is proved in the paper [10] by He and Wong, and in the book [20] by Wong that

$$\langle \varphi u, \varphi v \rangle = (2\pi)^{-n} \int_{\mathbb{R}^n} \langle u, \pi(\xi)\varphi \rangle \langle \pi(\xi)\varphi, v \rangle d\xi$$

for all u and v in the Schwartz space \mathcal{S} , where $\langle \cdot, \cdot \rangle$ is the inner product in $L^2(\mathbb{R}^n)$.

Let $\sigma \in L^1(\mathbb{R}^n) \cup L^\infty(\mathbb{R}^n)$. Then it is easy to prove that the linear operator $P_{\sigma, \varphi} : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$ defined by

$$(1.3) \quad \langle P_{\sigma, \varphi} u, v \rangle = \int_{\mathbb{R}^n} \sigma(\xi) \langle u, \pi(\xi)\varphi \rangle \langle \pi(\xi)\varphi, v \rangle d\xi$$

for all u and v in $L^2(\mathbb{R}^n)$ is a bounded linear operator on $L^2(\mathbb{R}^n)$. Using the Riesz-Thorin theorem again, it is proved in [10] by He and Wong, and in [20] by Wong that if $\sigma \in L^p(\mathbb{R}^n)$, $1 \leq p \leq \infty$, then there exists a unique bounded linear operator $P_{\sigma, \varphi} : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$ such that the formula (1.3) is valid for all σ in $L^1(\mathbb{R}^n) \cup L^\infty(\mathbb{R}^n)$, and u and v in $L^2(\mathbb{R}^n)$. Moreover,

$$\|P_{\sigma, \varphi}\|_{B(L^2(\mathbb{R}^n))} \leq (2\pi)^{-\frac{n}{p}} \|\varphi\|_{L^\infty(\mathbb{R}^n)}^{\frac{2}{p'}} \|\varphi\|_{L^p(\mathbb{R}^n)},$$

where p' is the conjugate index of p and $\|\cdot\|_{B(L^2(\mathbb{R}^n))}$ is the norm in $B(L^2(\mathbb{R}^n))$.

Remark 1.3. In order to understand the linear operator $P_{\sigma, \varphi} : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$ better, let $\sigma \in L^1(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$ and let $\varphi \in L^2(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$ be such that $\|\varphi\|_{L^2(\mathbb{R}^n)} = 1$. Then it is proved in [10] by He and Wong, and in [20] by Wong that the bounded linear operators $P_{\sigma, \varphi} : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$ and $\varphi T_\sigma \bar{\varphi} : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$ are equal. Thus, the function φ plays the role of the admissible wavelet in a localization operator. Had the admissible wavelet φ in the linear operator $P_{\sigma, \varphi} : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$ been replaced by the function φ_0 on \mathbb{R}^n given by

$$\varphi_0(x) = 1, \quad x \in \mathbb{R}^n,$$

we would have obtained the pseudo-differential operator or Fourier multiplier $T_\sigma : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$ defined by

$$(T_\sigma u)(x) = (2\pi)^{-\frac{n}{2}} \int_{\mathbb{R}^n} e^{ix \cdot \xi} \sigma(\xi) \hat{u}(\xi) d\xi, \quad x \in \mathbb{R}^n.$$

See, for instance, the books [11] and [19] by Kumano-go and Wong for detailed expositions of pseudo-differential operators. Therefore it is reasonable to call the bounded linear operator $P_{\sigma, \varphi} : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$ a wavelet multiplier. That the Landau-Pollak-Slepian operator in signal analysis (studied in the fundamental papers [12, 13] by Landau and Pollak, [14, 15] by Slepian, and [16] by Slepian and Pollak) is in fact a wavelet multiplier is a result proved in [10] by He and Wong, and in [20] by Wong. As in the case of localization operators, it is proved in [10] by He and Wong, and in [20] by Wong that if $\sigma \in L^p(\mathbb{R}^n)$, $1 \leq p \leq \infty$, then the wavelet multiplier $P_{\sigma, \varphi} : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$ is in the Schatten-von Neumann class S_p .

In this paper, we let $B(L^p(G))$, $1 \leq p \leq \infty$, be the Banach algebra of all bounded linear operators on $L^p(G)$ and look at admissible wavelets associated to left regular representations $\pi : G \rightarrow B(L^p(G))$ of unimodular, locally compact and Hausdorff groups G on $L^p(G)$, and define localization operators from $L^p(G)$ into $L^p(G)$ in terms of these admissible wavelets and symbols F in $L^r(G)$, $1 \leq r \leq \infty$. An L^p boundedness result for all these localization operators is established for $1 \leq p \leq \infty$. As an application, we show that wavelet multipliers can be looked at as such localization operators when the underlying group is taken to be the additive group \mathbb{R}^n and we obtain the L^p boundedness of wavelet multipliers from $L^p(\mathbb{R}^n)$ into $L^p(\mathbb{R}^n)$ for $1 \leq p \leq \infty$.

2. L^1 BOUNDEDNESS AND L^∞ BOUNDEDNESS

Let G be a unimodular, locally compact and Hausdorff group on which the left Haar measure is denoted by μ . Let $\pi : G \rightarrow B(L^p(G))$ be the left regular representation of G on $L^p(G)$, $1 \leq p \leq \infty$, i.e., $(\pi(g)u)(h) = u(g^{-1}h)$, $g, h \in G$, for all u in $L^p(\mathbb{R}^n)$. Let $\varphi \in \bigcap_{1 \leq p \leq \infty} L^p(G)$ be such that $\|\varphi\|_{L^2(G)} = 1$. By Young's inequality on, say, page 52 of the book [7] by Folland, we get

$$(2.1) \quad c_\varphi = \int_G |(\varphi, \pi(g)\varphi)|^2 d\mu(g) \leq \|\varphi\|_{L^1(G)}^2 < \infty.$$

In view of (2.1), the fact that $\|\varphi\|_{L^2(G)} = 1$ and the notion of admissible wavelets given in Section 1, we call the function φ an admissible wavelet for the left regular representation π of G on $L^p(G)$.

Remark 2.1. In order to obtain an explicit formula for c_φ , we assume that G is a second countable, unimodular, type I, locally compact and Hausdorff group. We let \hat{G} be the set of all equivalence classes of irreducible and unitary representations of G on $L^2(G)$. Let ν be the Plancherel measure on \hat{G} . Then by Plancherel's theorem,

$$(2.2) \quad \begin{aligned} c_\varphi &= \int_G |(\varphi, \pi(g)\varphi)|^2 d\mu(g) \\ &= \int_G |(\varphi * \varphi^*)(g)|^2 d\mu(g) \\ &= \int_{\hat{G}} \text{tr}\{\hat{\psi}(\omega)\hat{\psi}(\omega)^*\} d\nu(\omega), \end{aligned}$$

where φ^* is the function on G defined by

$$(2.3) \quad \begin{aligned} \varphi^*(g) &= \overline{\varphi(g^{-1})}, \quad g \in G, \\ \psi(g) &= (\varphi * \varphi^*)(g), \quad g \in G, \end{aligned}$$

$\text{tr}\{\dots\}$ is the trace of the trace class operator $\{\dots\}$ and $\{\dots\}^*$ denotes the adjoint of the bounded linear operator $\{\dots\}$. Now,

$$(2.4) \quad \hat{\psi}(\omega) = \hat{\varphi}(\omega)^* \hat{\varphi}(\omega), \quad \omega \in \hat{G}.$$

Thus, by (2.2)-(2.4),

$$(2.5) \quad c_\varphi = \int_{\hat{G}} \text{tr}\{|\hat{\varphi}(\omega)|^4\} d\nu(\omega),$$

where

$$(2.6) \quad |\hat{\varphi}(\omega)| = (\hat{\varphi}(\omega)^* \hat{\varphi}(\omega))^{\frac{1}{2}}.$$

Therefore by (2.5) and (2.6), we get

$$(2.7) \quad c_\varphi = \int_{\hat{G}} \|\hat{\varphi}(\omega)\|_{S_4}^4 d\nu(\omega),$$

where $\|\cdot\|_{S_4}$ is the norm in the Schatten-von Neumann class S_4 . Finally, we can write (2.7) as

$$c_\varphi = \|\hat{\varphi}\|_{L^4(\hat{G}, S_4)}^4,$$

where $L^4(\hat{G}, S_4)$ is the Banach space of all S_4 -valued functions f on \hat{G} for which

$$\int_{\hat{G}} \|f(\omega)\|_{S_4}^4 d\nu(\omega) < \infty$$

and the norm $\|\cdot\|_{L^4(\hat{G}, S_4)}$ in it is given by

$$\|f\|_{L^4(\hat{G}, S_4)} = \left\{ \int_{\hat{G}} \|f(\omega)\|_{S_4}^4 d\nu(\omega) \right\}^{\frac{1}{4}}$$

for all f in $L^4(\hat{G}, S_4)$.

Let $F \in L^1(G) \cup L^\infty(G)$. Then for $1 \leq p \leq \infty$, we define the localization operator $L_{F,\varphi} : L^p(G) \rightarrow L^p(G)$ associated to the symbol F and the admissible wavelet φ by

$$(2.8) \quad (L_{F,\varphi}u, v) = \frac{1}{c_\varphi} \int_G (u, \pi(g)\varphi)(\pi(g)\varphi, v) d\mu(g)$$

for all u in $L^p(G)$ and v in $L^{p'}(G)$, where

$$(u, v) = \int_G u(g)\overline{v(g)} d\mu(g), \quad u \in L^p(G), v \in L^{p'}(G).$$

Proposition 2.2. *Let $F \in L^1(G)$. Then for $1 \leq p \leq \infty$, the localization operator $L_{F,\varphi} : L^p(G) \rightarrow L^p(G)$ is a bounded linear operator and*

$$\|L_{F,\varphi}\|_{B(L^p(G))} \leq \frac{1}{c_\varphi} \|\varphi\|_{L^p(G)} \|\varphi\|_{L^{p'}(G)} \|F\|_{L^1(G)},$$

where $\|\cdot\|_{B(L^p(G))}$ is the norm in $B(L^p(G))$.

Proof. By (2.8), Hölder's inequality and the fact that π is the left regular representation of G , we get

$$\begin{aligned} |(L_{F,\varphi}u, v)| &= \frac{1}{c_\varphi} \int_G |F(g)| |(u, \pi(g)\varphi)| |(\pi(g)\varphi, v)| d\mu(g) \\ &\leq \frac{1}{c_\varphi} \|\varphi\|_{L^p(G)} \|\varphi\|_{L^{p'}(G)} \|F\|_{L^1(G)} \|u\|_{L^p(G)} \|v\|_{L^{p'}(G)} \end{aligned}$$

for all u in $L^p(G)$ and v in $L^{p'}(G)$. This completes the proof. □

Proposition 2.3. *Let $F \in L^\infty(G)$. Then for $1 \leq p \leq \infty$, the localization operator $L_{F,\varphi} : L^p(G) \rightarrow L^p(G)$ is a bounded linear operator and*

$$\|L_{F,\varphi}\|_{B(L^p(G))} \leq \frac{1}{c_\varphi} \|\varphi\|_{L^1(G)}^2 \|F\|_{L^\infty(G)}.$$

Proof. By (2.8), we get

$$(2.9) \quad |(L_{F,\varphi}u, v)| = \frac{1}{c_\varphi} \int_G |F(g)| |(u, \pi(g)\varphi)| |(\pi(g)\varphi, v)| d\mu(g)$$

for all u in $L^p(G)$ and v in $L^{p'}(G)$. Now, using the fact that π is the left regular representation of G and Young's inequality on, say, page 52 of the book [7] by Folland again, we get

$$(2.10) \quad \left\{ \int_G |(u, \pi(g)\varphi)|^p d\mu(g) \right\}^{\frac{1}{p}} \leq \|u\|_{L^p(G)} \|\varphi\|_{L^1(G)}$$

and

$$(2.11) \quad \left\{ \int_G |(\pi(g)\varphi, v)|^{p'} d\mu(g) \right\}^{\frac{1}{p'}} \leq \|\varphi\|_{L^1(G)} \|v\|_{L^{p'}(G)}$$

for all u in $L^p(G)$ and v in $L^{p'}(G)$. So, by (2.9), Hölder's inequality, (2.10) and (2.11), we get

$$|(L_{F,\varphi}u, v)| \leq \frac{1}{c_\varphi} \|\varphi\|_{L^1(G)}^2 \|F\|_{L^\infty(G)} \|u\|_{L^p(G)} \|v\|_{L^{p'}(G)}$$

for all u in $L^p(G)$ and v in $L^{p'}(G)$. This completes the proof. □

3. L^p BOUNDEDNESS

We begin with the following result, which is known as the Riesz-Thorin theorem. References on this theorem include [17] by Stein and Weiss and [18] by Wong.

Theorem 3.1. *Let (X, μ) be a measure space and (Y, ν) a σ -finite measure space. Let T be a linear transformation with domain \mathcal{D} consisting of all μ -simple functions f on X such that*

$$\mu\{x \in X : f(x) \neq 0\} < \infty$$

and such that the range of T is contained in the set of all ν -measurable functions on Y . Suppose that $\alpha_1, \alpha_2, \beta_1$ and β_2 are real numbers in $[0, 1]$ and there exist positive constants M_1 and M_2 such that

$$\|Tf\|_{L^{\frac{1}{\beta_j}}(Y)} \leq M_j \|f\|_{L^{\frac{1}{\alpha_j}}(X)}, \quad f \in \mathcal{D}, \quad j = 1, 2.$$

Then for $0 < \theta < 1$,

$$\alpha = (1 - \theta)\alpha_1 + \theta\alpha_2$$

and

$$\beta = (1 - \theta)\beta_1 + \theta\beta_2,$$

we have

$$\|Tf\|_{L^{\frac{1}{\beta}}(Y)} \leq M_1^{1-\theta} M_2^\theta \|f\|_{L^{\frac{1}{\alpha}}(X)}, \quad f \in \mathcal{D}.$$

Theorem 3.2. *Let $F \in L^r(G)$, $1 \leq r \leq \infty$. Then for $1 \leq p \leq \infty$, there exists a unique bounded linear operator $L_{F,\varphi} : L^p(G) \rightarrow L^p(G)$ such that formula (2.2) is valid for all u in $L^p(G)$, v in $L^{p'}(G)$ and μ -simple functions F on G for which*

$$\mu\{g \in G : F(g) \neq 0\} < \infty.$$

Moreover,

$$\|L_{F,\varphi}\|_{B(L^p(G))} \leq \frac{1}{c_\varphi} \|\varphi\|_{L^p(G)}^{\frac{1}{r}} \|\varphi\|_{L^{p'}(G)}^{\frac{1}{r}} \|\varphi\|_{L^1(G)}^{\frac{2}{r}} \|F\|_{L^r(G)}.$$

Proof. We only need to prove the theorem for $1 < r < \infty$. To this end, let $u \in L^p(G)$. Let T_u be the linear transformation with domain \mathcal{D} consisting of all μ -simple functions on G with the property that

$$\mu\{g \in G : F(g) \neq 0\} < \infty,$$

and defined by

$$T_u F = L_{F,\varphi} u, \quad u \in \mathcal{D}.$$

Then by Proposition 2.2,

(3.1)

$$\|T_u F\|_{L^p(G)} = \|L_{F,\varphi} u\|_{L^p(G)} \leq \frac{1}{c_\varphi} \|\varphi\|_{L^p(G)} \|\varphi\|_{L^{p'}(G)} \|u\|_{L^p(G)} \|F\|_{L^1(G)},$$

and by Proposition 2.3,

$$(3.2) \quad \|T_u F\|_{L^p(G)} = \|L_{F,\varphi} u\|_{L^p(G)} \leq \frac{1}{c_\varphi} \|\varphi\|_{L^1(G)}^2 \|u\|_{L^p(G)} \|F\|_{L^\infty(G)}$$

for all F in \mathcal{D} . In order to apply the Riesz-Thorin theorem, we let $\alpha_1 = 1$, $\alpha_2 = 0$ and $\beta_1 = \beta_2 = \frac{1}{p}$. Let $\alpha = \frac{1}{r}$. Then $\theta = \frac{1}{r'}$, where r' is the conjugate index of r . Hence $\alpha = \frac{1}{r}$ and $\beta = \frac{1}{p}$, and we get by (3.1), (3.2) and the Riesz-Thorin theorem,

$$\begin{aligned} \|L_{F,\varphi} u\|_{L^p(G)} &= \|T_u F\|_{L^p(G)} \\ &\leq \frac{1}{c_\varphi} \|\varphi\|_{L^p(G)}^{\frac{1}{r}} \|\varphi\|_{L^{p'}(G)}^{\frac{1}{r}} \|\varphi\|_{L^1(G)}^{\frac{2}{r}} \|F\|_{L^r(G)} \|u\|_{L^p(G)} \end{aligned}$$

for all F in \mathcal{D} . Since \mathcal{D} is dense in $L^r(G)$, we can use a density argument and the proof is complete. \square

4. WAVELET MULTIPLIERS

For $1 \leq p \leq \infty$, we let $\pi : \mathbb{R}^n \rightarrow B(L^p(\mathbb{R}^n))$ be the left regular representation of the additive group \mathbb{R}^n on $L^p(\mathbb{R}^n)$, i.e.,

$$(\pi(y)u)(x) = u(x - y), \quad x, y \in \mathbb{R}^n.$$

Let $\varphi \in \bigcap_{1 \leq p \leq \infty} L^p(\mathbb{R}^n)$ be such that $\|\varphi\|_{L^2(\mathbb{R}^n)} = 1$. Then, by Plancherel's theorem,

$$(4.1) \quad \begin{aligned} c_\varphi &= \int_{\mathbb{R}^n} |\langle \varphi, \pi(y)\varphi \rangle|^2 dy \\ &= \int_{\mathbb{R}^n} |\langle \hat{\varphi}, M_y \hat{\varphi} \rangle|^2 dy, \end{aligned}$$

where

$$(4.2) \quad (M_y \hat{\varphi})(\xi) = e^{iy \cdot \xi} \hat{\varphi}(\xi), \quad \xi \in \mathbb{R}^n.$$

So, by (4.1), (4.2) and Plancherel's theorem,

$$(4.3) \quad \begin{aligned} c_\varphi &= \int_{\mathbb{R}^n} \left| \int_{\mathbb{R}^n} e^{-iy \cdot \xi} |\hat{\varphi}(\xi)|^2 d\xi \right|^2 dx \\ &= (2\pi)^n \|\hat{\varphi}\|_{L^4(\mathbb{R}^n)}^4. \end{aligned}$$

The following result follows immediately from Theorem 3.2.

Theorem 4.1. *Let $\sigma \in L^r(\mathbb{R}^n)$, $1 \leq r \leq \infty$, and let $\varphi \in \bigcap_{1 \leq p \leq \infty} L^p(\mathbb{R}^n)$ be such that $\|\varphi\|_{L^2(G)} = 1$. Then for $1 \leq p \leq \infty$, there exists a unique localization operator $L_{\sigma, \varphi} : L^p(\mathbb{R}^n) \rightarrow L^p(\mathbb{R}^n)$ such that*

$$\langle L_{\sigma, \varphi} u, v \rangle = (2\pi)^{-n} \|\hat{\varphi}\|_{L^4(\mathbb{R}^n)}^{-4} \int_{\mathbb{R}^n} \sigma(y) \langle u, \pi(y)\varphi \rangle \langle \pi(y)\varphi, v \rangle dy$$

for all u and v in \mathcal{S} . Moreover,

$$\|L_{\sigma, \varphi}\|_{B(L^p(\mathbb{R}^n))} \leq (2\pi)^{-n} \|\hat{\varphi}\|_{L^4(\mathbb{R}^n)}^{-4} \|\varphi\|_{L^p(\mathbb{R}^n)}^{\frac{1}{r}} \|\varphi\|_{L^{p'}(\mathbb{R}^n)}^{\frac{1}{r}} \|\varphi\|_{L^1(\mathbb{R}^n)}^{\frac{2}{r}} \|\sigma\|_{L^r(\mathbb{R}^n)}.$$

Now, for all σ in $L^r(\mathbb{R}^n)$, $1 \leq r \leq \infty$, and u and v in the Schwartz space \mathcal{S} , we get by (2.2), (4.3), and Plancherel's theorem,

$$\begin{aligned} \langle L_{\sigma, \varphi} u, v \rangle &= \frac{1}{c_\varphi} \int_{\mathbb{R}^n} \sigma(y) \langle u, \pi(y)\varphi \rangle \langle \pi(y)\varphi, v \rangle dy \\ &= (2\pi)^{-n} \|\hat{\varphi}\|_{L^4(\mathbb{R}^n)}^{-4} \int_{\mathbb{R}^n} \sigma(y) \langle \hat{u}, M_y \hat{\varphi} \rangle \langle M_y \hat{\varphi}, \hat{v} \rangle dy \\ &= \|\hat{\varphi}\|_{L^4(\mathbb{R}^n)}^{-4} \int_{\mathbb{R}^n} \sigma(y) \widehat{(\hat{u}\hat{\varphi})}(y) \overline{\widehat{(\hat{v}\hat{\varphi})}(y)} dy \\ &= (2\pi)^{\frac{n}{2}} \langle T_\sigma(\hat{\varphi}\hat{u}), \hat{\varphi}\hat{v} \rangle \\ (4.4) \quad &= (2\pi)^{\frac{n}{2}} \|\hat{\varphi}\|_{L^4(\mathbb{R}^n)}^{-4} \langle \mathcal{F}^{-1} \overline{\hat{\varphi}} T_\sigma \hat{\varphi} \mathcal{F} u, v \rangle. \end{aligned}$$

Thus, in the case when $p = 2$, (4.4) tells us that the localization operator $L_{\sigma, \varphi} : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$ is unitarily equivalent to the wavelet multiplier $P_{\sigma, \varphi} : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$.

REFERENCES

- [1] A. L. Carey, Square integrable representations of non-unimodular groups, *Bull. Austral. Math. Soc.* **15** (1976), 1-12. MR **55**:3153
- [2] I. Daubechies, Time-frequency localization operators: a geometric phase space approach, *IEEE Trans. Inform. Theory* **34** (1988), 605-612. CMP 21:03
- [3] I. Daubechies, *Ten Lectures on Wavelets*, SIAM, 1992. MR **93e**:42045
- [4] M. Dufo and C. C. Moore, On the regular representation of a nonunimodular locally compact group, *J. Funct. Anal.* **21** (1976), 209-243. MR **52**:14145
- [5] J. Du and M. W. Wong, Traces of localization operators, *C. R. Math. Rep. Acad. Sci. Canada* **22** (2000), 92-95. MR **2001c**:47053
- [6] J. Du and M. W. Wong, Traces of wavelet multipliers, *C.R. Math. Acad. Sci. Soc. R. Can.* **23** (2001), 148-152.
- [7] G. B. Folland, *A Course in Abstract Harmonic Analysis*, CRC Press, 1995. MR **98c**:43001
- [8] A. Grossmann, J. Morlet and T. Paul, Transforms associated to square integrable group representations I: general results, *J. Math. Phys.* **26** (1985), 2473-2479. MR **86k**:22013
- [9] Z. He and M. W. Wong, Localization operators associated to square integrable group representations, *Panamer Math. J.* **6**(1) (1996), 93-104. MR **97d**:43009
- [10] Z. He and M. W. Wong, Wavelet multipliers and signals, *J. Austral. Math. Soc. Ser. B* **40** (1999), 437-446. MR **2001a**:42040
- [11] H. Kumano-go, *Pseudo-Differential Operators*, MIT Press, 1981. MR **84c**:35113
- [12] H. J. Landau and H. O. Pollak, Prolate spheroidal wave functions, Fourier analysis and uncertainty II, *Bell Syst. Tech. J.* **40** (1961), 65-84. MR **25**:4147
- [13] H. J. Landau and H. O. Pollak, Prolate spheroidal wave functions, Fourier analysis and uncertainty III, *Bell Syst. Tech. J.* **41** (1962), 1295-1336. MR **26**:5200
- [14] D. Slepian, On bandwidth, *Proc. IEEE* **64** (1976), 292-300. MR **57**:2738
- [15] D. Slepian, Some comments on Fourier analysis, uncertainty and modeling, *SIAM Rev.* **25** (1983), 379-393. MR **84i**:94016

- [16] D. Slepian and H. O. Pollak, Prolate spheroidal wave functions, Fourier analysis and uncertainty I, Bell Syst. Tech. J. **40** (1961), 43-64. MR **25**:4146
- [17] E. M. Stein and G. Weiss, Introduction to Fourier Analysis on Euclidean Spaces, Princeton University Press, 1971. MR **46**:4102
- [18] M. W. Wong, Weyl Transforms, Springer-Verlag, 1998. MR **2000c**:47098
- [19] M. W. Wong, An Introduction to Pseudo-Differential Operators, Second Edition, World Scientific, 1999. MR **2000c**:35002
- [20] M. W. Wong, Localization Operators, Lecture Notes Series **47**, Seoul National University, Research Institute of Mathematics, Global Analysis Research Center, Seoul, 1999. MR **2002c**:22010

DEPARTMENT OF MATHEMATICS AND STATISTICS, YORK UNIVERSITY, 4700 KEELE STREET,
TORONTO, ONTARIO, CANADA M3J 1P3

E-mail address: `mwwong@pascal.math.yorku.ca`