

## TRIQUOTIENT MAPS VIA ULTRAFILTER CONVERGENCE

MARIA MANUEL CLEMENTINO AND DIRK HOFMANN

(Communicated by Alan Dow)

ABSTRACT. In this paper we characterize triquotient maps as those that are surjective on chains of convergent ultrafilters, extending the result known for triquotient maps between finite topological spaces.

### 1. INTRODUCTION

Triquotient maps, introduced by Michael [11], fit very nicely among classes of special quotient maps:

- proper maps and open maps are triquotient maps;
- triquotient maps are effective descent, which in turn are biquotient maps.

A recent study of Janelidze and Sobral on the behaviour of the mentioned classes of morphisms, when defined between finite topological spaces, led to very interesting characterizations based on point convergence. Among these characterizations, the following was established (see [3] and [7]):

**Theorem I.** *If  $X$  and  $Y$  are finite topological spaces, a continuous map  $f : X \rightarrow Y$  is a triquotient map if and only if it is surjective on chains of convergent points.*

To pass from the finite to the infinite case one must replace points by (ultra)filters – or (ultra)nets – and, except for the cases of effective descent and triquotient maps (and partially local homeomorphisms), the characterizations are straightforward. To establish a general characterization of triquotient maps that includes the former theorem, some new notions and techniques are needed.

This is the central part of this paper: we introduce and study a new category, defined via ultrafilter convergence, endowed with a special endofunctor that is used to define *chains of convergent ultrafilters*, and that finally leads to

**Theorem II.** *A continuous map  $f : X \rightarrow Y$  is a triquotient map if and only if it is surjective on chains of convergent ultrafilters.*

Moreover, these characterizations turn out to be very effective for proving stability properties for special kinds of limits, since initial structures – in particular, limit

---

Received by the editors November 7, 2000 and, in revised form, June 25, 2001.

2000 *Mathematics Subject Classification.* Primary 54C10, 54A20, 54B30, 18A20, 18B30.

*Key words and phrases.* Biquotient map, effective descent map and triquotient map, convergent structure.

The authors acknowledge partial financial assistance by Centro de Matemática da Universidade de Coimbra. The first author also thanks Project PRAXIS XXI 2/2.1/MAT/46/94.

structures – are easily described by convergent ultrafilters. This gives rise to unified proofs of results obtained separately, and will be the subject of a forthcoming note.

2. BASIC DEFINITIONS AND RESULTS

For a topological space  $X$  we denote its topology by  $\mathcal{O}X$ . For  $x \in X$ ,  $\mathcal{O}(x)$  denotes the set of open subsets of  $X$  containing  $x$ .

**Definitions 2.1.** A topological continuous map  $f : X \rightarrow Y$  is:

1. a *biquotient map* if, whenever  $y \in Y$  and  $\mathcal{A}$  is an open covering of  $f^{-1}(y)$ , then finitely many  $f(A)$ , with  $A \in \mathcal{A}$ , cover some neighbourhood of  $y$  in  $Y$ ;
2. *effective descent (descent)* if the pullback functor  $f^* : \mathbf{Top}/Y \rightarrow \mathbf{Top}/X$ , that assigns to each  $g : Z \rightarrow Y$  its pullback along  $f$ , is (pre)monadic (see [8]);
3. a *triquotient map* if there exists a map  $(-)^{\#} : \mathcal{O}X \rightarrow \mathcal{O}Y$  such that
  - (T1)  $(\forall U \in \mathcal{O}X) U^{\#} \subseteq f(U)$ ,
  - (T2)  $X^{\#} = Y$ ,
  - (T3)  $(\forall U, V \in \mathcal{O}X) U \subseteq V \Rightarrow U^{\#} \subseteq V^{\#}$ ,
  - (T4)  $(\forall U \in \mathcal{O}X) (\forall y \in U^{\#}) (\forall \Sigma \subseteq \mathcal{O}X \text{ directed}) f^{-1}(y) \cap U \subseteq \bigcup \Sigma \Rightarrow \exists S \in \Sigma : y \in S^{\#}$ ;
4. *proper (perfect)* if it is closed and has compact fibres (and Hausdorff, i.e. if  $f(x) = f(x')$  and  $x \neq x'$  there exist  $U \in \mathcal{O}(x)$  and  $V \in \mathcal{O}(x')$  with  $U \cap V = \emptyset$ ).


Concerning the notion of triquotient map, we note that (T3) is implied by (T4).

We also remark that every proper map  $f : X \rightarrow Y$  is triquotient: take  $U^{\#} := Y - f(X - U)$  for  $U \in \mathcal{O}X$ , as well as every open map: take  $U^{\#} := f(U)$ . Plewe showed that – both on topological spaces and locales – triquotient maps are effective descent (see [12]). These latter are descent maps, which are exactly the biquotient maps (see [8]) introduced independently by Michael [10], Hájek [5], by the name of *limit lifting maps*, and Day and Kelly [4], as *universal quotient maps*.

The characterizations of biquotient, open, proper and perfect maps between finite topological spaces stated in [7] can be easily generalized to arbitrary spaces using ultrafilters, while a possible generalization of the characterization of effective descent maps is the well-known Reiterman-Tholen characterization [13]:

**Theorem 2.2.** *If  $f : X \rightarrow Y$  is a continuous map, then:*


1.  *$f$  is a biquotient map if and only if, for each ultrafilter  $\mathfrak{b} \rightarrow y$  in  $Y$ , there exists an ultrafilter  $\mathfrak{a} \rightarrow x$  in  $X$  with  $f(\mathfrak{a}) = \mathfrak{b}$  and  $f(x) = y$ :*




2.  *$f$  is an effective descent map if and only if, for each family of ultrafilters  $(\mathfrak{b}_i)_{i \in I}$  and each ultrafilter  $\mathfrak{u}$  on  $I$ , whenever  $\mathfrak{b}_i \rightarrow y_i$ , for  $i \in I$ , and  $y_i \xrightarrow{\mathfrak{u}} y$  in  $Y$ , there exists an ultrafilter  $\mathfrak{a}$  on  $X$  such that  $\mathfrak{a} \rightarrow x \in f^{-1}(y)$  and, for each  $U \in \mathfrak{u}$ ,*

$$\bigcup_{i \in U} (f^{-1}(y_i) \cap \text{adh}(f^{-1}(\mathfrak{b}_i))) \in \mathfrak{a}.$$

3.  $f$  is a proper (perfect) map if and only if, for each ultrafilter  $\mathfrak{a}$  on  $X$  with  $f(\mathfrak{a}) \rightarrow y$  in  $Y$ , there exists (a unique)  $x \in f^{-1}(y)$  such that  $\mathfrak{a} \rightarrow x$ :



4.  $f$  is an open map if and only if, for each  $x \in X$  and each ultrafilter  $\mathfrak{b}$  on  $Y$  with  $\mathfrak{b} \rightarrow f(x)$  in  $Y$ , there exists an ultrafilter  $\mathfrak{a}$  such that  $\mathfrak{a} \rightarrow x$  in  $X$  and  $f(\mathfrak{a}) = \mathfrak{b}$ :



By similarity with the finite case, and also with the characterizations of proper and perfect maps, it seems natural that one may obtain a characterization of local homeomorphisms imposing in the one above a uniqueness condition. We do not know if this really holds. In fact, what we know at the moment is:

**Theorem 2.3.** *For a continuous map  $f : X \rightarrow Y$ , if  $f$  is a local homeomorphism, then, for each  $x \in X$  and each ultrafilter  $\mathfrak{b}$  on  $Y$  with  $\mathfrak{b} \rightarrow f(x)$  in  $Y$ , there exists a unique ultrafilter  $\mathfrak{a}$  such that  $\mathfrak{a} \rightarrow x$  in  $X$  and  $f(\mathfrak{a}) = \mathfrak{b}$ .*

It is also possible to characterize these classes of maps by their behaviour on lifting convergent (ultra)nets:

**Theorem 2.4.** *Let  $f : X \rightarrow Y$  be a topological continuous map.*

1.  $f$  is a biquotient map if and only if, for each net  $y_\lambda \rightarrow y$  in  $Y$ , there exists a net  $x_\gamma \rightarrow x$  in  $X$  such that  $(f(x_\gamma))$  is a subnet of  $(y_\lambda)$  and  $f(x) = y$ .
2.  $f$  is a proper (perfect) map if and only if, for each ultranet  $(x_\lambda)_{\lambda \in \Lambda}$  in  $X$  with  $f(x_\lambda) \rightarrow y$  in  $Y$ , there exists (a unique)  $x \in f^{-1}(y)$  such that  $x_\lambda \rightarrow x$ .
3.  $f$  is an open map if and only if, for each  $x \in X$  and each ultranet  $(y_\lambda)_{\lambda \in \Lambda}$  converging to  $f(x)$  in  $Y$ , there exists  $(x_\gamma)_{\gamma \in \Gamma}$  in  $X$  and  $\phi : \Gamma \rightarrow \Lambda$  such that  $x_\gamma \rightarrow x$  and  $(f(x_\gamma))$  is a subnet of  $(y_\lambda)$ , via  $\phi$ , such that  $\phi(\uparrow \gamma) = \uparrow \phi(\gamma)$ .
4.  $f$  is a local homeomorphism if and only if, for each  $x \in X$  and each net  $y_\lambda \rightarrow f(x)$  in  $Y$ , there exists an essentially unique net  $(x_\lambda)_{\lambda \in \Lambda}$  in  $X$  such that  $x_\lambda \rightarrow x$  and  $f(x_\lambda) = y_\lambda$  for every  $\lambda \in \Lambda$ .

(By *essentially unique* we mean that, if  $(x_\lambda)$  and  $(x'_\lambda)$  satisfy the conditions above, then there exists  $\lambda_0 \in \Lambda$  such that, for  $\lambda \geq \lambda_0$ ,  $x_\lambda = x'_\lambda$ .)

The approach with ultrafilter convergence gives a more elegant and unified way of describing biquotient, proper, perfect and open maps. This is the reason why we preferred them to nets, and investigated similar characterizations for effective descent and triquotient maps. This is the aim of the following sections.

First we recall some known facts on ultrafilters. If  $X$  is a set, the set  $\mathcal{U}(X)$  of ultrafilters on  $X$  may be endowed with the Zariski closure, becoming a compact Hausdorff space (it is in fact the Čech-Stone compactification of the discrete space  $X$ ; see [9]). For a map  $f : X \rightarrow Y$  and  $\mathfrak{a} \in \mathcal{U}(X)$ ,  $f(\mathfrak{a})$  denotes the filter generated by  $\{f(A) \mid A \in \mathfrak{a}\}$ , which is automatically an ultrafilter since  $\mathfrak{a}$  is. The map  $\mathcal{U}f : \mathcal{U}(X) \rightarrow \mathcal{U}(Y)$ ,  $\mathfrak{a} \mapsto f(\mathfrak{a})$  is continuous.

3. THE CATEGORY **URS**

In order to characterize triquotient maps using convergence, we will need a combination of ultrafilters convergence, as it is already the case of effective descent maps, but in a higher order: 2-chains for effective descent maps between finite spaces give rise to the combination of 2-sorts of convergent ultrafilters while  $n$ -chains, for triquotient maps, will give rise to infinite chains of convergent ultrafilters.

To make the description as simple as possible, we introduce the category of ultrarelatational spaces, whose particularity – that distinguishes them from pseudotopological spaces – is the fact that principal ultrafilters do not need to converge.

**Definition 3.1.** An *ultrarelation* on a set  $X$  is a subset  $r \subseteq \mathcal{U}(X) \times X$ . An *ultrarelatational space* is a set  $X$  equipped with an ultrarelation  $r$  on  $X$ . Given ultrarelatational spaces  $(X, r)$  and  $(Y, s)$ , a map  $f : X \rightarrow Y$  is *continuous* if, for each  $(\mathbf{a}, x) \in r$ ,  $(f(\mathbf{a}), f(x)) \in s$ .

We denote by **URS** the category of ultrarelatational spaces and continuous maps.

We will often use the more suggestive notation  $\mathbf{a} \rightarrow x$  instead of  $(\mathbf{a}, x) \in r$ .

The category **URS** is equipped with a canonical faithful functor  $|\_|\_ : \mathbf{URS} \rightarrow \mathbf{Set}$  sending  $(X, r)$  to  $X$ . The construct  $(\mathbf{URS}, |\_|\_)$  is topological (in the sense of [1]) and therefore concretely complete and cocomplete. It contains **Top** as a full and concrete subcategory: each topology  $\tau$  on  $X$  defines an ultrarelation  $r_{(X, \tau)}$  by

$$r_{(X, \tau)} = \{(\mathbf{a}, x) \mid \mathbf{a} \rightarrow x \text{ w.r.t. the topology } \tau\}.$$

For an ultrarelatational space  $(X, r)$ , we consider the projection map  $p_{(X, r)} : r \rightarrow X$ , with  $(\mathbf{a}, x) \mapsto x$ , and define the following ultrarelation  $R_{(X, r)}$  on  $r$ :

$$R_{(X, r)} = \{(\mathfrak{A}, (\mathbf{a}, x)) \in \mathcal{U}(r) \times r \mid p_{(X, r)}(\mathfrak{A}) = \mathbf{a}\}.$$

We denote the ultrarelatational space  $(r, R_{(X, r)})$  by  $\text{Ult}(X, r)$ . Obviously, the map  $p_{(X, r)} : \text{Ult}(X, r) \rightarrow (X, r)$  is continuous, by definition of the structure on  $r$ .

Some extra conditions on these spaces will give us back well-known structures:

**Definition 3.2.** An ultrarelatational space  $(X, r)$  is called

1. *weak reflexive* if, for each  $x \in X$ , there exists an  $\mathbf{a} \in \mathcal{U}(X)$  such that  $(\mathbf{a}, x) \in r$ ;
2. *reflexive* if, for each  $x \in X$ ,  $(\dot{x}, x) \in r$ ;
3. *fibre-closed* if, for each  $x \in X$ ,  $\{\mathbf{a} \in \mathcal{U}(X) \mid (\mathbf{a}, x) \in r\}$  is closed in  $\mathcal{U}(X)$  with respect to the Zariski topology;
4. *transitive* if the map

$$\mu_{(X, r)} : R_{(X, r)} \rightarrow \mathcal{U}(X) \times X, (\mathfrak{A}, (\mathbf{a}, x)) \mapsto \left( \bigcup_{\mathcal{A} \in \mathfrak{A}} \bigcap_{(\mathbf{a}', x') \in \mathcal{A}} \mathbf{a}', x \right)$$

factors via the inclusion  $r \hookrightarrow \mathcal{U}(X) \times X$ .

An ultrarelatational space  $(X, r)$  is weak reflexive if and only if  $p_{(X, r)} : \text{Ult}(X, r) \rightarrow (X, r)$  is surjective. Hence  $\text{Ult}(X, r)$  is weak reflexive provided that  $(X, r)$  is: for an  $(\mathbf{a}, x)$ ,  $p_{(X, r)}^{-1}(\mathbf{a})$  is a filter base and any ultrafilter  $\mathfrak{A}$  in  $\text{Ult}(X, r)$  containing it converges to  $(\mathbf{a}, x)$ . Moreover,  $\text{Ult}(X, r)$  is always fibre-closed. However,  $\text{Ult}$  does not preserve reflexivity or transitivity.

These properties define full subcategories of **URS**: the category **PsTop** (**PrTop**; **Top**) of pseudotopological (pretopological; topological) spaces is concretely isomorphic to the full subcategory of **URS** consisting of all reflexive (reflexive and fibre-closed; reflexive and transitive) ultrarelatational spaces.

4. THE FUNCTOR  $\text{Ult}$

Each ultrarelatational continuous map  $f : (X, r) \rightarrow (Y, s)$  induces a map

$$\text{Ult}(f) : r \rightarrow s, (\mathbf{a}, x) \mapsto (f(\mathbf{a}), f(x)),$$

and the diagram

$$\begin{array}{ccc} r & \xrightarrow{\text{Ult}(f)} & s \\ p_{(X,r)} \downarrow & & \downarrow p_{(Y,s)} \\ X & \xrightarrow{f} & Y \end{array}$$

commutes. It is then clear that  $\text{Ult}(f) : \text{Ult}(X, r) \rightarrow \text{Ult}(Y, s)$  is continuous. Moreover, the equalities  $\text{Ult}(1) = 1$  and  $\text{Ult}(f \circ g) = \text{Ult}(f) \circ \text{Ult}(g)$  hold, hence  $\text{Ult} : \mathbf{URS} \rightarrow \mathbf{URS}$  is a functor and  $(p_{(X,r)})_{(X,r) \in \text{Ob } \mathbf{URS}} : \text{Ult} \rightarrow \text{Id}_{\mathbf{URS}}$  is a natural transformation.

Note that we have  $\text{Ult}(p_{(X,r)}) = p_{\text{Ult}(X,r)}$  for each ultrarelatational space  $(X, r)$ , that is,  $(\text{Ult}, p)$  is a *well copointed endofunctor* (see [6]).

Hence, we may define endofunctors  $\text{Ult}^\alpha$  and natural transformations  $p_\beta^\alpha$  for ordinal numbers  $\alpha, \beta$  with  $\beta \leq \alpha$ , by:

- $\text{Ult}^0 = \text{Id}_{\mathbf{URS}}, p_0^0 = 1_{\text{Ult}^0}$ ;
- $\text{Ult}^{\alpha+1} = \text{Ult}(\text{Ult}^\alpha), p_\beta^{\alpha+1} = p_\beta^\alpha \cdot p_{\text{Ult}^\alpha}$  and  $p_{\alpha+1}^{\alpha+1} = 1_{\text{Ult}^{\alpha+1}}$ , for  $\beta \leq \alpha$ ;
- $\text{Ult}^\lambda = \lim_{\beta \leq \alpha < \lambda} p_\beta^\alpha, p_\beta^\lambda =$  the limit projection and  $p_\lambda^\lambda = 1_{\text{Ult}^\lambda}$ , for every limit ordinal  $\lambda$  and every  $\beta < \lambda$ .

From now on, since we usually work with only one ultrarelation on a set  $X$ , for an ultrarelatational space we relax our notation and write  $X$  instead of  $(X, r)$ . Also, we will denote  $\text{Ult}^\alpha(X)$  by  $X_\alpha$  and  $\text{Ult}^\alpha(f)$  by  $f_\alpha$ , for every continuous map  $f : X \rightarrow Y$  between ultrarelatational spaces.

This transfinite construction can be easily described: for each ultrarelatational space  $X$  and each ordinal  $\alpha$ ,

$$X_\alpha = \{((\mathbf{a}_\beta)_{\beta \in \alpha}, x) \in \prod_{\beta \in \alpha} \mathcal{U}(X_\beta) \times X \mid \mathbf{a}_0 \rightarrow x \text{ and } (\forall \gamma \leq \beta < \alpha) (p_\gamma^\beta)_X(\mathbf{a}_\beta) = \mathbf{a}_\gamma\},$$

for each  $\beta \leq \alpha$ , the projection  $(p_\beta^\alpha)_X : X_\alpha \rightarrow X_\beta$  is defined by

$$(p_\beta^\alpha)_X((\mathbf{a}_\gamma)_{\gamma \in \alpha}, x) = ((\mathbf{a}_\gamma)_{\gamma \in \beta}, x),$$

and the ultrarelatational structure in  $X_\alpha$  is defined by

$$\mathbf{a}_\alpha \rightarrow ((\mathbf{a}_\beta)_{\beta \in \alpha}, x) \iff (\forall \beta \in \alpha) (p_\beta^\alpha)_X(\mathbf{a}_\alpha) = \mathbf{a}_\beta.$$

Finally, if  $f : X \rightarrow Y$  is a continuous map, then, for each ordinal  $\alpha$  and each  $((\mathbf{a}_\beta)_{\beta \in \alpha}, x) \in X_\alpha$ ,  $f_\alpha((\mathbf{a}_\beta)_{\beta \in \alpha}, x) = ((f_\beta(\mathbf{a}_\beta))_{\beta \in \alpha}, f(x))$ .

We remark that, for each ordinal  $\alpha$  and each ultrarelatational space  $X$ , an element of  $X_{\alpha+1}$  is given by an ultrafilter  $\mathbf{a}_\alpha \in \mathcal{U}(X_\alpha)$  and an element  $x \in X$  such that  $(p_0^\alpha)_X(\mathbf{a}_\alpha) \rightarrow x$ . The map  $f_{\alpha+1} : X_{\alpha+1} \rightarrow Y_{\alpha+1}$  is surjective if and only if, for each ultrafilter  $\mathbf{b}_\alpha$  on  $Y_\alpha$  and each  $y \in Y$  such that  $(p_0^\alpha)_Y(\mathbf{b}_\alpha) \rightarrow y$ , there exist an ultrafilter  $\mathbf{a}_\alpha$  on  $X_\alpha$  and an  $x \in f^{-1}(y)$  such that  $(p_0^\alpha)_X(\mathbf{a}_\alpha) \rightarrow x$  and  $f_\alpha(\mathbf{a}_\alpha) = \mathbf{b}_\alpha$ .

Hence, by Theorem 2.2, if  $X$  and  $Y$  are topological spaces, a continuous map  $f : X \rightarrow Y$  is a biquotient map if and only if  $f_1$ , and then also  $f_0$ , is surjective.

Next we will show that this kind of surjectivity condition also characterizes effective descent and triquotient maps in **Top**. Therefore, we introduce the following

**Definition 4.1.** If  $\alpha$  is an ordinal number, an ultrarelatational continuous map  $f : X \rightarrow Y$  is said to be  $\alpha$ -surjective if  $f_\beta : X_\beta \rightarrow Y_\beta$  is surjective for every  $\beta \in \alpha$ . The map  $f$  is called  $\Omega$ -surjective if  $f_\alpha$  is surjective for every ordinal  $\alpha$ .

Hence, 1-surjective maps are just surjective maps, while our observation above means that a biquotient map in **Top** is a 2-surjective map.

5. 3-SURJECTIVE MAPS

The continuous maps  $f : X \rightarrow Y$  between topological spaces such that  $f_2$  (and then also  $f_0$  and  $f_1$ ) is surjective are very well-known: they are exactly the effective descent maps in **Top**, as we show below. For that we will make use of Reiterman-Tholen characterization (Theorem 2.2). We first start showing that the data they used may be easily interpreted using the functor **Ult**.

**Lemma 5.1.** *If  $Y$  is a topological space and  $\mathcal{F}_Y = \{(I, \mathbf{u}, (f_i), (y_i), y) \mid \mathbf{u} \text{ ultrafilter on } I, f_i \rightarrow y_i \text{ and } y_i \xrightarrow{\mathbf{u}} y\}$ , there are maps  $\Phi : Y_2 \rightarrow \mathcal{F}_Y$  and  $\Psi : \mathcal{F}_Y \rightarrow Y_2$  such that  $\Psi \cdot \Phi = 1_{Y_2}$ .*

*Proof.* For any  $(\mathfrak{B}, (\mathbf{b}, y))$  in  $Y_2$ ,  $\mathfrak{B}$  is an ultrafilter on  $Y_1$  and, for  $\phi = p : Y_1 \rightarrow Y$ ,

$$\Phi(\mathfrak{B}, (\mathbf{b}, y)) := (Y_1, \mathfrak{B}, (f)_{(f, y') \in Y_1}, (y')_{(f, y') \in Y_1}, y)$$

belongs to  $\mathcal{F}_Y$  since  $(f, y') \in Y_1$ , that is  $f \rightarrow y'$ , and  $y' \xrightarrow{\mathfrak{B}} y$  by the definition of the ultrarelatational structure on  $Y_1$ .

On the other hand, if  $(I, \mathbf{u}, (f_i), (y_i), y) \in \mathcal{F}_Y$ , with  $\phi : I \rightarrow Y$  inducing  $y_i \xrightarrow{\mathbf{u}} y$ , for  $\psi : I \rightarrow Y_1$  defined by  $\psi(i) = (f_i, y_i)$ , we may define

$$\Psi(I, \mathbf{u}, (f_i), (y_i), y) := (\psi(\mathbf{u}), (\phi(\mathbf{u}), y)),$$

and it is easy to check that  $\Psi \cdot \Phi = 1_{Y_2}$ . □

**Theorem 5.2.** *A topological continuous map  $f : X \rightarrow Y$  is effective descent if and only if it is 3-surjective, that is:*

$$\begin{array}{ccccc} X & & \mathfrak{A} & \dashrightarrow & \mathfrak{a} & \dashrightarrow & x \\ f \downarrow & & \downarrow & & \downarrow & & \downarrow \\ Y & & \mathfrak{B} & \longrightarrow & \mathbf{b} & \longrightarrow & y \end{array}$$

*Proof.* Assume first that  $f_2$  is surjective and let  $I$  be an index set, let  $\mathbf{b}_i$  ( $i \in I$ ) be a family of ultrafilters on  $Y$  converging to  $y_i$  and  $y_i \xrightarrow{\mathbf{u}} y$  with  $\mathbf{u}$  ultrafilter on  $I$ . Considering its corresponding element  $(\mathfrak{B}, (\mathbf{b}, y))$  in  $Y_2$ , since  $f_2$  is surjective, there exist an element  $x \in f^{-1}(y)$  and ultrafilters  $\mathfrak{a}$  on  $X$  and  $\mathfrak{A}$  on  $X_1$  such that

$$\mathfrak{a} \rightarrow x, \mathfrak{A} \rightarrow (\mathfrak{a}, x), f_1(\mathfrak{A}) = \mathfrak{B}.$$

Hence we have, for each  $U \in \mathbf{u}$ ,

$$\bigcup_{i \in U} (f^{-1}(y_i) \cap \text{adh}(f^{-1}(\mathbf{b}_i))) = p_X(\text{Ult}(f)^{-1}(\psi(U))) \in \mathfrak{a}.$$

Assume now that  $f$  is effective descent. Let  $(\mathfrak{B}, (\mathfrak{b}, y)) \in Y_2$ . For its corresponding data  $(I, \mathbf{u}, (f_i, y_i, y))$ , since  $f$  is effective descent, there exist an ultrafilter  $\mathfrak{a}$  on  $X$  and an element  $x \in f^{-1}(y)$  such that  $\mathfrak{a} \rightarrow x$  and, for each  $\mathcal{B} \in \mathfrak{B}$ ,

$$p_X(\text{Ult}(f)^{-1}(\mathcal{B})) = \bigcup_{(\mathfrak{b}', y') \in \mathcal{B}} (f^{-1}(y') \cap \text{adh}(f^{-1}(\mathfrak{b}')))) \in \mathfrak{a}.$$

Hence  $\text{Ult}(f)^{-1}(\mathfrak{B}) \cup p_X^{-1}(\mathfrak{a})$  induces a filter on  $X_1$  which can be refined to an ultrafilter  $\mathfrak{A}$ , that clearly satisfies the conditions  $\mathfrak{A} \rightarrow (\mathfrak{a}, x)$  and  $f_1(\mathfrak{A}) = \mathfrak{B}$ .  $\square$

We remark that these techniques also make the proofs of the “Key Lemmas” of [13] become substantially easier.

### 6. $\Omega$ -SURJECTIVE MAPS

We are now going to characterize topological triquotient maps inside **URS** as the  $\Omega$ -surjective maps. First we state an auxiliary result.

**Lemma 6.1.** *Let  $X$  be a weak reflexive ultrarelatational space. Then, for each ordinal  $\alpha$ ,  $X_\alpha$  is weak reflexive and  $(p_\beta^\alpha)_X$  is a surjection for each  $\beta \leq \alpha$ .*

*Proof.* It follows immediately from the preservation of weak reflexivity by  $\text{Ult}$  and from the construction of  $X_\lambda$  for every limit ordinal  $\lambda$ .  $\square$

**Proposition 6.2.** *Let  $f : X \rightarrow Y$  be a topological continuous map together with a map  $(\cdot)^\# : \mathcal{O}X \rightarrow \mathcal{O}Y$  satisfying (T1) and (T4).<sup>1</sup> Then, for each ordinal  $\alpha$  and each  $U \in \mathcal{O}X$ ,  $(p_0^\alpha)^{-1}(U^\#) \subseteq f_\alpha((p_0^\alpha)^{-1}(U))$ .*

*Proof.* For  $\alpha = 0$ , the assertion follows from the fact that  $U^\# \subseteq f(U)$  for each  $U \in \mathcal{O}X$ . For  $\alpha > 0$  assume that the condition above holds for each  $\beta \in \alpha$ . Let  $U \in \mathcal{O}X$ ,  $y \in U^\#$  and  $((\mathfrak{b}_\beta)_{\beta \in \alpha}, y) \in Y_\alpha$ . We define

$$\Sigma = \{S \in \mathcal{O}X \mid \exists \beta \in \alpha : f_\beta((p_0^\beta)_X^{-1}(S)) \notin \mathfrak{b}_\beta\}.$$

$\Sigma$  is directed since all  $\mathfrak{b}_\beta$  ( $\beta \in \alpha$ ) are ultrafilters and all  $(p_\gamma^\beta)_X$  ( $\gamma \leq \beta < \alpha$ ) are surjective. We are now going to show that  $y \notin S^\#$  for each  $S \in \Sigma$ . Assume that  $y \in S^\#$  for some  $S \in \Sigma$ . Then we have  $S^\# \in \mathfrak{b}_0$  and therefore, for all  $\beta \in \alpha$ ,  $(p_0^\beta)_Y^{-1}(S^\#) \in \mathfrak{b}_\beta$ . But this is impossible since, by induction hypothesis, we have

$$(p_0^\beta)_Y^{-1}(S^\#) \subseteq f_\beta((p_0^\beta)_X^{-1}(S)) \notin \mathfrak{b}_\beta.$$

By (T4), there exists  $x \in f^{-1}(y) \cap U$  such that, for all  $S \in \Sigma$ ,  $x \notin S$ . Hence for each  $V \in \mathcal{O}(x)$  and each  $\beta \in \alpha$  we have  $f_\beta((p_0^\beta)_X^{-1}(V)) \in \mathfrak{b}_\beta$  and therefore  $(p_0^\beta)_X^{-1}(\mathcal{O}(x)) \cup f_\beta^{-1}(\mathfrak{b}_\beta)$  induces a filter  $\mathfrak{f}_\beta$  on  $X_\beta$ . For each  $\beta \in \alpha$  we put

$$\mathcal{M}_\beta = \{\mathfrak{a} \in \mathcal{U}(X_\beta) \mid \mathfrak{a} \supseteq \mathfrak{f}_\beta\}.$$

Each  $\mathcal{M}_\beta$  ( $\beta \in \alpha$ ) is non-empty and Zariski-closed, hence, since a codirected limit of non-empty compact Hausdorff spaces is non-empty (cf. [2]), there exists  $(\mathfrak{a}_\beta)_{\beta \in \alpha}$  with  $\mathfrak{a}_\beta$  ultrafilter on  $X_\beta$  such that  $(p_{\beta'}^\beta)_X(\mathfrak{a}_\beta) = \mathfrak{a}_{\beta'}$  for all  $\beta' \leq \beta \in \alpha$ . We have by definition  $\mathfrak{a}_0 \supseteq \mathcal{O}(x)$  and  $f_\beta(\mathfrak{a}_\beta) = \mathfrak{b}_\beta$  for all  $\beta \in \alpha$ , hence  $((\mathfrak{a}_\beta)_{\beta \in \alpha}, x) \in X_\alpha$  and  $f_\alpha((\mathfrak{a}_\beta)_{\beta \in \alpha}, x) = ((\mathfrak{b}_\beta)_{\beta \in \alpha}, y)$ .  $\square$

<sup>1</sup>See Definition 2.1.

Since this shows, in particular, that every triquotient map (between topological spaces) is  $\Omega$ -surjective, we conclude immediately that triquotient maps are effective descent.

For a set  $Y$ , let  $\lambda_Y$  be the least regular cardinal larger than the cardinal of  $Y$ .

**Proposition 6.3.** *Let  $f : X \rightarrow Y$  be a topological continuous map. Then, for each  $U \in \mathcal{O}X$ , the set*

$$(1) \quad U^\sharp = \{y \in Y \mid (\forall \alpha \in \lambda_Y) (p_0^\alpha)^{-1}(y) \subseteq f_\alpha((p_0^\alpha)^{-1}(U))\}$$

*is open and the map  $(-)^\sharp : \mathcal{O}X \rightarrow \mathcal{O}Y$  satisfies (T1) and (T4).*


*Proof.* First we show that  $U^\sharp$  is open for every  $U \in \mathcal{O}X$ . For that, let  $y_0 \in \text{cl}(Y - U^\sharp)$ . There exists an ultrafilter  $\mathfrak{b}_0$  on  $Y$  converging to  $y_0$  such that  $Y - U^\sharp \in \mathfrak{b}_0$ . For each  $y \in Y - U^\sharp$  there exist  $\alpha_y \in \lambda_Y$  and  $((\mathfrak{b}_\beta)_{\beta \in \alpha_y}, y) \in Y_{\alpha_y}$  such that

$$\ddagger((\mathfrak{a}_\beta)_{\beta \in \alpha_y}, x) \in X_{\alpha_y} : (x \in U \wedge f_\alpha((\mathfrak{a}_\beta)_{\beta \in \alpha_y}, x) = ((\mathfrak{b}_\beta)_{\beta \in \alpha_y}, y),$$

by definition of  $U^\sharp$ . For  $\alpha := \sup_{y \in (Y - U^\sharp)} \alpha_y$ , we have  $\alpha \in \lambda_Y$ . Considering  $\mathcal{B} = (p_0^\alpha)^{-1}(Y - U^\sharp) - f_\alpha((p_0^\alpha)^{-1}(U))$ , one has  $(p_0^\alpha)_Y(\mathcal{B}) = Y - U^\sharp \in \mathfrak{b}_0$ . Let  $\mathfrak{b}_\alpha$  be any ultrafilter containing  $\{\mathcal{B}\} \cup (p_0^\alpha)^{-1}(\mathfrak{b}_0)$ . Assume that there exist an ultrafilter  $\mathfrak{a}_\alpha$  on  $X_\alpha$  and an element  $x_0 \in f^{-1}(y_0)$  such that  $\mathfrak{a}_0 = p_0^\alpha(\mathfrak{a}_\alpha) \rightarrow x_0$ ,  $x_0 \in U$  and  $f_\alpha(\mathfrak{a}_\alpha) = \mathfrak{b}_\alpha$ . Then we have  $U \in \mathfrak{a}_0$  and therefore  $(p_0^\alpha)^{-1}(U) \in \mathfrak{a}_\alpha$ . Hence  $f_\alpha((p_0^\alpha)^{-1}(U)) \cap \mathcal{B} \neq \emptyset$  which contradicts the definition of  $\mathcal{B}$ . Therefore, we have proved that  $y_0 \in Y - U^\sharp$ .

This means that (1) defines a map  $(-)^\sharp : \mathcal{O}X \rightarrow \mathcal{O}Y$ , that satisfies obviously (T1). So it remains to show that it also satisfies (T4). Let  $U \in \mathcal{O}X$ ,  $y \in U^\sharp$  and  $\Sigma \subseteq \mathcal{O}X$ , directed, be such that, for each  $S \in \Sigma$ ,  $y \notin S^\sharp$ . There exists an ordinal  $\alpha \in \lambda_Y$  such that, for each  $S \in \Sigma$ , the set  $\mathcal{B}_S = (p_0^\alpha)^{-1}(y) - f_\alpha((p_0^\alpha)^{-1}(S))$  is non-empty. Since  $\Sigma$  is filtered,  $\{\mathcal{B}_S \mid S \in \Sigma\}$  forms a filter base. Let  $\mathfrak{b}_\alpha$  be any ultrafilter containing  $\{\mathcal{B}_S \mid S \in \Sigma\} \cup (p_0^\alpha)^{-1}(\eta(y))$ . Since  $y \in U^\sharp$ , there exist an ultrafilter  $\mathfrak{a}_\alpha$  on  $X_\alpha$  and an element  $x \in U$  such that  $\mathfrak{a}_0 = (p_0^\alpha)_X(\mathfrak{a}_\alpha) \rightarrow x$ ,  $f(x) = y$  and  $f_\alpha(\mathfrak{a}_\alpha) = \mathfrak{b}_\alpha$ . Let  $V \in \mathcal{O}(x)$ . Then  $V \in \mathfrak{a}_0$  and therefore  $(p_0^\alpha)^{-1}(V) \in \mathfrak{a}_\alpha$ . Hence  $f_\alpha((p_0^\alpha)^{-1}(V)) \cap \mathcal{B}_S \neq \emptyset$  and therefore  $V \neq S$ , for each  $S \in \Sigma$ . Hence,  $\Sigma$  does not cover  $f^{-1}(y) \cap U$ , and (T4) follows.  $\square$

This proposition gives a general way of defining a map  $(-)^\sharp$  – in fact, the largest possible one –, as required in the definition of triquotient map, but that in general does not satisfy (T2):  $X^\sharp = Y$ . By the definition of  $(-)^\sharp$  it is clear what this condition means:  $f$  is  $\lambda_Y$ -surjective; that is:



Hence, we may now state the characterization of topological triquotient maps.

**Theorem 6.4.** *Let  $f : X \rightarrow Y$  be a continuous map between topological spaces. The following conditions are equivalent:*

- (i)  $f$  is a triquotient map;
- (ii)  $f$  is  $\Omega$ -surjective;
- (iii)  $f$  is  $\lambda_Y$ -surjective.

$\square$

In the finite case, since all ultrafilters are fixed,  $X_n$  may be described as the set of all  $(n+1)$ -chains  $x_n \rightarrow \cdots \rightarrow x_0$  of elements of  $X$ . The ultrarelatational structure is then described by

$$(x_n, \cdots, x_1, x_0) \rightarrow (x'_n, \cdots, x'_1, x'_0) : \iff (x_{n-1}, \cdots, x_0) = (x'_n, \cdots, x'_1).$$

From Theorem 6.4, we know that, if  $X$  and  $Y$  are finite, then  $f : X \rightarrow Y$  is a triquotient map if and only if  $f_n$  is surjective for every  $n \in \mathbb{N}$ , which is exactly Theorem I.

#### ACKNOWLEDGMENT

We thank George Janelidze, Manuela Sobral and Walter Tholen for valuable discussions on the subject of this paper.

#### REFERENCES

- [1] J. Adámek, H. Herrlich and G.E. Strecker, *Abstract and concrete categories* (Wiley Interscience, New York 1990). MR **91h**:18001
- [2] N. Bourbaki, *Topologie Générale* (Hermann, Paris 1961).
- [3] M. M. Clementino, On finite triquotient maps, *J. Pure Appl. Algebra* **168** (2002), 387-389.
- [4] B. J. Day and G. M. Kelly, On topological quotient maps preserved by pullbacks or products, *Proc. Cambridge Philos. Soc.* **67** (1970) 553-558. MR **40**:8024
- [5] O. Hájek, Notes on quotient maps, *Comment. Math. Univ. Carolinae* **7** (1966), 319-323. MR **34**:1992
- [6] G. M. Kelly, A unified treatment of transfinite constructions for free algebras, free monoids, colimits, associated sheaves, and so on, *Bull. Austral. Math. Soc.* **22** (1980) 1-83. MR **84a**:18011
- [7] G. Janelidze and M. Sobral, Finite preorders and topological descent, *J. Pure Appl. Algebra* (to appear).
- [8] G. Janelidze and W. Tholen, How algebraic is the change-of-base functor?, *Proceedings Conference Category Theory*, Como 1990, Lecture Notes in Mathematics **1488** (Springer, Berlin 1991) 174-186. MR **93f**:18011
- [9] E.G. Manes, A triple-theoretic construction of compact algebras, *Seminar on Triples and Categorical Homology Theory*, Lecture Notes in Mathematics **80** (Springer, Berlin 1969), pp. 91-118. MR **39**:5657
- [10] E. Michael, Bi-quotient maps and Cartesian products of quotient maps, *Ann. Inst. Fourier, Grenoble* **18** (1968) 287-302. MR **39**:6277
- [11] E. Michael, Complete spaces and tri-quotient maps, *Illinois J. of Math.* **21** (1977), 716-733. MR **57**:7543
- [12] T. Plewe, Localic triquotient maps are effective descent maps, *Math. Proc. Cambridge Philos. Soc.* **122** (1997), 17-43. MR **98b**:54022
- [13] J. Reiterman and W. Tholen, Effective descent maps of topological spaces, *Top. Appl.* **57** (1994), 53-69. MR **95f**:18008

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE DE COIMBRA, 3001-454 COIMBRA, PORTUGAL  
*E-mail address:* mmc@mat.uc.pt

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE DE COIMBRA, 3001-454 COIMBRA, PORTUGAL  
*E-mail address:* dirk@mat.uc.pt  
*Current address:* Departamento de Matemática, Universidade de Aveiro, 3810-193 Aveiro, Portugal  
*E-mail address:* dirk@mat.ua.pt