

P.I. ENVELOPES OF CLASSICAL SIMPLE LIE SUPERALGEBRAS

IAN M. MUSSON

(Communicated by Lance W. Small)

ABSTRACT. Let \mathfrak{g} be a classical simple Lie superalgebra. We describe the prime ideals P in the enveloping algebra $U(\mathfrak{g})$ such that $U(\mathfrak{g})/P$ satisfies a polynomial identity. If the factor algebra $U(\mathfrak{g})/P$ is not artinian, then it is an order in a matrix algebra over $K(z)$.

Throughout this paper we work over an algebraically closed field K of characteristic zero. All unadorned tensor products are taken over K . Let \mathfrak{g} be a finite dimensional classical simple Lie superalgebra over K . A factor algebra of the enveloping algebra $U(\mathfrak{g})$ satisfying a polynomial identity is called a *P.I. envelope* of \mathfrak{g} . Our aim is to describe all prime P.I. envelopes of \mathfrak{g} . If \mathfrak{g} has a nonartinian prime P.I. envelope it is not hard to show that the center of \mathfrak{g}_0 must be nonzero (Lemma 1.3). Thus by the classification theorem in [K1], $\mathfrak{g} = sl(m, n)$ with $m > n \geq 1$ or $\mathfrak{g} = osp(2, 2n)$.

It was shown by Bahturin and Montgomery that when $\mathfrak{g} = sl(m, n)$ with $m > n \geq 1$, \mathfrak{g} has a nonartinian P.I. envelope. In fact the proof of [BM, Theorem 4.2] shows that this is true also when $\mathfrak{g} = osp(2, 2n)$ although these algebras are omitted from the statement of [BM, Theorems 1.5 and 4.2]. If $\mathfrak{g} = sl(m, n)$ with $m > n \geq 1$ or $\mathfrak{g} = osp(2, 2n)$, then $\mathfrak{g}_0 = [\mathfrak{g}_0, \mathfrak{g}_0] \oplus Kz$ where z is central in \mathfrak{g}_0 . Furthermore as a \mathfrak{g}_0 -module via the adjoint action, $\mathfrak{g}_1 = \mathfrak{g}_1^+ \oplus \mathfrak{g}_1^-$, a direct sum of two simple submodules. If $\mathfrak{p} = \mathfrak{g}_0 \oplus \mathfrak{g}_1^+$, then $U(\mathfrak{g}_0)$ is a homomorphic image of $U(\mathfrak{p})$ and thus any $U(\mathfrak{g}_0)$ -module can be regarded as a $U(\mathfrak{p})$ -module. Choose a Cartan subalgebra \mathfrak{h} and a system of simple roots for \mathfrak{g}_0 . Let P^+ denote the corresponding set of dominant integral weights. If $\lambda \in P^+$ let L_λ be the finite dimensional simple $[\mathfrak{g}_0, \mathfrak{g}_0]$ -module with highest weight λ and set

$$V(\lambda) = U(\mathfrak{g}) \otimes_{U(\mathfrak{p})} (L_\lambda \otimes K[z]).$$

Our main result is as follows.

Main Theorem. *Set $P_\lambda = \text{ann}_{U(\mathfrak{g})} V(\lambda)$, $n = \dim_K L_\lambda$ and $N = n2^{\dim \mathfrak{g}_1^-}$. Then P_λ is a prime ideal of $U(\mathfrak{g})$ such that $U(\mathfrak{g})/P_\lambda$ is a subring of the matrix algebra $M_N(K[z])$ with Goldie quotient ring $M_N(K(z))$. In particular $U(\mathfrak{g})/P_\lambda$ is a prime P.I. algebra.*

Conversely, if P is a prime ideal in $U(\mathfrak{g})$ such that $U(\mathfrak{g})/P$ satisfies a polynomial identity, then $P = P_\lambda$ for a unique $\lambda \in P^+$.

Received by the editors June 12, 2001.

2000 *Mathematics Subject Classification.* Primary 17B20, 17B35.

If $\mathcal{C} = K[z] \setminus \{0\}$, then \mathcal{C} is an Ore set of regular elements in $U(\mathfrak{g})$. A key step in the proof that P_λ is prime is to show that the localized module $V(\lambda)_{\mathcal{C}}$ is in a natural way a $U(\mathfrak{g}) - F$ bimodule where $F = K(z)$, and then a simple Kac module over $U(\mathfrak{g}) \otimes F$. For the converse we use some results of E.S. Letzter concerning prime ideals in finite extensions of Noetherian rings.

1.1. Let R and S be prime Noetherian rings. An $R - S$ bimodule M is a *bond* from R to S if M is finitely generated and torsionfree both as a left R -module and as a right S -module.

Lemma. *Suppose M is a bond from R to S . Then:*

- (a) *R is artinian if and only if S is artinian.*
- (b) *R is a P.I. ring if and only if S is a P.I. ring.*

Proof. (a) This is [J, Theorem 5.2.9].

(b) This is Remark (2) after [BS, Prop. 2.5]. We give some details for the convenience of the reader. Let $D = \text{Fract } R$ and $E = \text{Fract } S$. By [BS, Prop. 2.5] there exists an integer t such that D embeds in the ring of $t \times t$ matrices $M_t(E)$ and E embeds in $M_t(D)$. Thus if S is P.I., then R embeds in $M_t(E)$ which is a central simple algebra by Posner's Theorem [McR, Theorem 13.6.5], so R is P.I. by the Amitsur-Levitzki Theorem [McR, Corollary 13.3.5]. Similarly if R is P.I. so is S .

1.2. Until the end of section 1.4 suppose that $R \subseteq S$ is an extension of Noetherian K -algebras of finite Gel'fand-Kirillov dimension. Assume that S is finitely generated and free as a right R -module. The following definitions are due to Letzter [L2], [L3].

i) Suppose P is a prime ideal of S and set $B = \text{Fract}(S/P)$. Let V_P be the set of prime ideals of R which are right annihilators of simple $B - R$ factor bimodules of B .

ii) Suppose Q is a prime ideal of R and set $A = \text{Fract}(R/Q)$. Let W_Q be the set of prime ideals of S which are left annihilators of simple $S - A$ factor bimodules of $S \otimes_R A$.

In addition set $J_Q = \ell - \text{ann}(S/SQ)$ and

$$X_Q = \{P \in \text{Spec } R \mid P \text{ is minimal over } J_Q\}.$$

These definitions are related by the following results.

Theorem. (a) *If $Q \in \text{Spec } R$ and $P \in \text{Spec } S$, then*

$$Q \in V_P \text{ if and only if } P \in W_Q.$$

Furthermore if this condition holds there is a bond from S/P to R/Q .

(b) $W_Q \subseteq X_Q$.

Proof. (a) follows from [L2, Lemma 3.2] and [L1, Lemma 1.1], while (b) follows from the proof of [L2, Proposition 4.2].

1.3. Lemma. *Let \mathfrak{g} be a classical simple Lie superalgebra such that there is a prime ideal P in $U(\mathfrak{g})$ with $U(\mathfrak{g})/P$ a nonartinian P.I. algebra. Then the center of \mathfrak{g}_0 is nonzero.*

Proof. We apply the results of the two previous subsections, with $R = U(\mathfrak{g}_0)$ and $S = U(\mathfrak{g})$. Choose $Q \in V_P$. Then there is a bond from S/P to R/Q by Theorem 1.2. Hence by Lemma 1.1, R/Q is a non-artinian P.I. algebra. The result follows from a result of Bahturin; see [BM, page 2837].

1.4. If the equivalent conditions of Theorem 1.2(a) hold we say that P lies directly over Q . Recall that a module over a prime Noetherian ring is *fully faithful* if every nonzero submodule is faithful. We require another result of Letzter [L3, Lemma 2.6 (iv)].

Lemma. *Suppose that P lies directly over Q and that M is a fully faithful S/P -module. Then there exists an R -submodule N of M such that $Q = \text{ann}_R N$ and N is a fully faithful R/Q -module.*

1.5. Again suppose that \mathfrak{g} is classical simple. We often write U for $U(\mathfrak{g})$. The simple artinian factor rings of U correspond to the finite dimensional simple \mathfrak{g} -modules and these have been classified [K1, Theorem 8].

For the remainder of this paper we assume therefore that P is a prime ideal of U , such that U/P is a nonartinian P.I. algebra. By Lemma 1.3 this means that \mathfrak{g}_0 has a nonzero center.

As noted in the introduction we have $\mathfrak{g}_0 = [\mathfrak{g}_0, \mathfrak{g}_0] \oplus Kz$, and we can choose z in such a way that $[z, x] = \pm x$ for all $x \in \mathfrak{g}^\pm$. Set $\mathfrak{h}' = \mathfrak{h} \oplus Kz$, so that \mathfrak{h}' is a Cartan subalgebra of \mathfrak{g}_0 and \mathfrak{g} . Fix a non-degenerate invariant bilinear form $(,)$ on $(\mathfrak{h}')^*$. For $\alpha \in (\mathfrak{h}')^*$, we write \mathfrak{g}^α for the corresponding root space. There is a unique $h_\alpha \in \mathfrak{h}'$ such that $(\mu, \alpha) = \mu(h_\alpha)$ for all $\mu \in (\mathfrak{h}')^*$. Let Δ_1^+ be the set of roots of \mathfrak{g}_1^+ . For $\alpha \in \Delta_1^+$, choose $e_{\pm\alpha}$ such that $\mathfrak{g}_1^{\pm\alpha} = Ke_{\pm\alpha}$ and $h_\alpha = [e_\alpha, e_{-\alpha}]$.

1.6. We construct a functor T between categories of left modules:

$$T : U(\mathfrak{g}_0)\text{-mod} \longrightarrow U(\mathfrak{g})\text{-mod}.$$

First set $\mathfrak{p} = \mathfrak{g}_0 \oplus \mathfrak{g}_1^+$ and $J = U(\mathfrak{p})\mathfrak{g}_1^+$. Then J is a nilpotent ideal of $U(\mathfrak{p})$ with $U(\mathfrak{p})/J \cong U(\mathfrak{g}_0)$. Thus we can regard $U(\mathfrak{g}_0)\text{-mod}$ as a subcategory of $U(\mathfrak{p})\text{-mod}$ and define $T(_) = U(\mathfrak{g}) \otimes_{U(\mathfrak{p})} (_)$.

We use the functor T to construct some examples of P.I. envelopes of \mathfrak{g} . If M is any $K[z]$ -module and $\lambda \in P^+$ we regard $L(\lambda, M) = L_\lambda \otimes M$ as a $U(\mathfrak{g}_0)$ -module by allowing $[\mathfrak{g}_0, \mathfrak{g}_0]$ (resp. Kz) to act on the first (resp. second) factor of the tensor product. For $a \in K$, let $\mathcal{O}_a = K[z]/(z - a)$ and set

$$L(\lambda) = L(\lambda, K[z]), \quad L(\lambda, a) = L(\lambda, \mathcal{O}_a),$$

$$V(\lambda) = T(L(\lambda)), \quad V(\lambda, a) = T(L(\lambda, a)).$$

The natural map $K[z] \longrightarrow \mathcal{O}_a$ induces an epimorphism of $U(\mathfrak{g})$ -modules

$$V(\lambda) \longrightarrow V(\lambda, a).$$

The pair $\lambda' = (\lambda, a)$ can be viewed as the element of $(\mathfrak{h}')^*$ with $\lambda'|_{\mathfrak{h}} = \lambda$ and $\lambda(z) = a$. The module $V(\lambda, a)$ is called the Kac module with highest weight λ' . By [K2, Proposition 2.9] $V(\lambda, a)$ is a simple $U(\mathfrak{g})$ -module if and only if $\lambda' = (\lambda, a)$ satisfies $(\lambda' + \rho, \alpha) \neq 0$ for all odd positive roots α . Here $\rho = \rho_0 - \rho_1$, where ρ_0 (resp. ρ_1) is the half-sum of the positive even (resp. odd) roots. Since $V(\lambda)$ maps onto any Kac module of the form $V(\lambda, a)$ we call $V(\lambda)$ the universal Kac-module with highest weight $\lambda \in \mathfrak{h}^*$.

1.7. The enveloping algebra U has a \mathbb{Z} -grading, $U = \bigoplus_{n \in \mathbb{Z}} U(n)$ extending the \mathbb{Z} -grading on \mathfrak{g} given by $\deg \mathfrak{g}_0 = 0$, $\deg \mathfrak{g}_1^\pm = \pm 1$. Henceforth the adjective “graded” refers to this grading. We use this grading to construct a useful localization of U .

Suppose that $M = \bigoplus M(n)$ is a graded U -module which is torsionfree as a $K[z]$ -module. We can make M into a $U - K[z]$ -bimodule via the rule

$$(1) \quad mf(z) = f(z - n)m$$

for $m \in M(n)$, $f(z) \in K[z]$. Let $F = K(z)$ and give $M^F = M \otimes_{K[z]} F$ the right F -module structure obtained by localization. In particular U^F becomes a $U - F$ -bimodule in this way and we can extend the algebra structure on U to U^F by

$$(u \otimes f_1(z))(v \otimes f_2(z)) = uv \otimes f_1(z + n)f_2(z)$$

for $u \in U$, $v \in U(n)$ and $f_1, f_2 \in F$. It is now easy to verify the following:

Lemma. *The multiplicative set $\mathcal{C} = K[z] \setminus \{0\}$ is Ore in U and $U_{\mathcal{C}} \cong U^F$ with the above algebra structure. If M is a graded left U -module which is torsionfree as a $K[z]$ -module, then $M_{\mathcal{C}} \cong M \otimes_{K[z]} F$ as a $U - F$ bimodule via the map*

$$f^{-1}(z)m \longrightarrow m \otimes f^{-1}(z + n)$$

for $m \in M(n)$, $f(z) \in K[z]$. In addition if N is any graded U -submodule of $M_{\mathcal{C}}$, then N is a $U_{\mathcal{C}}$ -submodule if and only if it is a $U - F$ sub-bimodule of $M_{\mathcal{C}}$.

1.8. If V is a vector space over K we write V_F for $V \otimes F$. If A is a K -algebra and M is a left A -module, then A_F is an F -algebra and M_F an A_F -module by extension of scalars.

We apply these remarks to the universal Kac module $V(\lambda) = U(\mathfrak{g}) \otimes_{U(\mathfrak{p})} L(\lambda)$. If $\Lambda = \bigoplus \Lambda^n$ is the exterior algebra on \mathfrak{g}_1^- , then as a left $U(\mathfrak{g}_0)$ -module

$$V(\lambda) \cong \bigoplus \Lambda^n \otimes L(\lambda).$$

By definition $L(\lambda) = L_\lambda \otimes K[z]$, and so $L(\lambda)$ and $V(\lambda)$ are in an obvious way right $K[z]$ -modules. Since $\deg \mathfrak{g}_1^- = -1$ the gradings on Λ and U satisfy $\Lambda^n \subseteq U(-n)$. Observe that the $K[z]$ -bimodule structure on $V(\lambda)$ satisfies (1) in section 1.7.

Note also that $V(\lambda)$ is torsionfree as a left (and right) $K[z]$ -module. Thus $V(\lambda)_{\mathcal{C}}$ is a $U - F$ bimodule or equivalently a left U_F -module and we have

$$\begin{aligned} V(\lambda)_{\mathcal{C}} &\cong (U(\mathfrak{g}) \otimes_{U(\mathfrak{p})} L(\lambda)) \otimes_{K[z]} F \\ &\cong U(\mathfrak{g}) \otimes_{U(\mathfrak{p})} (L(\lambda) \otimes_{K[z]} F) \\ &\cong U(\mathfrak{g})_F \otimes_{U(\mathfrak{p})_F} (L(\lambda) \otimes_{K[z]} F). \end{aligned}$$

Next we consider the $U(\mathfrak{p})_F$ -module $L(\lambda) \otimes F$. This is annihilated by \mathfrak{g}_{1F}^+ and so is a $U(\mathfrak{g}_0)_F$ -module. In fact it is the finite dimensional simple module over this algebra whose highest weight λ' is the unique F -linear map $\mathfrak{h}'_F \rightarrow F$ such that $\lambda'|_{\mathfrak{h}} = \lambda$ and $\lambda'(z) = z$. Thus $V(\lambda)_{\mathcal{C}}$ is a Kac module over $U(\mathfrak{g}) \otimes F$. In fact we have

Proposition. (a) *The module $V(\lambda)_{\mathcal{C}}$ is a simple Kac module with highest weight λ' module over the algebra $U(\mathfrak{g}) \otimes F$.*

(b) *The module $V(\lambda)_{\mathcal{C}}$ is a simple module over the algebra $U(\mathfrak{g})_{\mathcal{C}}$.*

Proof. It remains to show simplicity in both cases.

(a) We extend $(,)$ to a bilinear form $(,)_F$ on $(\mathfrak{h}')^*_F$. It suffices to show that $(\lambda' + \rho, \alpha) \neq 0$ for all odd positive roots α . Since the highest exterior power of \mathfrak{g}_1^+

is trivial as a \mathfrak{g}_0 -module we have $(\rho_1, \beta) = 0$ for all even roots β . Thus $\mathfrak{h}^\perp = K\rho_1$. If α is an odd root, it follows that $(\rho_1, \alpha) = \rho_1(h_\alpha) \neq 0$. Hence $h_\alpha \notin (K\rho_1)^\perp = \mathfrak{h}$, so $h_\alpha - bz \in \mathfrak{h}$ for some nonzero $b \in K$. Therefore

$$(\lambda' + \rho, \alpha)_F = \lambda'(h_\alpha - bz + bz)_F + (\rho, \alpha) = bz + \lambda(h_\alpha - bz) + (\rho, \alpha),$$

which is a linear polynomial in z .

(b) Fix an order on Δ_1^+ and for $I \subseteq \Delta_1^+$ set

$$e_I = \prod_{\alpha \in I} e_{-\alpha}$$

where the product is taken with respect to this order. Let N be a nonzero $U(\mathfrak{g})_{\mathcal{C}}$ -submodule of $V(\lambda)_{\mathcal{C}}$ and suppose

$$n = \sum_I e_I n_I$$

is nonzero with $n_I \in L(\lambda) \otimes_{K[z]} F$ for all I . Choose m minimal such that $n_I \neq 0$ for some subset I with $|I| = m$. Set $I' = \Delta_1^+ \setminus I$. Then

$$e_{I'} n = \pm e_{\Delta_1^+} n_I$$

is a nonzero homogeneous element of N , so generates a graded submodule. It follows from Lemma 1.7 that $N = V(\lambda)_{\mathcal{C}}$.

Corollary. (a) *If N is any nonzero $U(\mathfrak{g})$ -submodule of $V(\lambda)$, then N contains a submodule isomorphic to $V(\lambda)$.*

(b) $P_\lambda = \text{ann}_{U(\mathfrak{g})} V(\lambda)$ is a prime ideal of $U(\mathfrak{g})$.

Proof. (a) By the Proposition $N_{\mathcal{C}} = (V(\lambda))_{\mathcal{C}}$. Hence $(L_\lambda \otimes 1) \subseteq N_{\mathcal{C}}$ so $L_\lambda \otimes (f) \subseteq N$ for some nonzero f . The submodule of $V(\lambda)$ generated by $L_\lambda \otimes (f)$ is isomorphic to $T(L(\lambda, (f))) \cong V(\lambda)$.

(b) This follows since any nonzero submodule of $V(\lambda)$ has annihilator P_λ , by part (a).

1.9. Lemma. *Identify $L(\lambda)$ with the $U(\mathfrak{p})$ -submodule $1 \otimes L(\lambda)$ of $V(\lambda)$ and let $J = U(\mathfrak{p})\mathfrak{g}_1^+$. Then $\text{ann}_{V(\lambda)} J = L(\lambda)$.*

Proof. We use the same notation as in the proof of Proposition 1.8(b).

For $0 \leq m \leq |\Delta_1^+|$ set

$$V(m) = \bigoplus_{|I|=m} e_I L(\lambda).$$

Then $V(\lambda) = \bigoplus_m V(m)$ and $\mathfrak{g}_1^+ V(m) \subseteq V(m-1)$. Therefore it suffices to show that if $m > 0$, then $\mathfrak{g}_1^+ (\sum_{|I|=m} e_I w_I) \neq 0$ provided the $w_I \in L(\lambda)$ are not all zero. Now $L(\lambda) = L_\lambda \otimes K[z]$ has a filtration given by setting $\text{deg}(L_\lambda \otimes z^n) = n$. Choose I so that w_I has maximum degree in this filtration, choose $\alpha \in I$ and set $H = I \setminus \{\alpha\}$. Note that

$$\text{deg}(h_\alpha w_I) = \text{deg}(w_I) + 1.$$

Using the formula

$$[e_\alpha, ab] = [e_\alpha, a]b \pm a[e_\alpha, b]$$

for homogeneous $a, b \in U(\mathfrak{g})$ we see that

$$e_\alpha e_I w_I = \pm e_H h_\alpha w_I$$

plus a sum of terms of smaller degree. The result follows easily from this.

1.10. The next result is an easy consequence of the Artin-Wedderburn theorem.

Lemma. *Let U be a K -algebra and L a finite dimensional simple U -module. Then for any field extension K' of K we have*

$$\text{End}_{U \otimes_K K'}(L \otimes_K K') = K'.$$

1.11. Lemma. *For any $\lambda \in P^+$,*

$$\text{End}_{U(\mathfrak{g})}(V(\lambda)) \cong K[z].$$

Proof. By the adjoint isomorphism $f \in \text{End}_{U(\mathfrak{g})}(V(\lambda))$ is determined by

$$f_1 = f|_{L(\lambda)} \in \text{Hom}_{U(\mathfrak{p})}(L(\lambda), V(\lambda)).$$

If J is as in Lemma 1.9, then $f_1(L(\lambda)) \subseteq \text{ann}_{V(\lambda)} J = L(\lambda)$, and hence

$$f_1 \in \text{End}_{U(\mathfrak{p})}(L_\lambda \otimes K[z]) = \text{End}_{U(\mathfrak{g}_0)}(L_\lambda \otimes K[z]) \cong K[z],$$

using Lemma 1.10.

1.12. Suppose that \mathfrak{k} is a semisimple Lie algebra over K and C is a commutative K -algebra. We describe the prime ideals Q of $R = U(\mathfrak{k}) \otimes C$ such that R/Q is P.I. Since C is central, $q = Q \cap C$ is prime in C and by replacing R by the factor ring R/Rq we can assume that $Q \cap C = 0$. There is a one-one correspondence between prime ideals Q of R such that $Q \cap C = 0$ and prime ideals of $U(\mathfrak{k}) \otimes \text{Fract}(C)$. Thus we may assume that C is a field extension of K . By the argument on page 2837 of [BM] Q is the annihilator of a finite dimensional simple module over $U(\mathfrak{k}) \otimes C$.

To apply this to our reductive algebra \mathfrak{g}_0 with center Kz set $\mathfrak{k} = [\mathfrak{g}_0, \mathfrak{g}_0]$, $R = U(\mathfrak{g}_0)$ and $C = K[z]$. If $q \neq 0$, then $R/Rq \cong U(\mathfrak{k})$ and R/Q is artinian. Thus if R/Q is nonartinian, then $q = 0$ and Q corresponds to the annihilator of a finite dimensional simple module over $U(\mathfrak{k}) \otimes \text{Fract}(C)$. This gives the following result.

Lemma. *Suppose Q is a prime ideal of $U(\mathfrak{g}_0)$ such that $U(\mathfrak{g}_0)/Q$ is a nonartinian P.I. ring. Then for some uniquely determined $\lambda \in P^+$, $Q = \text{ann}_{U(\mathfrak{g}_0)} L(\lambda)$.*

Proof of the Main Theorem. Suppose $\lambda \in P^+$ and let $n = \dim_K L_\lambda$. By Corollary 1.8 P_λ is a prime ideal in $U(\mathfrak{g})$. Set $U_\lambda = U(\mathfrak{g})/P_\lambda$. Note that $V(\lambda)$ is a torsionfree $K[z]$ -module and thus U_λ embeds in $(U_\lambda)_C$. Since $V(\lambda)$ is a $U(\mathfrak{g}) - K[z]$ -bimodule which is free of rank $N = n2^{\dim \mathfrak{g}_1}$ on the right, U_λ embeds in $M_N(K[z])$. This embedding induces an embedding of $(U_\lambda)_C$ into $M_N(F)$ which is surjective since $V(\lambda)_C$ is a simple $(U_\lambda)_C$ -module of dimension N over its endomorphism ring F .

Conversely suppose P is a prime ideal of $U(\mathfrak{g})$ with $U(\mathfrak{g})/P$ a nonartinian P.I. ring. We apply the results in sections 1.1 and 1.2 with $R = U(\mathfrak{p})$ and $S = U(\mathfrak{g})$. If $Q \in V_P$ there is a bond from S/P to R/Q , so R/Q is a nonartinian P.I. ring, by Lemma 1.1. Lemma 1.12 implies that $Q = Q_\lambda$ for some $\lambda \in P^+$. Since $P \in X_Q$, P is minimal over $\text{ann}_S(S/SQ)$ which equals $\text{ann}_S V(\lambda) = P_\lambda$ by [BGR, Satz 10.4]. As P_λ is prime we get $P = P_\lambda$. To show that λ is uniquely determined by P it suffices to show that $V_{P_\lambda} = \{Q_\lambda\}$. However if $Q' \in V_{P_\lambda}$, then by Lemma 1.4 $Q' = \text{ann}_R N$ for some R -submodule N of $V(\lambda)$ which is fully faithful as an R/Q' -submodule. Since $J = U(\mathfrak{p})\mathfrak{g}_1^+$ is nilpotent $J \subseteq Q'$, so using Lemma 1.9 $N \subseteq \text{ann}_{V(\lambda)} J = L(\lambda)$. However every nonzero submodule of $L(\lambda)$ has annihilator Q_λ , so $Q' = Q_\lambda$ as desired.

REFERENCES

- [BM] Y. Bahturin and S. Montgomery, PI-envelopes of Lie superalgebras, Proc. Amer. Math. Soc., 127 (1999), 2829-2839. MR **2000a**:17029
- [BGR] W. Borho, P. Gabriel and R. Rentschler, Primideale in Einhüllenden auflösbarer Lie-Algebren, Lecture notes in Mathematics, 357, Springer-Verlag, Berlin, 1973. MR **51**:12965
- [BS] K.A. Brown and S.P. Smith, Bimodules over a solvable algebraic Lie algebra, Quart. J. Math. 36 (1985), 129-139. MR **87d**:17009
- [J] A.V. Jategaonkar, Localization in Noetherian Rings, LMS Lecture Notes 98, Cambridge University Press, Cambridge 1986. MR **88c**:16005
- [K1] V.G. Kac, Lie Superalgebras, Adv. in Math., 26 (1977), 8-96. MR **58**:5803
- [K2] V.G. Kac, Representations of classical Lie superalgebras, pages 597-626. Differential Geometric Methods in Mathematical Physics II, ed. K. Bleuler et al. Lecture Notes in Mathematics, 676, Springer-Verlag, Berlin, 1978. MR **80f**:17006
- [L1] E.S. Letzter, Primitive ideals in finite extensions of Noetherian rings, J. London Math. Soc. 39 (1989), 427-435. MR **90f**:16013
- [L2] E.S. Letzter, Finite correspondence of spectra in Noetherian ring extensions, Proc. Amer. Math. Soc. 116 (1992), 645-652. MR **93a**:16003
- [L3] E.S. Letzter, A bijection of spectra for classical Lie superalgebras of Type I, J. London Math. Soc. 53 (1996), 39-49. MR **96k**:17016
- [McR] J.C. McConnell and J.C. Robson, Noncommutative Noetherian rings, Wiley-Interscience, Chichester 1987. MR **89j**:16023; revised edition MR **2001i**:16039

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF WISCONSIN, MILWAUKEE, WISCONSIN 53211

E-mail address: musson@csd.uwm.edu