

INFINITE DETERMINACY ON A CLOSED SET FOR SMOOTH GERMS WITH NON-ISOLATED SINGULARITIES

VINCENT THILLIEZ

(Communicated by Mei-Chi Shaw)

ABSTRACT. We give necessary and sufficient conditions for infinite determinacy of a smooth function germ whose critical locus contains a given set. This set is assumed to be the zero variety X of some analytic map-germ having maximal rank on a dense subset of X . We obtain a result in terms of Łojasiewicz estimates which extends, in particular, previous works by Sun and Wilson on line singularities, and by Grandjean on singularities of codimension 1 or 2.

INTRODUCTION

Infinite determinacy is a way to express the stability of smooth function-germs under flat perturbations. Let \mathcal{E}_n denote the ring of C^∞ function-germs at the origin in \mathbb{R}^n and let \underline{m} denote its maximal ideal. Consider the ideal of flat germs $\underline{m}^\infty = \bigcap_{k \geq 0} \underline{m}^k$. An element f of \mathcal{E}_n is said to be *infinitely determined* if, for any element u of \underline{m}^∞ , there exists a germ Φ of a C^∞ -diffeomorphism at the origin such that $f + u = f \circ \Phi$. In abbreviated form, this can be written

$$(1) \quad f + \underline{m}^\infty \subset f \circ \mathcal{R},$$

where \mathcal{R} denotes the group of germs at the origin of C^∞ -diffeomorphisms of \mathbb{R}^n . In what follows, for any subset \mathcal{R}' of \mathcal{R} , we shall always use the notation $f \circ \mathcal{R}'$ to denote the set $\{f \circ \Phi; \Phi \in \mathcal{R}'\}$.

It is known (see [7], [12] or part II of [11]) that (1) holds if and only if

$$(2) \quad \underline{m}^\infty \subset J_f,$$

where J_f denotes the Jacobian ideal $(\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n})\mathcal{E}_n$. Note that condition (2) implies that ∇f has at most an isolated singularity at 0.

The case of non-isolated singularities is much less understood. Typically, one considers germs with a prescribed critical locus X , or germs belonging to a given ideal I . If X is a line, necessary and sufficient conditions for the corresponding version of infinite determinacy have been stated first by Sun and Wilson [9]. The main theorem of [9] appears as a nice extension of the pioneering work of Siersma [8] on finite determinacy for line singularities. Unfortunately, the proof of the crucial lemma 4.4 in the paper of Sun and Wilson is not correct, and there is currently no known correction [13]. Among other applications, here we shall provide a complete proof of a slightly modified version of their main statement. To be more precise, a

Received by the editors December 13, 2004.

2000 *Mathematics Subject Classification*. Primary 58K40.

©2005 American Mathematical Society
 Reverts to public domain 28 years from publication

part of the results in [9] can be described as follows. The set X is the x_n -axis, and I is the ideal generated in \mathcal{E}_n by the $n-1$ first coordinate functions x_1, \dots, x_{n-1} . It is easy to see that the critical locus of an element f of \mathcal{E}_n contains X (as a set germ) if and only if f belongs to I^2 , and that the set of flat elements of I^2 is precisely the ideal $\underline{m}^\infty I^2$. Accordingly, an element f of I^2 is said to be *infinitely determined relative to I^2* if

$$(3) \quad f + \underline{m}^\infty I^2 \subset f \circ \mathcal{R}^X,$$

where \mathcal{R}^X denotes the subgroup of \mathcal{R} given by the elements which preserve X (or, equivalently in this situation, which preserve I). It is stated in [9] that (3) holds if and only if

$$(4) \quad \underline{m}^\infty I \subset J_f.$$

The “only if” part relies on another characterization of (3) involving Lojasiewicz inequalities for ∇f and for a suitable partial Hessian of f , in the spirit of the classical formulation of (2) in terms of Lojasiewicz estimates for ∇f . This is the so-called *real isolated line singularity* condition in [9]. In order to show that (3) implies this condition, some information on the non-degeneracy of partial Hessians along the critical locus is required, and this is where a gap occurs¹ in [9]. Our main result implies, as a particular case, that the real isolated line singularity condition characterizes a determinacy property which, instead of \mathcal{R}^X , involves the (smaller) subgroup $\mathcal{R}_{\text{fix}}^X$ of diffeomorphisms preserving X *pointwise*.

Although quite attractive, the real isolated line singularity condition is not easy to extend to wider classes of sets X . A first step towards more general situations has been made by Grandjean [3], who studied the case of ideals I generated by real-analytic germs ψ_1, \dots, ψ_p defining a complete intersection variety X with at most an isolated singular point at the origin. It is shown in [3] that the implication (3) \implies (4) essentially still holds in this setting, under the additional assumption $\text{codim } X \leq 2$. Note that the proof of Grandjean is not affected by the defective lemma in [9], since working in codimension 1 or 2 makes it possible to bypass the argument. On the other hand, [3] adds an *a priori* assumption on the partial Hessians, namely non-degeneracy on $X \setminus \{0\}$. As another particular case of our main result, this extra requirement can be suppressed, provided, once again, \mathcal{R}^X is replaced by $\mathcal{R}_{\text{fix}}^X$.

Beyond these adjustments, the main purpose of the present paper is to extend the aforementioned results by removing the restrictive assumption on the codimension of X and, to some extent, on its singular part. In order to achieve this latter point, we involve flatness on a given closed set Y , maybe larger than the single point 0. This viewpoint has been recently used in [5] to extend the equivalence (1) \iff (2): indeed, if \underline{m}_Y^∞ denotes the ideal of germs in \mathcal{E}_n which are flat on Y , theorem 36 of [5] asserts that an element f of \mathcal{E}_n satisfies $f + \underline{m}_Y^\infty \subset f \circ \mathcal{R}$ if and only if $\underline{m}_Y^\infty \subset J_f$. In order to generalize (3) \iff (4) in the same way, here we shall investigate the relationship between the conditions $f + \underline{m}_Y^\infty I^2 \subset f \circ \mathcal{R}'$ (for some suitable subgroup \mathcal{R}' of \mathcal{R}) and $\underline{m}_Y^\infty I \subset J_f$. This will lead us to the statement of Theorem 2.1. The geometric requirements are described precisely in Subsection 1.2 below: roughly speaking, the regular part of X has to be dense in X , and its singular part has

¹The computation of $\det A'_m$ in Lemma 4.4 of [9] is erroneous because the addition of an extra term $\frac{\det A_m}{\det B} y_m y_l$ to the germ f modifies the Hessian symmetrically, and not only the coefficient located at the m -th row and the l -th column.

to be contained in Y . There is no restriction on $\text{codim } X$. Of course, the key fact is that we are able to obtain a characterization of infinite determinacy in terms of Lojasiewicz estimates, quite in the same spirit as the real isolated line singularity condition of Sun and Wilson.

1. DEFINITIONS AND TECHNICAL FRAMEWORK

1.1. Notation. For any multi-index $J = (j_1, \dots, j_n)$ in \mathbb{N}^n , of length $|J| = j_1 + \dots + j_n$, we put $D^J = \partial^j / \partial x_1^{j_1} \dots \partial x_n^{j_n}$. For any finite family $\lambda_1, \dots, \lambda_p$ of real numbers, we denote by $\text{Diag}(\lambda_1, \dots, \lambda_p)$ the $p \times p$ diagonal matrix whose j -th diagonal coefficient is λ_j for $j = 1, \dots, p$. The differential of a map-germ $G : (\mathbb{R}^n, 0) \longrightarrow (\mathbb{R}^m, 0)$ will be denoted by dG and its jacobian matrix by G' .

Definition 1.1.1. Let Z be a germ of subset of \mathbb{R}^n at the origin. We denote by \mathcal{R}^Z (resp. $\mathcal{R}_{\text{fix}}^Z$) the set of elements Φ of \mathcal{R} which preserve Z , that is, $\Phi(Z) \subset Z$ (resp. which preserve Z pointwise, that is, $\Phi(x) = x$ for any $x \in Z$).

We recall from the Introduction that \mathcal{R} denotes the group of germs of C^∞ -diffeomorphisms of $(\mathbb{R}^n, 0)$. The set $\mathcal{R}_{\text{fix}}^Z$ is obviously a subgroup of \mathcal{R} , whereas \mathcal{R}^Z may not be a subgroup.

Definition 1.1.2. Let V and W be two germs of subsets at the origin in \mathbb{R}^n . We say that a function $g : V \longrightarrow \mathbb{R}$ satisfies the Lojasiewicz inequality $\mathfrak{L}(V, W)$ if there exist constants $C > 0$ and $\alpha \geq 0$ such that $|g(x)| \geq C \text{dist}(x, W)^\alpha$ for any $x \in V$.

Remark 1.1.3. Throughout the article, properties holding on a given subset V of \mathbb{R}^n are always understood in the sense of germs, that is, on a sufficiently small representative of V .

1.2. Non-isolated singularities. Let $\psi = (\psi_1, \dots, \psi_p)$ be a real-analytic map-germ $(\mathbb{R}^n, 0) \longrightarrow (\mathbb{R}^p, 0)$ with $p \leq n$. With ψ we associate the zero variety $X = \psi^{-1}(\{0\})$ and the ideal $I = (\psi_1, \dots, \psi_p)\mathcal{E}_n$. We define

$$\Sigma = \{x \in X ; d\psi_1(x) \wedge \dots \wedge d\psi_p(x) = 0\}$$

and we make the following assumption:

$$(5) \quad X \setminus \Sigma \text{ is dense in } X.$$

Any sufficiently general version of the Lojasiewicz inequality [1] shows that

$$(6) \quad |d\psi_1 \wedge \dots \wedge d\psi_p| \text{ satisfies } \mathfrak{L}(X, \Sigma).$$

The analyticity of ψ also ensures that I and its powers are closed [6]. Using section V.4 of [10], we therefore have

$$(7) \quad \underline{m}_X^\infty \subset I^2.$$

For any germ of a closed subset Y at the origin of \mathbb{R}^n , we also have, by proposition V.2.3 of [10],

$$(8) \quad \underline{m}_Y^\infty \cap I^2 = \underline{m}_Y^\infty I^2.$$

Let J_X denote the set of germs f satisfying $f(0) = 0$ and whose critical locus contains X . Using local coordinates on the smooth submanifold $X \setminus \Sigma$, it is easy to show that any element f of J_X vanishes on $X \setminus \Sigma$, hence on X by (5). Thus, if I_X denotes the ideal of elements of \mathcal{E}_n vanishing on X , it follows that J_X is the so-called *primitive ideal* of I_X . Obviously, J_X contains I^2 . It coincides with I^2

in certain situations, as in [9]. The following lemma provides a simple sufficient condition for this equality.

Lemma 1.2.1. *One has $J_X = I^2$ whenever $I_X = I$.*

Proof. Assume $I_X = I$, and let f be an element of J_X . Then we have $f = \sum_{j=1}^p f_j \psi_j$ with $\sum_{j=1}^p f_j(x) d\psi_j(x) = 0$ for any $x \in X$. Since the differentials $d\psi_j(x)$ are independent at any point x of $X \setminus \Sigma$, we derive that each f_j vanishes on $X \setminus \Sigma$, hence on X by (5). Thus each f_j belongs to I , and f belongs to I^2 . \square

In what follows, just as in [3], [4], [8] and [9], the study will be limited to elements f of I^2 . Beside technical considerations, the preceding lemma provides a motivation for this approach. Note that the difference of two elements of I^2 is flat on Y if and only if it belongs to $\underline{m}_Y^\infty I^2$, by virtue of (8). Thus, we are led to consider a notion of determinacy involving $f + \underline{m}_Y^\infty I^2$, as can be seen in the statement of Theorem 2.1 below.

1.3. Transversal Hessians. The Hessian matrix of an element f of \mathcal{E}_n will be denoted by f'' . Assuming that f belongs to I^2 , we can write $f = \sum_{1 \leq i, j \leq p} f_{ij} \psi_i \psi_j$ for some suitable elements f_{ij} of \mathcal{E}_n satisfying $f_{ij} = f_{ji}$. Of course, the f_{ij} are not unique. However, uniqueness holds in restriction to X , as shown below.

Lemma 1.3.1. *The matrix $H_f = (f_{ij})_{1 \leq i, j \leq p}$ satisfies*

$$(9) \quad {}^t(\psi'(x))H_f(x)\psi'(x) = \frac{1}{2}f''(x) \text{ for any } x \in X.$$

In particular, for $x \in X$, the matrix $H_f(x)$ is fully determined by ψ and f .

Proof. The identity (9) follows from a direct computation. The uniqueness statement is an obvious consequence of (9) since ψ has maximal rank on a dense subset of X , namely on $X \setminus \Sigma$. \square

From now on, H_f will be considered in restriction to X . At any point of the smooth submanifold $X \setminus \Sigma$, it can be viewed as a Hessian of f with respect to transversal directions.

Definition 1.3.2. For $x \in X$, we put $D_f(x) = \det H_f(x)$.

As shown by Lemma 1.3.1, the function $D_f : X \rightarrow \mathbb{R}$ depends only on ψ and f . The following elementary property will play an important role in Section 3.

Lemma 1.3.3. *Let Z be a subset of $X \setminus \Sigma$, let Φ be an element of \mathcal{R}^Z such that both f and $f \circ \Phi$ belong to I^2 , and let x be any point of Z . Then one has $D_{f \circ \Phi}(x) = 0$ if and only if $D_f(\Phi(x)) = 0$.*

Proof. Note first that $D_f(\Phi(x))$ makes sense since $\Phi(Z) \subset Z \subset X \setminus \Sigma$; in particular, both x and $\Phi(x)$ belong to X . A direct computation then yields $(f \circ \Phi)''(x) = {}^t(\Phi'(x))f''(\Phi(x))\Phi'(x)$. Using Lemma 1.3.1, we get ${}^t(\psi'(x))H_{f \circ \Phi}(x)\psi'(x) = {}^t((\psi \circ \Phi)'(x))H_f(\Phi(x))(\psi \circ \Phi)'(x)$. The result follows, since both $\psi'(x)$ and $(\psi \circ \Phi)'(x)$ have maximal rank. \square

1.4. Jacobian and Fitting ideals. In the setting of Subsection 1.2, consider the map $\sigma : \mathcal{E}_n^p \longrightarrow \mathcal{E}_n$ given by $\sigma(f_1, \dots, f_p) = \sum_{i=1}^p f_i \psi_i$. Since \mathcal{E}_n is flat over the ring \mathcal{O}_n of analytic function-germs, the module of smooth relations between ψ_1, \dots, ψ_p , that is, $\ker \sigma$, admits a finite system of generators k^1, \dots, k^q belonging to \mathcal{O}_n^p . We can assume that this system includes all the trivial relations $\psi_s e_r - \psi_r e_s$, where (e_1, \dots, e_p) is the canonical basis of \mathcal{E}_n^p and r, s are integers with $1 \leq r < s \leq p$.

Now let f be an element of I^2 and put $M_f = \sigma^{-1}(J_f)$, where J_f denotes the Jacobian ideal of f , as in the Introduction. For any $j = 1, \dots, n$, it is easy to check that the element $h^j = (h_1^j, \dots, h_p^j)$ of \mathcal{E}_n^p defined by

$$(10) \quad h_i^j = \sum_{k=1}^p \left(2f_{ik} \frac{\partial \psi_k}{\partial x_j} + \frac{\partial f_{ik}}{\partial x_j} \psi_k \right) \quad \text{for } i = 1, \dots, p$$

satisfies $\sigma(h^j) = \partial f / \partial x_j$. Then, obviously, M_f is the submodule of \mathcal{E}_n^p generated by $h^1, \dots, h^n, k^1, \dots, k^q$, so that we can write $M_f = \lambda(\mathcal{E}_n^{n+q})$, where $\lambda : \mathcal{E}_n^{n+q} \longrightarrow \mathcal{E}_n^p$ is the morphism of free modules defined by

$$\lambda(\xi_1, \dots, \xi_{n+q}) = \sum_{j=1}^n \xi_j h^j + \sum_{j=1}^q \xi_{n+j} k^j.$$

Denote by Λ the matrix of λ in the canonical bases of \mathcal{E}_n^{n+q} and \mathcal{E}_n^p , and consider, as in [3] or [9], the ideal K_f generated in \mathcal{E}_n by the minors of order p of Λ . Following section 20.2 of [2], we see that K_f is precisely the Fitting ideal $\text{Fitt}_0(\mathcal{E}_n^p / M_f)$ as can be seen from the free presentation

$$\mathcal{E}_n^{n+q} \xrightarrow{\lambda} \mathcal{E}_n^p \longrightarrow \mathcal{E}_n^p / M_f \longrightarrow 0.$$

In particular, K_f depends only on ψ and f , and it annihilates \mathcal{E}_n^p / M_f , hence the inclusion $K_f \mathcal{E}_n^p \subset M_f$. Taking the image by σ , we therefore get

$$(11) \quad K_f I \subset J_f.$$

We now have all the tools required for the statement of our result.

2. A THEOREM OF INFINITE DETERMINACY

Theorem 2.1. *Let ψ , X and I be defined as in Subsection 1.2, and let Y be a germ of closed subset of \mathbb{R}^n at the origin satisfying*

$$(12) \quad \Sigma \subset Y.$$

Then for any element f of I^2 , the following conditions are equivalent:

$$(13) \quad f + \underline{m}_Y^\infty I^2 \subset f \circ \mathcal{R}_{\text{fix}}^{X \setminus Y},$$

$$(14) \quad \underline{m}_Y^\infty I \subset J_f,$$

$$(15) \quad \underline{m}_Y^\infty \subset K_f,$$

$$(16) \quad |\nabla f| \text{ satisfies } \mathfrak{L}(\mathbb{R}^n, X \cup Y), \text{ and } D_f \text{ satisfies } \mathfrak{L}(X, Y).$$

Moreover, if one assumes additionally

$$(17) \quad D_f^{-1}(\{0\}) \subset Y,$$

then (13) is equivalent to the (a priori weaker) condition

$$(18) \quad f + \underline{m}_Y^\infty I^2 \subset f \circ \mathcal{R}^{X \setminus Y}.$$

All of Section 3 will be devoted to proving the theorem. We shall first complete Section 2 with some comments and examples.

2.2. Comments.

Remark 2.2.1. The particular case $Y = \{0\}$ corresponds to the natural notion of determinacy by the Taylor jet at a single point. In view of (12), our result covers this case, provided X has at most an isolated singular point. Note that we then obviously have $\mathcal{R}^{X \setminus Y} = \mathcal{R}^X$.

Remark 2.2.2. In this special situation, it would not be difficult to add to the statement of Theorem 2.1 another equivalent condition involving suitable substitutes for the tangent spaces to group orbits which are customary in such problems. We shall not describe this condition nor use it here (see, however, Example 2.3.2 below).

2.3. Examples.

Example 2.3.1 (isolated critical points). From Theorem 2.1, we can easily recover the equivalence (1) \iff (2) of the Introduction, and more generally theorem 36 of [5]. Indeed, take $p = n$ and $\psi_j(x) = x_j$ so that $I = \underline{m}$, $X = \{0\}$ and $\Sigma = \emptyset$. Then (12) is trivial, and it is easy to check that $\underline{m}_Y^\infty I^2 = \underline{m}_Y^\infty I = \underline{m}_Y^\infty$. The result follows.

Example 2.3.2 (isolated critical lines). The situation studied in [9] corresponds to $p = n - 1$, $\psi_j(x) = x_j$, so that X is the x_n -axis. For $Y = \{0\}$, condition (16) coincides with the real isolated line singularity condition of Sun and Wilson. Taking Remark 2.2.1 into account, we recover, as announced in the Introduction, the main result of [9] with \mathcal{R}^X replaced by $\mathcal{R}_{\text{fix}}^X$. Explicitly, an element Φ of \mathcal{R} belongs to \mathcal{R}^X if and only if the components $\Phi_1, \dots, \Phi_{n-1}$ belong to I , and an element Φ of \mathcal{R}^X belongs to $\mathcal{R}_{\text{fix}}^X$ if and only if the component Φ_n satisfies $\Phi_n(x) = x_n + \theta(x)$ with $\theta \in I$. The corresponding tangent spaces are respectively given by

$$\mathcal{T}^X f = \left(\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_{n-1}} \right) I + \frac{\partial f}{\partial x_n} \underline{m} \quad \text{and} \quad \mathcal{T}_{\text{fix}}^X f = I J_f.$$

Example 2.3.3 (complete intersections with isolated singularities). In the case studied in [3], one has $\Sigma = \{0\}$ and $Y = \{0\}$. In this situation, [3] provides the implication (18) \implies (16) under the additional assumptions (17) and $\text{codim } X \leq 2$. Thus, our Theorem 2.1 removes the restriction on the codimension p of X . It also removes the assumption on $D_f^{-1}(\{0\})$ as soon as \mathcal{R}^X is replaced by $\mathcal{R}_{\text{fix}}^X$.

Problem. Is it possible to show that (18) implies (16) without the extra assumption (17), at least when $Y = \{0\}$? One might also hope for a result involving the subgroup of diffeomorphism-germs that induce bijections on $X \setminus Y$.

3. PROOFS

3.1. Proof of (13) \implies (16). We separate the estimates for ∇f and D_f .

3.1.1. Łojasiewicz estimate for ∇f . We follow, with suitable modifications, the proof of the corresponding inequality in [9]. Assume that the estimate does not hold. One can then find a sequence $(x_\nu)_{\nu \geq 1}$ of points of \mathbb{R}^n converging to 0 and such that

$$(19) \quad |\nabla f(x_\nu)| < \text{dist}(x_\nu, X \cup Y)^\nu \text{ for any } \nu \geq 1.$$

Extracting a subsequence if necessary, one can assume that

$$|x_{\nu+1}| < \frac{1}{3} \operatorname{dist}(x_\nu, X \cup Y)$$

(which implies, in particular, that $\operatorname{dist}(x_\nu, X \cup Y)$ decreases). Denote by B_ν the open euclidean ball centered at x_ν , with radius $\frac{1}{3} \operatorname{dist}(x_\nu, X \cup Y)$. Then the B_ν are pairwise disjoint, the open set $\mathcal{U} = \bigcup_{\nu \geq 1} B_\nu$ is contained in $\mathbb{R}^n \setminus (X \cup Y)$ and we have $\overline{\mathcal{U}} \cap (X \cup Y) = \{0\}$. Using a classical construction of cutoff functions (see, e.g. [6], chapter I, lemma 4.2), we also have a sequence $(\chi_\nu)_{\nu \geq 1}$ of C^∞ functions such that each χ_ν is supported in B_ν , identically equal to 1 in a neighborhood of the center x_ν , and satisfies, for any integer $j \geq 0$, any multi-index J of length j and any point x in \mathbb{R}^n , the estimate

$$(20) \quad |D^J \chi_\nu(x)| \leq C(j) \operatorname{dist}(x_\nu, X \cup Y)^{-j},$$

where $C(j)$ is a constant depending only on j . Beside this, Sard's lemma ensures the existence of a regular value c_ν of f such that

$$|f(x_\nu) - c_\nu| \leq \operatorname{dist}(x_\nu, X \cup Y)^\nu.$$

Consider the affine function u_ν given by $u_\nu(x) = f(x_\nu) - c_\nu + \langle \nabla f(x_\nu), x - x_\nu \rangle$. Taking into account (19) and (20), as well as the support condition for χ_ν , one checks that the series $u = \sum_{\nu \geq 1} \chi_\nu u_\nu$ defines an element of $\underline{m}_{X \cup Y}^\infty$ such that $u(x_\nu) = f(x_\nu) - c_\nu$ and $\nabla u(x_\nu) = \nabla f(x_\nu)$ for any $\nu \geq 1$. Since $\underline{m}_{X \cup Y}^\infty = \underline{m}_X^\infty \cap \underline{m}_Y^\infty$, we see, using (7) and (8), that u belongs to $\underline{m}_Y^\infty I^2$. By (13), one can therefore find a germ of diffeomorphism Φ such that $f \circ \Phi = f - u$. But this is impossible since each c_ν is a regular value of f and a singular value of $f - u$.

3.1.2. Łojasiewicz estimate for D_f . Assume that D_f does not satisfy $\mathfrak{L}(X, Y)$. One can then find a sequence $(y_\nu)_{\nu \geq 1}$ of points of X converging to 0 and such that

$$(21) \quad |D_f(y_\nu)| < \operatorname{dist}(y_\nu, Y)^\nu \quad \text{for any } \nu \geq 1.$$

Denote by $\lambda_1^{(\nu)}, \dots, \lambda_p^{(\nu)}$ the eigenvalues of the symmetric matrix $H_f(y_\nu)$, counted with multiplicities. Since we have $D_f(y_\nu) = \lambda_1^{(\nu)} \cdots \lambda_p^{(\nu)}$, the estimate (21) implies, for each $\nu \geq 1$, the existence of at least one index i_ν such that

$$(22) \quad |\lambda_{i_\nu}^{(\nu)}| < \operatorname{dist}(y_\nu, Y)^{\nu/p}.$$

Beside this, one can also find an orthogonal matrix P_ν such that

$$(23) \quad P_\nu^{-1} H_f(y_\nu) P_\nu = \operatorname{Diag}(\lambda_1^{(\nu)}, \dots, \lambda_p^{(\nu)}).$$

We define a $p \times p$ symmetric matrix V_ν by putting

$$(24) \quad P_\nu^{-1} V_\nu P_\nu = \operatorname{Diag}(0, \dots, 0, \lambda_{i_\nu}^{(\nu)}, 0, \dots, 0),$$

where $\lambda_{i_\nu}^{(\nu)}$ is at the i_ν -th position. Using the fact that $\|P_\nu^{-1}\| = \|P_\nu\| = 1$ for the euclidean norm, it is easy to see that the coefficients $v_{ij}^{(\nu)}$ of V_ν satisfy $|v_{ij}^{(\nu)}| \leq |\lambda_{i_\nu}^{(\nu)}|$, hence, by virtue of (22),

$$(25) \quad |v_{ij}^{(\nu)}| \leq \operatorname{dist}(y_\nu, Y)^{\nu/p}.$$

Now choose real numbers $\varepsilon_1^{(\nu)}, \dots, \varepsilon_p^{(\nu)}$ satisfying, for any $i = 1, \dots, p$,

$$(26) \quad \lambda_i^{(\nu)} \neq \varepsilon_i^{(\nu)} \quad \text{and} \quad |\varepsilon_i^{(\nu)}| < \operatorname{dist}(y_\nu, Y)^\nu.$$

Then define a $p \times p$ symmetric matrix W_ν by putting

$$(27) \quad P_\nu^{-1} W_\nu P_\nu = \text{Diag}(\varepsilon_1^{(\nu)}, \dots, \varepsilon_p^{(\nu)}).$$

In view of the second condition in (26), the same argument as for V_ν shows that the coefficients $w_{ij}^{(\nu)}$ of W_ν satisfy

$$(28) \quad |w_{ij}^{(\nu)}| \leq \text{dist}(y_\nu, Y)^\nu.$$

We can also assume that $|y_{\nu+1}| < \frac{1}{3} \text{dist}(y_\nu, Y)$ and consider the open ball C_ν centered at y_ν with radius $\frac{1}{3} \text{dist}(y_\nu, Y)$. With each C_ν we associate a cutoff function η_ν supported in C_ν , identically equal to 1 in a neighborhood of y_ν , and such that

$$(29) \quad |D^J \eta_\nu(x)| \leq C(j) \text{dist}(y_\nu, Y)^{-j}$$

for any integer $j \geq 0$, any multi-index J of length j and any $x \in \mathbb{R}^n$. The open set $\mathcal{V} = \bigcup_{\nu \geq 1} C_\nu$ is contained in $\mathbb{R}^n \setminus Y$ and satisfies $\overline{\mathcal{V}} \cap Y = \{0\}$. By (25), (28) and the support properties of η_ν , both series $V = \sum_{\nu \geq 1} \eta_\nu V_\nu$ and $W = \sum_{\nu \geq 1} \eta_\nu W_\nu$ define symmetric $p \times p$ matrices whose respective coefficients v_{ij} and w_{ij} all belong to \underline{m}_Y^∞ . Therefore, the functions $v = \sum_{1 \leq i, j \leq p} v_{ij} \psi_i \psi_j$ and $w = \sum_{1 \leq i, j \leq p} w_{ij} \psi_i \psi_j$ belong to $\underline{m}_Y^\infty I^2$ and, by assumption, one can then find Φ_v and Φ_w in $\mathcal{R}_{\text{fix}}^{X \setminus Y}$ such that

$$(30) \quad f - v = f \circ \Phi_v \quad \text{and} \quad f - w = f \circ \Phi_w.$$

Beside this, we have $H_{f-v}(y_\nu) = H_f(y_\nu) - V(y_\nu) = H_f(y_\nu) - V_\nu$ by construction. Therefore, (23) and (24) imply that 0 is an eigenvalue of $H_{f-v}(y_\nu)$, hence $D_{f-v}(y_\nu) = 0$. Thanks to (12), we can now use Lemma 1.3.3 with $Z = X \setminus Y$ and $x = y_\nu$. Thus, (30) implies

$$(31) \quad D_f(\Phi_v(y_\nu)) = 0 \quad \text{for any } \nu \geq 1.$$

We similarly have $H_{f-w}(y_\nu) = H_f(y_\nu) - W(y_\nu) = H_f(y_\nu) - W_\nu$, hence $D_{f-w}(y_\nu) = \prod_{i=1}^p (\lambda_i^{(\nu)} - \varepsilon_i^{(\nu)}) \neq 0$ by (23), (27) and the first condition in (26). By (30) and Lemma 1.3.3, we now get

$$(32) \quad D_f(\Phi_w(y_\nu)) \neq 0 \quad \text{for any } \nu \geq 1.$$

Finally, recall that the diffeomorphisms Φ_v and Φ_w provided by (13) preserve $X \setminus Y$ pointwise, hence $\Phi_v(y_\nu) = \Phi_w(y_\nu) = y_\nu$ and (31) contradicts (32). The proof is complete.

3.2. Proof of (16) \implies (15). We follow the pattern of the corresponding proof in [9], with some necessary adaptations. First, one checks that K_f contains all the germs $\psi_i^{p-2} \partial f / \partial x_j$ with $1 \leq i \leq p$ and $1 \leq j \leq n$: it suffices to compute the determinant whose first column is given by h^j and the other columns by the trivial relations $\psi_i e_k - \psi_k e_i$ with $k \neq i$. Therefore, K_f contains the germ $v = |\psi|^{2(p-2)} |\nabla f|^2$. For $0 < \varepsilon < 1$ and $s \geq 1$, now put

$$V_{\varepsilon,s} = \{x \in \mathbb{R}^n; \text{dist}(x, X) \geq \varepsilon \text{dist}(x, Y)^s\} \quad \text{and} \quad W_{\varepsilon,s} = \mathbb{R}^n \setminus V_{\varepsilon,s}.$$

Since $|\psi|^2$ satisfies $\mathfrak{L}(\mathbb{R}^n, X)$ by the classical Łojasiewicz inequality for analytic functions, and since ∇f satisfies $\mathfrak{L}(\mathbb{R}^n, X \cup Y)$ by assumption, we see that for any given choice of ε and s , the germ v satisfies $\mathfrak{L}(V_{\varepsilon,s}, Y)$.

Now, let \mathcal{M} be the set of $p \times p$ minors of the jacobian matrix ψ' . Using (10) and elementary linear algebra, one obtains, for any μ in \mathcal{M} ,

$$(33) \quad 2^p \mu \det(f_{ij}) = a_\mu + b_\mu,$$

where a_μ is a suitable $p \times p$ minor of the matrix Λ defined in Subsection 1.4, hence an element of K_f , and b_μ is an element of I . Put $w = \sum_{\mu \in \mathcal{M}} a_\mu^2$ and, for any $x \in (\mathbb{R}^n, 0)$, denote by \hat{x} a point of X such that $|x - \hat{x}| = \text{dist}(x, X)$. We obviously have

$$(34) \quad |w(x) - w(\hat{x})| \leq C \text{dist}(x, X)$$

for some constant $C > 0$. By (33), we also have

$$(35) \quad w(\hat{x}) = 2^{2p} (D_f(\hat{x}))^2 \sum_{\mu \in \mathcal{M}} (\mu(\hat{x}))^2$$

since each b_μ vanishes on X . Now, by (6) and (12), there exist constants $C' > 0$ and $\alpha \geq 1$ such that $\sum_{\mu \in \mathcal{M}} (\mu(\hat{x}))^2 \geq C' \text{dist}(\hat{x}, \Sigma)^\alpha \geq C' \text{dist}(\hat{x}, Y)^\alpha$. Beside this, D_f satisfies $\mathfrak{L}(X, Y)$ by assumption. From (35) and these remarks, we get

$$(36) \quad |w(\hat{x})| \geq C'' \text{dist}(\hat{x}, Y)^\beta$$

for some suitable constants $C'' > 0$ and $\beta \geq 1$. Assume now that x belongs to some set $W_{\epsilon, s}$. In this situation, we have $|\text{dist}(x, Y) - \text{dist}(\hat{x}, Y)| \leq |x - \hat{x}| \leq \epsilon \text{dist}(x, Y)^s$, therefore $\text{dist}(\hat{x}, Y)^\beta$ can be replaced by $\text{dist}(x, Y)^\beta$ in (36), up to a modification of C'' . From (34) and (36), we derive that w satisfies $\mathfrak{L}(W_{\epsilon, s}, Y)$ for $s > \beta$. Finally, $v + w$ is an element of K_f which satisfies $\mathfrak{L}(\mathbb{R}^n, Y)$, and proposition V.4.3 of [10] yields the conclusion.

3.3. Proof of (15) \implies (14). It is an immediate consequence of (11).

3.4. Proof of (14) \implies (13). This can be proved by a Mather-type homotopy argument, using Nakayama's lemma and tangent spaces as in [4] for instance. Alternatively, here we use the somewhat shorter approach based on Tougeron's implicit function theorem; see [10], theorem III.3.2 and remark VIII.3.7.1. Let u be an element of $\underline{m}_Y^\infty I^2$ and put $g = f + u$. For x and y in $(\mathbb{R}^n, 0)$, define $F(x, y) = f(x + y) - g(x)$. Put $F_i(x) = \partial F / \partial y_i(x, 0)$ for $i = 1, \dots, n$. Since $F_i = \partial f / \partial x_i$, it turns out that $(F_1, \dots, F_n) \mathcal{E}_n = J_f$. Now remark that $\underline{m}_Y^\infty = \underline{m}_Y^\infty \underline{m}_Y^\infty$ by proposition V.2.3 of [10], hence $\underline{m}_Y^\infty I^2 = \underline{m}_Y^\infty (\underline{m}_Y^\infty I)^2$. Therefore, (14) implies that $F(\cdot, 0)$ belongs to $\underline{m}_Y^\infty J_f^2$. Tougeron's implicit function theorem yields a map $\varphi : (\mathbb{R}^n, 0) \longrightarrow (\mathbb{R}^n, 0)$ with components in $\underline{m}_Y^\infty J_f$, hence in $\underline{m}_Y^\infty I$, such that $F(x, \varphi(x)) = 0$. Put $\Phi(x) = x + \varphi(x)$. Clearly Φ is a germ of diffeomorphism at the origin, and it coincides with the identity on $X \cup Y$. In particular it belongs to $\mathcal{R}_{\text{fix}}^{X \setminus Y}$. This ends the proof, since we also have $f \circ \Phi = g$ by construction.

3.5. Proof of (13) \iff (18) under the extra assumption (17). The implication (13) \implies (18) is obvious (and does not require (17)). In order to prove the converse, it suffices to show that (17) together with (18) imply (16). Now, a simple inspection of the proof of the implication (13) \implies (16) reveals that the pointwise preservation of $X \setminus Y$ is used only to obtain a contradiction between (31) and (32). If (13) is replaced by (18), we obtain in the same way (31) for some Φ_v preserving $X \setminus Y$ globally. Taking (17) into account, we obtain another contradiction, since D_f has no zero on $X \setminus Y$, hence no zero at $\Phi_v(y_\nu)$. The result follows.

REFERENCES

- [1] J. Bochnak and J.-J. Risler, *Sur les exposants de Łojasiewicz*, Comment. Math. Helvetici **50** (1975), 493–507. MR0404674 (53:8474)
- [2] D. Eisenbud, *Commutative Algebra with a view toward Algebraic Geometry*, Graduate Texts in Math. 150, Springer-Verlag (1995). MR1322960 (97a:13001)
- [3] V. Grandjean, *Infinite relative determinacy of smooth function germs with transverse isolated singularities and relative Łojasiewicz conditions*, J. London Math. Soc. **69** (2004), 518–530. MR2040619 (2005a:58060)
- [4] S. Izumiya and S. Matsuoka, *Notes on smooth function germs on varieties*, Proc. Amer. Math. Soc. **97** (1986), 146–150. MR0831404 (87g:58013)
- [5] L. Kushner and B. Terra Leme, *Finite relative determination and relative stability*, Pacific J. Math. **192** (2000), 315–328. MR1744573 (2000m:58069)
- [6] B. Malgrange, *Ideals of Differentiable Functions*, Oxford University Press (1965).
- [7] T.C. Nguyen, H.D. Nguyen, S.M. Nguyen, and H.V. Ha, *Sur les germes de fonctions infiniment déterminés*, C. R. Acad. Sci. Paris **285** (1977), 1045–1048. MR0649266 (58:31180)
- [8] D. Siersma, *Isolated line singularities*, Proc. Symp. Pure Math. **40** (1983), 485–495. MR0713274 (85d:32017)
- [9] B. Sun and L.C. Wilson, *Determinacy of smooth germs with real isolated line singularities*, Proc. Amer. Math. Soc. **129** (2001), 2789–2797. MR1838804 (2002d:58050)
- [10] J.-C. Tougeron, *Idéaux de fonctions différentiables*, Springer-Verlag (1972). MR0440598 (55:13472)
- [11] C.T.C. Wall, *Finite determinacy of smooth map-germs*, Bull. London Math. Soc. **13** (1981), 481–539. MR0634595 (83i:58020)
- [12] L.C. Wilson, *Infinitely determined mapgerms*, Can. J. Math. **33** (1981), 671–684. MR0627650 (82k:58023)
- [13] L.C. Wilson, personal communication (October 2003).

MATHÉMATIQUES - BÂTIMENT M2, UNIVERSITÉ DES SCIENCES ET TECHNOLOGIES DE LILLE,
F-59655 VILLENEUVE D'ASCQ CEDEX, FRANCE

E-mail address: thilliez@math.univ-lille1.fr