

A CHARACTERIZATION OF MODULES LOCALLY OF FINITE INJECTIVE DIMENSION

RYO TAKAHASHI

(Communicated by Bernd Ulrich)

ABSTRACT. In this note, we characterize finite modules locally of finite injective dimension over commutative Noetherian rings in terms of vanishing of Ext modules.

1. INTRODUCTION

Let R be a commutative Noetherian ring. Goto [2] proved the following theorem.

Theorem 1.1 (Goto). *The following are equivalent:*

- (1) R is Gorenstein;
- (2) For every finite R -module M , there exists an integer n such that $\text{Ext}_R^i(M, R) = 0$ for all $i > n$.

We should note that this theorem remains valid even in the case where the ring R has infinite Krull dimension.

The purpose of this note is to give a characterization of finite modules locally of finite injective dimension. Our theorem is the following.

Theorem 1.2. *The following are equivalent for a finite R -module N :*

- (1) $\text{id}_{R_{\mathfrak{p}}} N_{\mathfrak{p}} < \infty$ for every $\mathfrak{p} \in \text{Spec } R$;
- (2) For every finite R -module M , there exists an integer n such that $\text{Ext}_R^i(M, N) = 0$ for all $i > n$.

This theorem is a generalization of Goto's. In fact, applying our theorem to $N = R$, we immediately obtain Goto's theorem.

2. PROOF OF THE THEOREM

We denote by $\text{CM}(R)$ the *Cohen-Macaulay locus* of R , that is, the set of prime ideals \mathfrak{p} of R such that the local ring $R_{\mathfrak{p}}$ is Cohen-Macaulay. The following lemma can be shown in a similar way to the proof of [3, Theorem 24.5].

Lemma 2.1. *Let \mathfrak{p} be a prime ideal of R such that both $R_{\mathfrak{p}}$ and R/\mathfrak{p} are Cohen-Macaulay rings. Then there exists an element $f \in R - \mathfrak{p}$ such that $D(f) \cap V(\mathfrak{p}) \subseteq \text{CM}(R)$.*

Received by the editors March 13, 2006 and, in revised form, August 19, 2006.

2000 *Mathematics Subject Classification.* Primary 13D05, 13D07.

Key words and phrases. Injective dimension, Cohen-Macaulay locus.

©2007 American Mathematical Society
Reverts to public domain 28 years from publication

It is known that the Cohen-Macaulay locus of a homomorphic image of a Cohen-Macaulay ring is an open subset; see [3, Exercises 24.2]. To prove our theorem, we need to generalize this fact. For an ideal I of R , let $\text{CM}_R(R/I)$ denote the set of prime ideals $\mathfrak{p} \in V(I)$ such that the local ring $(R/I)_{\mathfrak{p}}$ is Cohen-Macaulay.

Lemma 2.2. *Let I and J be ideals of R . Suppose that $V(J)$ is contained in $\text{CM}(R)$. Then the following hold:*

- (1) *For any prime ideal $\mathfrak{p} \in \text{CM}_R(R/I) \cap V(J)$, there exists an element $f \in R - \mathfrak{p}$ such that $D(f) \cap V(\mathfrak{p}) \subseteq \text{CM}_R(R/I)$.*
- (2) *There exists an ideal K of R such that*

$$\text{CM}_R(R/I) \cap V(J) = D(K) \cap V(I) \cap V(J).$$

In other words, $\text{CM}_R(R/I) \cap V(J)$ is an open subset of $V(I) \cap V(J)$ in the relative topology induced by the Zariski topology of $\text{Spec } R$.

Proof. (1) Let $\mathfrak{p} \in \text{CM}_R(R/I) \cap V(J)$. Then by the assumption that $V(J)$ is contained in $\text{CM}(R)$, the ring $R_{\mathfrak{p}}$ is a Cohen-Macaulay local ring. Making a similar argument to the proof of [3, Theorem 24.5], we can assume without loss of generality that there is an R -regular sequence $\mathbf{x} = x_1, x_2, \dots, x_n$ in \mathfrak{p} with $\mathfrak{p}^r \subseteq \mathbf{x}R$ for some $r > 0$ and that $\overline{\mathfrak{p}}^i / \overline{\mathfrak{p}}^{i+1}$ is a free $\overline{R} / \overline{\mathfrak{p}}$ -module for all $i > 0$, where $\overline{R} = R / \mathbf{x}R$ and $\overline{\mathfrak{p}} = \mathfrak{p} / \mathbf{x}R$.

We have only to prove that the residue ring R / \mathfrak{p} is Cohen-Macaulay. In fact, if R / \mathfrak{p} is Cohen-Macaulay, then so are $(R/I) / (\mathfrak{p}/I)$ and $(R/I)_{\mathfrak{p}/I}$ since \mathfrak{p} is in $\text{CM}_R(R/I)$. Hence Lemma 2.1 implies that there is an element $f \in R - \mathfrak{p}$ such that $D(\overline{f}) \cap V(\mathfrak{p}/I)$ is contained in $\text{CM}(R/I)$, where \overline{f} denotes the residue class of f in R/I . We easily see that $D(f) \cap V(\mathfrak{p})$ is contained in $\text{CM}_R(R/I)$.

Let us show that R / \mathfrak{p} is a Cohen-Macaulay ring. It is easy to see from [3, Exercises 24.1] that $R / \mathfrak{p} = \overline{R} / \overline{\mathfrak{p}}$ is Cohen-Macaulay if and only if so is \overline{R} . Take a prime ideal $\mathfrak{q} \in V(\mathbf{x}) = V(\mathfrak{p})$. Then we have $\mathfrak{q} \supseteq \mathfrak{p} \supseteq J$; hence $\mathfrak{q} \in V(J) \subseteq \text{CM}(R)$. Therefore $R_{\mathfrak{q}}$ is a Cohen-Macaulay local ring, and so is $\overline{R}_{\mathfrak{q}}$, because \mathbf{x} is an $R_{\mathfrak{q}}$ -regular sequence. This shows that \overline{R} is a Cohen-Macaulay ring. Thus we conclude that the residue ring R / \mathfrak{p} is Cohen-Macaulay, as desired.

(2) Set $U = \{ \mathfrak{p}/I + J \mid \mathfrak{p} \in \text{CM}_R(R/I) \cap V(J) \}$. This is a subset of $\text{Spec } R/I + J$. Note that this subset is stable under generalization. Let $P \in U$. Then there is a prime ideal $\mathfrak{p} \in \text{CM}_R(R/I) \cap V(J)$ such that $P = \mathfrak{p}/I + J$. By the assertion (1) of the lemma, the set $D(f) \cap V(\mathfrak{p})$ is contained in $\text{CM}_R(R/I)$ for some $f \in R - \mathfrak{p}$. Denote by \overline{f} the residue class of f in R/I . It is easy to see that $P \in D(\overline{f}) \cap V(P) \subseteq U$. Thus U contains a nonempty open subset of $V(P)$. By virtue of the topological Nagata criterion [3, Theorem 24.2], U is an open subset of $\text{Spec } R/I + J$; we have $U = D(K/I + J)$ for some ideal K of R containing $I + J$. Then it is easily checked that $\text{CM}_R(R/I) \cap V(J) = D(K) \cap V(I) \cap V(J)$. □

Now, we can prove our theorem.

Proof of Theorem 1.2. (2) \Rightarrow (1): Let \mathfrak{p} be a prime ideal of R . Then there is an integer n such that $\text{Ext}_R^i(R/\mathfrak{p}, N) = 0$ for all $i > n$. Hence we have $\text{Ext}_{R_{\mathfrak{p}}}^i(\kappa(\mathfrak{p}), N_{\mathfrak{p}}) = 0$ for all $i > n$. Therefore by [1, Theorem 3.1.14] we obtain $\text{id}_{R_{\mathfrak{p}}} N_{\mathfrak{p}} \leq n < \infty$.

(1) \Rightarrow (2): First of all, note that (2) is equivalent to the statement that for each ideal I of R there is an integer n such that $\text{Ext}_R^i(R/I, N) = 0$ for all $i > n$. (This can easily be proved by induction on the number of generators of the R -module

M.) Suppose that there exists an ideal I of R such that for any integer n there is an integer $i > n$ such that $\text{Ext}_R^i(R/I, N) \neq 0$. We want to derive a contradiction. Since R is Noetherian, one can choose I to be a maximal one among such ideals. Making a similar argument to the proof of Theorem 1.1, we see that the ideal I is prime and that for any element $f \in R - I$, the map

$$\text{Ext}_R^i(R/I, N) \xrightarrow{f} \text{Ext}_R^i(R/I, N)$$

is an isomorphism for $i \gg 0$.

Claim 1. *One has $I \in \text{Supp}_R N \subseteq \text{CM}(R)$.*

Proof of Claim. Our assumption (1) implies that for any $\mathfrak{p} \in \text{Supp}_R N$, the nonzero finite $R_{\mathfrak{p}}$ -module $N_{\mathfrak{p}}$ has finite injective dimension. Hence $R_{\mathfrak{p}}$ is a Cohen-Macaulay local ring; see [1, Corollary 9.6.2 and Remarks 9.6.4]. Thus $\text{Supp}_R N$ is contained in $\text{CM}(R)$. On the other hand, assume that I is not in $\text{Supp}_R N$. Then there exists an element $f \in \text{Ann}_R N - I$, and the map $\text{Ext}_R^i(R/I, N) \xrightarrow{f} \text{Ext}_R^i(R/I, N)$ is an isomorphism for $i \gg 0$. Since $fN = 0$, this map is the zero map, and we get $\text{Ext}_R^i(R/I, N) = 0$ for $i \gg 0$. It follows from this contradiction that I belongs to $\text{Supp}_R N$. \square

Noting that $\text{Supp}_R N = V(\text{Ann}_R N)$, we see from Claim 1 and Lemma 2.2(2) that there is an ideal K of R such that $\text{CM}_R(R/I) \cap \text{Supp}_R N = D(K) \cap V(I) \cap \text{Supp}_R N$. The localization $(R/I)_I = \kappa(I)$ is a field, hence a Cohen-Macaulay ring. It is seen from Claim 1 again that $I \in \text{CM}_R(R/I) \cap \text{Supp}_R N \subseteq D(K)$. Thus there is an element $f \in K - I$.

Claim 2. *For any prime ideal $\mathfrak{p} \in D(f)$ and any integer $i > \text{ht } I$, one has $\text{Ext}_{R_{\mathfrak{p}}}^i(R_{\mathfrak{p}}/IR_{\mathfrak{p}}, N_{\mathfrak{p}}) = 0$.*

Proof of Claim. We may assume that \mathfrak{p} belongs to both $V(I)$ and $\text{Supp}_R N$, because otherwise the module $\text{Ext}_{R_{\mathfrak{p}}}^i(R_{\mathfrak{p}}/IR_{\mathfrak{p}}, N_{\mathfrak{p}})$ automatically vanishes. Hence Claim 1 implies that \mathfrak{p} belongs to $\text{CM}(R)$, namely the local ring $R_{\mathfrak{p}}$ is Cohen-Macaulay. Added to it, since $D(f)$ is contained in $D(K)$, we have $\mathfrak{p} \in D(K) \cap V(I) \cap \text{Supp}_R N \subseteq \text{CM}_R(R/I)$, and therefore $R_{\mathfrak{p}}/IR_{\mathfrak{p}}$ is Cohen-Macaulay. Thus we get the following equalities:

$$\text{depth } R_{\mathfrak{p}} - \text{depth } R_{\mathfrak{p}}/IR_{\mathfrak{p}} = \dim R_{\mathfrak{p}} - \dim R_{\mathfrak{p}}/IR_{\mathfrak{p}} = \text{ht } IR_{\mathfrak{p}} = \text{ht } I.$$

Since $N_{\mathfrak{p}}$ is a finite $R_{\mathfrak{p}}$ -module of finite injective dimension by assumption, it follows from the result of Ischebeck [1, Exercises 3.1.24] that $\text{Ext}_{R_{\mathfrak{p}}}^i(R_{\mathfrak{p}}/IR_{\mathfrak{p}}, N_{\mathfrak{p}}) = 0$ for every $i > \text{ht } I$. \square

Claim 2 means that $(\text{Ext}_{R_f}^i(R_f/IR_f, N_f))_{\mathfrak{P}} = 0$ for every $\mathfrak{P} \in \text{Spec } R_f$ and every $i > \text{ht } I$. Therefore, $\text{Ext}_{R_f}^i(R_f/IR_f, N_f) = 0$ for $i > \text{ht } I$. The R -module $\text{Ext}_R^i(R/I, N)$ is isomorphic to $\text{Ext}_{R_f}^i(R_f/IR_f, N_f)$ for $i \gg 0$, and thus $\text{Ext}_R^i(R/I, N) = 0$ for $i \gg 0$. This contradiction completes the proof of our theorem. \square

REFERENCES

[1] BRUNS, W.; HERZOG, J. Cohen-Macaulay rings, revised edition. Cambridge Studies in Advanced Mathematics, 39. Cambridge University Press, Cambridge, 1998. MR1251956 (95h:13020)

- [2] GOTO, S. Vanishing of $\text{Ext}_A^i(M, A)$. *J. Math. Kyoto Univ.* **22** (1982/83), no. 3, 481–484. MR674605 (84c:13019)
- [3] MATSUMURA, H. Commutative ring theory. Translated from the Japanese by M. Reid. Cambridge Studies in Advanced Mathematics, 8. *Cambridge University Press, Cambridge*, 1986. MR879273 (88h:13001)

DEPARTMENT OF MATHEMATICS, SCHOOL OF SCIENCE AND TECHNOLOGY, MEIJI UNIVERSITY,
1-1-1 HIGASHIMITA, TAMA-KU, KAWASAKI 214-8571, JAPAN
E-mail address: takahasi@math.meiji.ac.jp