

A MAXIMAL FUNCTION CHARACTERIZATION OF THE HARDY SPACE FOR THE GAUSS MEASURE

GIANCARLO MAUCERI, STEFANO MEDA, AND PETER SJÖGREN

(Communicated by Michael T. Lacey)

ABSTRACT. An atomic Hardy space $H^1(\gamma)$ associated to the Gauss measure γ in \mathbb{R}^n has been introduced by the first two authors. We first prove that it is equivalent to use $(1, r)$ - or $(1, \infty)$ -atoms to define this $H^1(\gamma)$. For $n = 1$, a maximal function characterization of $H^1(\gamma)$ is found. In arbitrary dimension, we give a description of the nonnegative functions in $H^1(\gamma)$ and use it to prove that $L^p(\gamma) \subset H^1(\gamma)$ for $1 < p \leq \infty$.

1. INTRODUCTION

Denote by γ the Gauss measure on \mathbb{R}^n , i.e., the probability measure with density $\gamma_0(x) = \pi^{-n/2} e^{-|x|^2}$ with respect to the Lebesgue measure λ . Harmonic analysis on the measured metric space $(\mathbb{R}^n, d, \gamma)$, where d denotes the Euclidean distance on \mathbb{R}^n , has been the object of many investigations. In particular, efforts have been made to study operators related to the Ornstein–Uhlenbeck semigroup, with emphasis on maximal operators [33, 16, 27, 13, 21, 22], Riesz transforms [29, 14, 28, 32, 30, 17, 15, 8, 9, 10, 31, 35, 7, 24] and functional calculus [11, 12, 18, 25].

In [24] the first two authors defined an atomic Hardy-type space $H^1(\gamma)$ and a space $BMO(\gamma)$ of functions of bounded mean oscillation, associated to γ . We briefly recall their definitions. A closed Euclidean ball B is called *admissible* at scale $s > 0$ if

$$r_B \leq s \min(1, 1/|c_B|);$$

here and in the sequel r_B and c_B denote the radius and the centre of B , respectively. We denote by \mathcal{B}_s the family of all balls admissible at scale s . For the sake of brevity, we shall refer to balls in \mathcal{B}_1 simply as admissible balls. Further, B will be called *maximal admissible* if $r_B = \min(1, 1/|c_B|)$.

Now let $r \in (1, \infty]$. A *Gaussian* $(1, r)$ -atom is either the constant function 1 or a function a in $L^r(\gamma)$ supported in an admissible ball B and such that

$$(1.1) \quad \int a \, d\gamma = 0 \quad \text{and} \quad \|a\|_r \leq \gamma(B)^{1/r-1};$$

here and in the whole paper, $\|\cdot\|_r$ denotes the norm in $L^r(\gamma)$. In the latter case, we say that the atom a is associated to the ball B . The space $H^{1,r}(\gamma)$ is then the vector

Received by the editors February 10, 2011 and, in revised form, September 7, 2011.

2010 *Mathematics Subject Classification*. Primary 42B30, 42B35; Secondary 42C10.

Key words and phrases. Gaussian measure, Gaussian Hardy space, maximal function, atomic decomposition.

This work was partially supported by PRIN 2009 “Analisi Armonica”.

space of all functions f in $L^1(\gamma)$ that admit a decomposition of the form $\sum_j \lambda_j a_j$, where the a_j are Gaussian $(1, r)$ -atoms and the sequence of complex numbers $\{\lambda_j\}$ is summable. The norm of f in $H^{1,r}(\gamma)$ is defined as the infimum of $\sum_j |\lambda_j|$ over all representations of f as above.

In [24] the spaces $H^{1,r}(\gamma)$ were defined and proved to coincide for all $1 < r < \infty$, with equivalent norms. In Section 2 we complement this by proving that they coincide also with the space $H^{1,\infty}(\gamma)$. Once this is established, we shall denote the space by $H^1(\gamma)$ and use the $H^{1,\infty}(\gamma)$ norm. Further, we shall frequently write atom for the $(1, \infty)$ -atom.

The space $BMO(\gamma)$ consists of all functions f in $L^1(\gamma)$ such that

$$\sup_{B \in \mathcal{B}_1} \frac{1}{\gamma(B)} \int_B |f - f_B| \, d\gamma < \infty,$$

where f_B denotes the mean value of f on B , taken with respect to the Gauss measure. The norm of a function in $BMO(\gamma)$ is

$$\|f\|_{BMO(\gamma)} = \|f\|_1 + \sup_{B \in \mathcal{B}_1} \frac{1}{\gamma(B)} \int_B |f - f_B| \, d\gamma.$$

If, in the definitions of $H^1(\gamma)$ and $BMO(\gamma)$, we replace the family \mathcal{B}_1 of admissible balls at scale 1 by \mathcal{B}_s for any fixed $s > 0$, we obtain the same spaces with equivalent norms; see [24]. We remark that a similar $H^1 - BMO$ theory for more general measured metric spaces has been developed by A. Carbonaro and the first two authors in [1, 2, 3]. See also the papers [19, 20] of L. Liu and D. Yang for related results.

The main motivation for introducing these two spaces was to provide endpoint estimates for singular integrals associated to the Ornstein-Uhlenbeck operator $\mathcal{L} = -(1/2)\Delta + x \cdot \nabla$, a natural self-adjoint Laplacian on $L^2(\gamma)$. Indeed, in [24] the first two authors proved that the imaginary powers of \mathcal{L} are bounded from $H^1(\gamma)$ to $L^1(\gamma)$ and from $L^\infty(\gamma)$ to $BMO(\gamma)$ and that Riesz transforms of the form $\nabla^\alpha \mathcal{L}^{-|\alpha|}$ and of any order are bounded from $L^\infty(\gamma)$ to $BMO(\gamma)$. In a recent paper [26], the authors proved that boundedness from $H^1(\gamma)$ to $L^1(\gamma)$ and from $L^\infty(\gamma)$ to $BMO(\gamma)$ holds for any first-order Riesz transform in dimension one, but not always in higher dimensions.

The definition of the space $H^1(\gamma)$ closely resembles the atomic definition of the classical Hardy space $H^1(\lambda)$ on \mathbb{R}^n endowed with the Lebesgue measure λ , but there are two basic differences. First, the measured metric space $(\mathbb{R}^n, d, \gamma)$ is nondoubling. Further, except for the constant atom, a Gaussian atom must have “small support”, i.e., support contained in an admissible ball. Despite these differences, $H^1(\gamma)$ shares many of the properties of $H^1(\lambda)$. In particular, the topological dual of $H^1(\gamma)$ is isomorphic to $BMO(\gamma)$, an inequality of John-Nirenberg type holds for functions in $BMO(\gamma)$ and the spaces $L^p(\gamma)$ are intermediate spaces between $H^1(\gamma)$ and $BMO(\gamma)$ for the real and the complex interpolation methods.

It is well known that the classical Hardy space $H^1(\lambda)$ can be defined in at least three different ways: the *atomic* definition, the *maximal* definition and the definition based on *Riesz transforms* [6, 34].

As shown in [26], in higher dimensions the first-order Ornstein-Uhlenbeck Riesz transforms $\partial_j \mathcal{L}^{-1/2}$ are unbounded from $H^1(\gamma)$ to $L^1(\gamma)$; here $\partial_j = \partial/\partial x_j$. Thus $H^1(\gamma)$ does not coincide with the space of all functions in $L^1(\gamma)$ such that $\partial_j \mathcal{L}^{-1/2} f \in L^1(\gamma)$ for $j = 1, \dots, n$.

This paper arose from the desire to find a maximal characterization of the space $H^1(\gamma)$. We recall that the classical space $H^1(\lambda)$ can be characterized as the space of all functions f in $L^1(\lambda)$ whose *grand maximal function*

$$(1.2) \quad \mathcal{M}f(x) = \sup \{ |\phi_t * f(x)| : \phi \in \Phi, t > 0 \}$$

is also in $L^1(\lambda)$. Here $\Phi = \{ \phi \in C_c^1(B(0, 1)) : |D^\alpha \phi| \leq 1 \text{ for } |\alpha| = 0, 1 \}$ and $\phi_t(x) = t^{-n} \phi(x/t)$.

To characterize $H^1(\gamma)$, we introduce the *local grand maximal function* defined on $L^1_{\text{loc}}(\mathbb{R}^n, \gamma)$ by

$$\mathcal{M}_{\text{loc}}f(x) = \sup \{ |\phi_t * f(x)| : \phi \in \Phi, 0 < t < \min(1, 1/|x|) \}.$$

In Section 3 we shall prove that, in arbitrary dimensions, $f \in H^1(\gamma)$ implies $\mathcal{M}_{\text{loc}}f \in L^1(\gamma)$. Moreover, *in dimension one*, $H^1(\gamma)$ can be characterized as the space of all functions f in $L^1(\gamma)$ satisfying $\mathcal{M}_{\text{loc}}f \in L^1(\gamma)$ and the following additional *global* condition:

$$(1.3) \quad E(f) = \int_0^\infty x \left(\left| \int_x^\infty f \, d\gamma \right| + \left| \int_{-\infty}^{-x} f \, d\gamma \right| \right) \, d\lambda(x) < \infty.$$

This is Theorem 3.3 below.

Roughly speaking, if we interpret a function f as a density of electrical charge on the real line, this global condition says that the positive and negative charges nearly balance out, so that the net charges inside the intervals $(-\infty, -x)$ and (x, ∞) decay sufficiently fast as x approaches $+\infty$. The condition is violated when the distance between the positive and the negative charges increases too much or the charges do not decay sufficiently fast at infinity. For instance, let $(a_n)_1^\infty$ and $(a'_n)_1^\infty$ be increasing sequences in $(2, \infty)$ such that

$$a_n + 2/a_n < a'_n \quad \text{and} \quad a'_n + 2/a'_n < a_{n+1} < 2a_n$$

for all n . Then set

$$(1.4) \quad f = \sum_1^\infty c_n \left(\frac{\chi_{(a_n, a_n + 1/a_n)}}{\gamma(a_n, a_n + 1/a_n)} - \frac{\chi_{(a'_n, a'_n + 1/a'_n)}}{\gamma(a'_n, a'_n + 1/a'_n)} \right)$$

for some $c_n > 0$. One easily verifies that $\mathcal{M}_{\text{loc}}f \in L^1(\gamma)$ if and only if $\sum c_n < \infty$. But the global condition $E(f) < \infty$ is equivalent to $\sum c_n a_n (a'_n - a_n) < \infty$, which is here a stronger condition.

We have not been able to find a similar characterization of $H^1(\gamma)$ in higher dimensions. However, in Section 4 we prove in all dimensions that if $\mathcal{M}_{\text{loc}}f \in L^1(\gamma)$ and the function f satisfies the stronger global condition

$$E_+(f) = \int |x|^2 |f(x)| \, d\gamma(x) < \infty,$$

then $f \in H^1(\gamma)$. Observe that for $n = 1$ and $f \geq 0$, Fubini's theorem implies that the conditions $E(f) < \infty$ and $E_+(f) < \infty$ are equivalent. In arbitrary dimensions, $E_+(f)$ can be used to characterize the nonnegative functions in $H^1(\gamma)$; see Theorem 4.2. This also leads to a simple proof of the inclusions $L^p(\gamma) \subset H^1(\gamma)$ and $BMO(\gamma) \subset L^p(\gamma)$ for $1 < p \leq \infty$.

We end the introduction with some technical observations and notation. In the following we use repeatedly the fact that on admissible balls at a fixed scale s ,

the Gauss and the Lebesgue measures are equivalent; i.e., there exists a positive constant $C(s)$ such that for every measurable subset E of $B \in \mathcal{B}_s$,

$$(1.5) \quad C(s)^{-1}\gamma(E) \leq \gamma_0(c_B)\lambda(E) \leq C(s)\gamma(E).$$

In particular this implies that the Gauss measure is doubling on balls in \mathcal{B}_s , with a constant that depends on s (see [24, Prop. 2.1]). Further, it is straightforward to see that if $B' \subset B$ are two balls and $B \in \mathcal{B}_s$, then B' is also in \mathcal{B}_s .

Given a ball B in \mathbb{R}^n and a positive number ρ , we shall denote by ρB the ball with the same centre and with radius ρr_B .

In the following C denotes a positive constant whose value may change from occurrence to occurrence and which depends only on the dimension n , except when otherwise explicitly stated.

2. COINCIDENCE OF $H^{1,\infty}(\gamma)$ AND $H^{1,r}(\gamma)$

First we need a lemma which will play a role also in the maximal characterization. It deals with the classical Hardy space $H^1(\lambda)$ with respect to the Lebesgue measure and the associated standard $(1, \infty)$ -atoms, called Lebesgue atoms below.

Lemma 2.1. *If $g \in H^1(\lambda)$ and the support of g is contained in a ball B , then g has an atomic decomposition $g = \sum_k \lambda_k a_k$, where the a_k are Lebesgue $(1, \infty)$ -atoms associated to balls contained in $2B$ and*

$$(2.1) \quad \sum_k |\lambda_k| \leq C \|g\|_{H^1(\lambda)}.$$

The lemma can be proved by applying [23, Theor. 4.13] to the space of homogeneous type B , endowed with Euclidean distance and Lebesgue measure. Some related results can be found in [4] and [5].

Theorem 2.2. *For every r in $(1, \infty)$, the spaces $H^{1,r}(\gamma)$ and $H^{1,\infty}(\gamma)$ coincide, with equivalent norms.*

Proof. In this proof, the constants C may depend on r and n . Since any Gaussian $(1, \infty)$ -atom is also a Gaussian $(1, r)$ -atom, $H^{1,\infty}(\gamma)$ is a subspace of $H^{1,r}(\gamma)$ and $\|f\|_{H^{1,r}(\gamma)} \leq \|f\|_{H^{1,\infty}(\gamma)}$. Conversely, suppose that a is a Gaussian $(1, r)$ -atom associated to the ball $B \in \mathcal{B}_1$. Then the function $a\gamma_0$ is a multiple of a Lebesgue $(1, r)$ -atom. Indeed, $\int a\gamma_0 d\lambda = \int a d\gamma = 0$ and, by the equivalence of the Gauss and Lebesgue measures on admissible balls,

$$\|a\gamma_0\|_{L^r(\lambda)} \leq C \lambda(B)^{1/r-1}.$$

Hence, $a\gamma_0$ is in $H^1(\lambda)$ with norm at most C . By Lemma 2.1, it has a decomposition

$$a\gamma_0 = \sum_j \lambda_j \alpha_j,$$

where each α_j is a Lebesgue $(1, \infty)$ -atom associated to a ball B_j contained in $2B$. Moreover

$$\sum_j |\lambda_j| \leq C,$$

and each B_j is admissible at scale 2. Define $a_j = \alpha_j \gamma_0^{-1}$. Then $\int a_j d\gamma = 0$, and by the equivalence of the Gauss and Lebesgue measures on B_j ,

$$\|a_j\|_\infty \leq C\gamma(B_j)^{-1}.$$

Thus the a_j are multiples of Gaussian $(1, \infty)$ -atoms. Since $a = \sum_j \lambda_j a_j$, we conclude that $a \in H^{1,\infty}(\gamma)$ and

$$(2.2) \quad \|a\|_{H^{1,\infty}(\gamma)} \leq C \sum_j |\lambda_j| \leq C. \quad \square$$

3. THE CHARACTERIZATION OF $H^1(\gamma)$ IN \mathbb{R}

In this section, we shall prove that $f \in H^1(\gamma)$ implies $\mathcal{M}_{\text{loc}}f \in L^1(\gamma)$ and that, in dimension one, functions in $H^1(\gamma)$ can be characterized by the two conditions $\mathcal{M}_{\text{loc}}f \in L^1(\gamma)$ and $E(f) < \infty$. We start with a simple but useful lemma dealing with the support of the local grand maximal function.

Lemma 3.1. *If $f \in L^1(\gamma)$ is supported in the admissible ball B , then $\text{supp } \mathcal{M}_{\text{loc}}f$ is contained in the ball $B' = B(c_B, R)$, where $R = 4 \min(1, 1/|c_B|)$.*

Proof. Let $x \in \text{supp } \mathcal{M}_{\text{loc}}f$. We write $\rho = |x|$ and $c = |c_B|$, so that $B \subset B(c_B, \min(1, 1/c))$. The balls B and $B(x, \min(1, 1/\rho))$ must intersect, and so

$$(3.1) \quad |x - c_B| \leq \min(1, 1/c) + \min(1, 1/\rho).$$

To prove the lemma, it is enough to show that

$$(3.2) \quad \min(1, 1/\rho) \leq 3 \min(1, 1/c),$$

since it then follows that $x \in B'$. Now $c - \rho \leq |x - c_B|$, so that (3.1) implies

$$c - \min(1, 1/c) \leq \rho + \min(1, 1/\rho).$$

Considering the cases $c \leq 1$ and $c > 1$, we conclude from this that

$$(3.3) \quad \max(1, c) - \min(1, 1/c) \leq \max(1, \rho) + \min(1, 1/\rho).$$

The function $t \mapsto t^{-1} - t$, $t > 0$, and its inverse are clearly decreasing. Considering the values of this function at $t = \min(1, 1/c)$ and $\min(1, 1/\rho)/3$, we see that (3.2) is equivalent to

$$\max(1, c) - \min(1, 1/c) \leq 3 \max(1, \rho) - \frac{1}{3} \min(1, 1/\rho).$$

Because of (3.3), this inequality follows if

$$\max(1, \rho) + \min(1, 1/\rho) \leq 3 \max(1, \rho) - \frac{1}{3} \min(1, 1/\rho)$$

or equivalently $\frac{4}{3} \min(1, 1/\rho) \leq 2 \max(1, \rho)$, which is trivially true. We have proved (3.2) and the lemma. \square

Lemma 3.2. *If f is in $H^1(\gamma)$, then $\mathcal{M}_{\text{loc}}f \in L^1(\gamma)$ and*

$$\|\mathcal{M}_{\text{loc}}f\|_1 \leq C \|f\|_{H^1(\gamma)}.$$

Proof. We shall prove that for any Gaussian atom a ,

$$(3.4) \quad \|\mathcal{M}_{\text{loc}}a\|_1 \leq C,$$

from which the lemma follows.

Since (3.4) is obvious if a is the constant function 1, we assume that a is associated to an admissible ball B . By the preceding lemma, $\text{supp } \mathcal{M}_{\text{loc}}f$ is contained in the ball denoted B' .

The integral of $\mathcal{M}_{\text{loc}}a$ over $2B$ with respect to γ is no larger than C , since $\mathcal{M}_{\text{loc}}a \leq C \sup |a| \leq C/\gamma(B)$. To estimate $\mathcal{M}_{\text{loc}}a$ at a point x in the remaining

set $B' \setminus 2B$, we take $\phi \in \Phi$ and $0 < t < \min(1, 1/|x|)$ and estimate $a * \phi_t(x)$. We can assume that $t > d(x, B)$ so that $t > |x - c_B|/2$, since otherwise $\phi_t * a(x)$ will vanish. Write

$$(3.5) \quad \phi_t * a(x) = t^{-n} \int \left(\phi\left(\frac{x-y}{t}\right) - \phi\left(\frac{x-c_B}{t}\right) \right) a(y) \, dy + t^{-n} \phi\left(\frac{x-c_B}{t}\right) \int a(y) \, dy.$$

Here the first term to the right can be estimated in a standard way by

$$Ct^{-n-1} \int_B |y - c_B| |a(y)| \, dy \leq C |x - c_B|^{-n-1} r_B \gamma_0(c_B)^{-1}.$$

To deal with the second term, we estimate $\int a(y) \, dy$, knowing that the integral of a against γ vanishes. Thus

$$\int a(y) \, dy = \int a(y) \frac{\gamma_0(c_B) - \gamma_0(y)}{\gamma_0(c_B)} \, dy.$$

The fraction appearing here is

$$(3.6) \quad e^{|c_B|^2} \left(e^{-|c_B|^2} - e^{-|y|^2} \right) = 1 - e^{(c_B-y) \cdot (c_B+y)},$$

and the last exponent stays bounded for $y \in B$. Thus the modulus of the right-hand side of (3.6) is at most $C|c_B - y||c_B + y| \leq Cr_B(1 + |c_B|)$. Since $\int |a| \, d\gamma \leq 1$, this implies that

$$\left| \int a(y) \, dy \right| \leq Cr_B(1 + |c_B|)\gamma_0(c_B)^{-1}.$$

For the last term in (3.5), we thus get the bound $C|x - c_B|^{-n}r_B(1 + |c_B|)\gamma_0(c_B)^{-1}$.

Putting things together, we conclude that for $x \in B' \setminus 2B$,

$$\mathcal{M}_{\text{loc}}a(x) \leq C|x - c_B|^{-n-1}r_B\gamma_0(c_B)^{-1} + C|x - c_B|^{-n}r_B(1 + |c_B|)\gamma_0(c_B)^{-1}.$$

An integration with respect to $d\gamma$, or equivalently $\gamma_0 \, d\lambda$, then leads to

$$\int_{B' \setminus 2B} \mathcal{M}_{\text{loc}}a(x) \, d\gamma(x) \leq C + Cr_B(1 + |c_B|) \log \frac{\min(1, 1/|c_B|)}{r_B} \leq C,$$

and (3.4) is proved. □

Theorem 3.3. *Let $n = 1$, and suppose that f is a function in $L^1(\gamma)$. Then f is in $H^1(\gamma)$ if and only if $\mathcal{M}_{\text{loc}}f \in L^1(\gamma)$ and $E(f) < \infty$. The norms $\|f\|_{H^1(\gamma)}$ and $\|\mathcal{M}_{\text{loc}}f\|_{L^1(\gamma)} + E(f)$ are equivalent.*

Proof. Suppose that $f \in H^1(\gamma)$. Then $\mathcal{M}_{\text{loc}}f \in L^1(\gamma)$ by Lemma 3.2. To prove the necessity of the condition $E(f) < \infty$, it suffices to show that $E(a) < C$ for all Gaussian atoms a . This is obvious for the exceptional atom 1. If a is associated to a ball $B \in \mathcal{B}_1$, it follows from the inequality

$$\left| \int_{-\infty}^{-x} a \, d\gamma \right| + \left| \int_x^{\infty} a \, d\gamma \right| \leq \mathbf{1}_{(-B) \cup B}(x).$$

Conversely, assume that f is a function in $L^1(\gamma)$ such that $\mathcal{M}_{\text{loc}}f \in L^1(\gamma)$ and $E(f) < \infty$. We shall prove that $f \in H^1(\gamma)$, by constructing a Gaussian atomic decomposition $f = \sum_j \lambda_j a_j$ such that $\sum_j |\lambda_j| \leq C(\|\mathcal{M}_{\text{loc}}f\|_1 + E(f))$.

Most of the following argument, up to the decomposition (3.15), works also in the n -dimensional setting. Since we shall need it in the next section, we carry out that part in \mathbb{R}^n .

By subtracting a multiple of the exceptional atom 1, we may without loss of generality assume that

$$(3.7) \quad \int f \, d\gamma = 0.$$

Let $\{B_j\}$ be a covering of \mathbb{R}^n by maximal admissible balls. We can choose this covering in such way that the family $\{\frac{1}{2}B_j\}$ is disjoint and $\{4B_j\}$ has bounded overlap [10, Lemma 2.4]. Fix a smooth nonnegative partition of unity $\{\eta_j\}$ in \mathbb{R}^n such that $\text{supp } \eta_j \subset B_j$ and $\eta_j = 1$ on $\frac{1}{2}B_j$ and verifying $|\nabla \eta_j| \leq C/r_{B_j}$. Thus $f = \sum_j f \eta_j$. We now need the following lemma.

Lemma 3.4. *For g in $L^1_{\text{loc}}(\gamma)$ and $x \in \mathbb{R}^n$ one has*

$$(3.8) \quad \mathcal{M}_{\text{loc}}(g\eta_j\gamma_0)(x) \leq C \gamma_0(c_{B_j}) \mathcal{M}_{\text{loc}}g(x) \mathbf{1}_{4B_j}(x) \quad \forall j.$$

Proof. Since the support of η_j is contained in B_j , the support of $\mathcal{M}_{\text{loc}}(g\eta_j\gamma_0)$ is contained in the ball $4B_j$, because of Lemma 3.1. Moreover, for $\phi \in \Phi$ and $x \in 4B_j$,

$$\phi_t * (g\eta_j\gamma_0)(x) = \gamma_0(c_{B_j}) \tilde{\phi}_t * g(x),$$

where $\tilde{\phi}(z) = \phi(z)\eta_j(x - tz)\gamma_0(x - tz)/\gamma_0(c_{B_j})$. Thus, to prove (3.8) it suffices to show that there exists a positive constant C such that $\tilde{\phi} \in C\Phi$ for $x \in 4B_j$ and $0 < t < \min(1, 1/|x|)$. The support of $\tilde{\phi}$ is contained in $B(0, 1)$ and

$$|\tilde{\phi}(z)| \leq \frac{\gamma_0(x - tz)}{\gamma_0(c_{B_j})} \leq C,$$

because for $|z| \leq 1$,

$$|x - tz - c_{B_j}| \leq |x - c_{B_j}| + |tz| \leq C \min(1, 1/|c_{B_j}|).$$

Similarly $|\nabla \tilde{\phi}(z)| \leq C$, because the gradients $\nabla_z \eta_j(x - tz)$ and $\nabla_z \gamma_0(x - tz)/\gamma_0(c_{B_j})$ give the factors $t(1 + |c_{B_j}|)$ and $t|x - tz|\gamma_0(x - tz)/\gamma_0(c_{B_j})$, respectively, both of which are bounded. This concludes the proof of Lemma 3.4. \square

Continuing the proof of Theorem 3.3, we define $b_j \in \mathbb{C}$ for each $j \in \mathbb{N}$ by

$$(3.9) \quad \int_{-\infty}^{\infty} (f - b_j)\eta_j \, d\gamma = 0.$$

Note that since $\eta_j = 1$ on $\frac{1}{2}B_j$,

$$(3.10) \quad |b_j| = \left| \frac{\int f \eta_j \, d\gamma}{\int \eta_j \, d\gamma} \right| \leq C \frac{1}{\gamma(B_j)} \int_{B_j} |f| \, d\gamma.$$

We now apply Lemma 3.4 with $g = f - b_j$ and use the subadditivity of \mathcal{M}_{loc} combined with (3.10) to get

$$(3.11) \quad \begin{aligned} \int \mathcal{M}_{\text{loc}}((f - b_j)\eta_j\gamma_0) \, d\lambda &\leq C \int_{4B_j} \mathcal{M}_{\text{loc}}f \gamma_0(c_{B_j}) \, d\lambda + C \frac{\gamma(4B_j)}{\gamma(B_j)} \int_{B_j} |f| \, d\gamma \\ &\leq C \int_{4B_j} \mathcal{M}_{\text{loc}}f \, d\gamma. \end{aligned}$$

Lemma 3.5. *The function $(f - b_j)\eta_j\gamma_0$ is in $H^1(\lambda)$ and*

$$(3.12) \quad \|(f - b_j)\eta_j\gamma_0\|_{H^1(\lambda)} \leq \int_{4B_j} \mathcal{M}_{\text{loc}}f \, d\gamma.$$

Proof. By the maximal characterization of the classical space $H^1(\lambda)$, it suffices to show that

$$(3.13) \quad \int \mathcal{M}((f - b_j)\eta_j\gamma_0) \, d\lambda(x) \leq C \int_{4B_j} \mathcal{M}_{\text{loc}}f \, d\gamma.$$

Because of (3.11), all that needs to be verified is that

$$(3.14) \quad \int \sup_{\phi \in \Phi} \sup_{t \geq \min(1, 1/|x|)} |((f - b_j)\eta_j\gamma_0) * \phi_t(x)| \, d\lambda(x) \leq C \int_{4B_j} \mathcal{M}_{\text{loc}}f \, d\gamma.$$

To prove (3.14), we split the integral in the left-hand side into the sum

$$\int_{4B_j} \cdots \, d\lambda(x) + \int_{(4B_j)^c} \cdots \, d\lambda(x).$$

If $x \in 4B_j$, then for $\phi \in \Phi$ and $t \geq \min(1, 1/|x|)$,

$$\begin{aligned} |\phi_t * ((f - b_j)\eta_j\gamma_0)(x)| &\leq t^{-n} \int_{B_j} |f(y) - b_j| \, d\gamma(y) \\ &\leq (1 + |x|)^n \int_{B_j} (|f(y)| + |b_j|) \, d\gamma(y) \\ &\leq C(1 + |c_{B_j}|)^n \int_{B_j} |f(y)| \, d\gamma(y), \end{aligned}$$

the last inequality because of (3.10). Hence

$$\int_{4B_j} \cdots \, d\lambda(x) \leq C |4B_j| (1 + |c_{B_j}|)^n \int_{B_j} |f| \, d\gamma \leq C \int_{B_j} \mathcal{M}_{\text{loc}}f \, d\gamma.$$

If $x \in (4B_j)^c$, we take ϕ and t as before and observe that we can assume that $t > d(x, B_j)$, since otherwise the convolution in (3.14) will vanish. In view of (3.9) and (3.10), we then get

$$\begin{aligned} |\phi_t * ((f - b_j)\eta_j\gamma_0)(x)| &\leq \int_{B_j} |\phi_t(x - y) - \phi_t(x - c_{B_j})| |f(y) - b_j| \eta_j(y) \, d\gamma(y) \\ &\leq C t^{-n-1} \int_{B_j} |y - c_{B_j}| |f(y) - b_j| \, d\gamma(y) \\ &\leq C \frac{1}{d(x, B_j)^{n+1} r_{B_j}} \int_{B_j} |f(y)| \, d\gamma(y). \end{aligned}$$

This implies that

$$\int_{(4B_j)^c} \cdots \, d\lambda(x) \leq C \int_{B_j} |f| \, d\gamma \leq C \int_{B_j} \mathcal{M}_{\text{loc}}f \, d\gamma.$$

We have proved (3.14) and the lemma. □

We can now finish the proof of Theorem 3.3. By Lemmata 3.5 and 2.1, each function $(f - b_j)\eta_j\gamma_0$ has an atomic decomposition $\sum_k \lambda_{jk}\alpha_{jk}$, where the α_{jk} are Lebesgue atoms with supports in $2B_j$ and

$$\sum_k |\lambda_{jk}| \leq C \int_{4B_j} \mathcal{M}_{\text{loc}}f \, d\gamma.$$

As we saw in the proof of Theorem 2.2, each $a_{jk} = \gamma_0^{-1} \alpha_{jk}$ is a multiple of a Gaussian atom, with a factor which is independent of j and k . Thus

$$(3.15) \quad f = \sum_j (f - b_j) \eta_j + \sum_j b_j \eta_j = \sum_j \sum_k \lambda_{jk} a_{jk} + \sum_j b_j \eta_j$$

and

$$\sum_{j,k} |\lambda_{jk}| \leq \sum_j \int_{4B_j} \mathcal{M}_{\text{loc}} f \, d\gamma \leq C \|\mathcal{M}_{\text{loc}} f\|_{L^1(\gamma)}.$$

To complete the proof of Theorem 3.3, we need to find an atomic decomposition of $\sum_j b_j \eta_j$. It is here that we must restrict ourselves to the one-dimensional case and that the global condition $E(f) < \infty$ plays a role.

Choose the intervals $I_0 = (-1, 1)$, $I_j = (\sqrt{j-1}, \sqrt{j+1})$ for $j \geq 1$ and $I_j = -I_{|j|}$ for $j \leq -1$. The intervals I_j have essentially the same properties as the balls B_j introduced above, and we can use them instead of the B_j to construct η_j and b_j as before. To decompose now $\sum_j b_j \eta_j$, we first normalise the functions η_j , letting

$$\tilde{\eta}_j = \frac{\eta_j}{\int \eta_j \, d\gamma}.$$

Then $b_j \eta_j = \int f \eta_j \, d\gamma \tilde{\eta}_j$, and clearly

$$\sum_{j \geq k} \int f \eta_j \, d\gamma = \int f \mu_k \, d\gamma, \quad k \in \mathbb{Z},$$

where $\mu_k(x) = \sum_{j \geq k} \eta_j(x)$. Notice that $\int f \mu_k \, d\gamma \rightarrow 0$ as $k \rightarrow \pm\infty$, in view of (3.7). A summation by parts now yields

$$(3.16) \quad \sum_{j \in \mathbb{Z}} \int f \eta_j \, d\gamma \tilde{\eta}_j = \sum_{k \in \mathbb{Z}} \int f \mu_k \, d\gamma (\tilde{\eta}_k - \tilde{\eta}_{k-1}).$$

But $\tilde{\eta}_k - \tilde{\eta}_{k-1}$ is C times a Gaussian atom if we use admissible balls at some scale $s > 1$ in the definition of atoms. Thus (3.16) is our desired atomic decomposition of $\sum_j b_j \eta_j$, provided we can estimate the coefficients by showing that

$$(3.17) \quad \sum_{k \in \mathbb{Z}} \left| \int f \mu_k \, d\gamma \right| \leq C (\|f\|_1 + E(f)).$$

To this end, observe that

$$\int f \mu_k \, d\gamma = \int f(x) \int_{-\infty}^x \mu'_k(y) \, d\lambda(y) \, d\gamma(x) = \int \mu'_k(y) \int_y^\infty f(x) \, d\gamma(x) \, d\lambda(y).$$

Since the support of μ'_k is contained in I_k and

$$|\mu'_k(y)| \leq \frac{C}{|I_k|} \leq C (1 + |c_{I_k}|),$$

we obtain, using also the bounded overlap of the I_j ,

$$\begin{aligned} \sum_{k \in \mathbb{Z}} \left| \int f \mu_k \, d\gamma \right| &\leq C \sum_k \int_{I_k} (1 + |c_{I_k}|) \left| \int_y^\infty f(x) \, d\gamma(x) \right| \, d\lambda(y) \\ &\leq C \int_{-\infty}^\infty (1 + |y|) \left| \int_y^\infty f \, d\gamma \right| \, d\lambda(y) \\ &= C \int_0^\infty (1 + y) \left(\left| \int_y^\infty f \, d\gamma \right| + \left| \int_{-y}^\infty f \, d\gamma \right| \right) \, d\lambda(y) \\ &\leq C(\|f\|_1 + E(f)); \end{aligned}$$

here we used (3.7). This concludes the proof of Theorem 3.3. □

4. A CHARACTERIZATION OF NONNEGATIVE FUNCTIONS IN $H^1(\gamma)$

The dimension n is now arbitrary. The following lemma will be needed.

Lemma 4.1. *Let $\phi_0 = \gamma(B(0, 1))^{-1} \mathbf{1}_{B(0,1)}$. If $g \in L^\infty$ is supported in a maximal admissible ball B , then*

$$\left\| g - \int g \, d\gamma \phi_0 \right\|_{H^1(\gamma)} \leq C(1 + |c_B|^2) \gamma(B) \|g\|_{L^\infty}.$$

Proof. We shall construct atoms whose supports form a chain connecting $B(0, 1)$ to B . First we define a finite sequence of maximal admissible balls

$$\tilde{B}_0 = B(0, 1), \tilde{B}_1, \dots, \tilde{B}_N,$$

all with centres $c_{\tilde{B}_j}$ on the segment $[0, c_B]$. The absolute values $\rho_j = |c_{\tilde{B}_j}|$ shall be increasing in j , and the boundary $\partial\tilde{B}_j$ shall contain $c_{\tilde{B}_{j-1}}$ for $j = 1, \dots, N - 1$, which means that

$$(4.1) \quad \rho_j - \frac{1}{\rho_j} = \rho_{j-1}, \quad j = 1, \dots, N - 1,$$

and $\rho_0 = 0, \rho_1 = 1$. Finally, N is defined so that \tilde{B}_{N-1} is the first ball of the sequence that contains c_B , and $\tilde{B}_N = B$. Squaring (4.1), we get

$$\rho_j^2 - \rho_{j-1}^2 = 2 - \frac{1}{\rho_j^2} \geq 1,$$

so that $\rho_{N-1}^2 \geq N - 1$. It follows that

$$(4.2) \quad N \leq |c_B|^2 + 1.$$

Next, we denote by $B_j, j = 1, \dots, N$, the largest ball contained in $\tilde{B}_j \cap \tilde{B}_{j-1}$. Notice that the three balls $\tilde{B}_j, \tilde{B}_{j-1}$ and B_j have comparable radii and comparable Gaussian measures. Now define functions ϕ_j and g_j by setting

$$\begin{aligned} \phi_j &= \gamma(B_j)^{-1} \mathbf{1}_{B_j} \quad j = 1, \dots, N, \\ g_j &= \int g \, d\gamma (\phi_j - \phi_{j-1}), \quad j = 1, \dots, N, \\ g_{N+1} &= g - \int g \, d\gamma \phi_N. \end{aligned}$$

Clearly,

$$(4.3) \quad g - \int g \, d\gamma \, \phi_0 = \sum_{j=1}^{N+1} g_j.$$

Each function g_j is a multiple of an atom. Indeed, its integral against γ vanishes. Moreover, if $1 \leq j \leq N$, the support of g_j is contained in \tilde{B}_{j-1} and

$$\|g_j\|_\infty \leq (\gamma(B_j)^{-1} + \gamma(B_{j-1})^{-1}) \int |g| \, d\gamma \leq C \gamma(\tilde{B}_{j-1})^{-1} \gamma(B) \|g\|_{L^\infty}.$$

The support of ϕ_{N+1} is contained in B and

$$\|g_{N+1}\|_\infty \leq \|g\|_\infty + \gamma(B)^{-1} \int |g| \, d\gamma \leq C \|g\|_{L^\infty}.$$

Thus

$$\|g_j\|_{H^1(\gamma)} \leq C \gamma(B) \|g\|_{L^\infty}, \quad j = 1, \dots, N + 1.$$

Summing the coefficients in the atomic decomposition (4.3), we then obtain via (4.2),

$$\left\| g - \int g \, d\gamma \, \phi_0 \right\|_{H^1(\gamma)} \leq C(N + 1) \gamma(B) \|g\|_{L^\infty} \leq C(1 + |c_B|^2) \gamma(B) \|g\|_{L^\infty}.$$

The proof of the lemma is complete. □

Theorem 4.2. *Suppose that f is a function in $L^1(\gamma)$. If $\mathcal{M}_{\text{loc}}f$ is in $L^1(\gamma)$ and*

$$(4.4) \quad E_+(f) = \int |x|^2 |f(x)| \, d\gamma(x) < \infty,$$

then f is in $H^1(\gamma)$ and

$$\|f\|_{H^1(\gamma)} \leq C \|\mathcal{M}_{\text{loc}}f\|_1 + CE_+(f).$$

If f is nonnegative, the conditions $\mathcal{M}_{\text{loc}}f \in L^1(\gamma)$ and $E_+(f) < \infty$ are also necessary for f to be in $H^1(\gamma)$.

Proof. Let f be a function in $L^1(\gamma)$ such that $\mathcal{M}_{\text{loc}}f \in L^1(\gamma)$ and $E_+(f) < \infty$. Write $f = c(f) + f_0$, where $c(f) = \int f \, d\gamma$. Since $c(f)$ is a multiple of the exceptional atom, it suffices to find an atomic decomposition of f_0 . Note that f_0 satisfies

$$\mathcal{M}_{\text{loc}}f_0 \in L^1(\gamma) \quad \text{and} \quad \int |x|^2 |f_0(x)| \, d\gamma(x) < \infty.$$

Let $\{B_j\}$ be the covering of \mathbb{R}^n by maximal admissible balls and $\{\eta_j\}$ the corresponding partition of unity introduced in the proof of Theorem 3.3. As there, we choose numbers $b_j \in \mathbb{C}$ such that

$$\int_{-\infty}^{\infty} (f_0 - b_j) \eta_j \, d\gamma = 0 \quad \forall j.$$

Then the argument leading to (3.15) shows that

$$f_0 = \sum_j \sum_k \lambda_{jk} a_{jk} + \sum_j b_j \eta_j,$$

where the a_{jk} are Gaussian atoms supported in $4B_j$ and

$$\sum_{j,k} |\lambda_{jk}| \leq C \|\mathcal{M}_{\text{loc}}f_0\|_{L^1(\gamma)}.$$

It remains only to prove that $\sum_j b_j \eta_j$ is in $H^1(\gamma)$. We write $g_j = b_j \eta_j$ and observe that

$$(4.5) \quad \int \sum_j g_j \, d\gamma = 0$$

because f_0 and the a_{ij} have integrals zero. Thus

$$\sum_j g_j = \sum_j \left(g_j - \int g_j \, d\gamma \phi_0 \right),$$

where ϕ_0 is as in Lemma 4.1. Since (3.10) remains valid for f_0 , we have

$$(4.6) \quad \|g_j\|_\infty \leq C \frac{1}{\gamma(B_j)} \int_{B_j} |f_0| \, d\gamma.$$

Lemma 4.1 thus applies to each g_j , and using also the bounded overlap of the B_j we conclude

$$\left\| \sum_j g_j \right\|_{H^1(\gamma)} \leq C \sum_j (1 + |c_{B_j}|^2) \int_{B_j} |f_0| \, d\gamma \leq C \int (1 + |x|^2) |f_0| \, d\gamma.$$

This concludes the proof of the sufficiency and the norm estimate.

The necessity of the condition $\mathcal{M}_{\text{loc}} f \in L^1(\gamma)$ was obtained in Lemma 3.2.

To prove the necessity of (4.4), let $0 \leq f \in H^1(\gamma)$. We first observe that the function $x \mapsto |x|^2$ is in $BMO(\gamma)$. Indeed, its oscillation on any admissible ball is bounded. Since $BMO(\gamma)$ is a lattice, the functions $g_k(x) = \min\{|x|^2, k\}$ are in $BMO(\gamma)$, uniformly for $k \geq 1$. By the monotone convergence theorem and the duality between $H^1(\gamma)$ and $BMO(\gamma)$,

$$\int |x|^2 f(x) \, d\gamma(x) = \lim_k \int g_k(x) f(x) \, d\gamma(x) \leq C \|f\|_{H^1(\gamma)}.$$

The theorem is proved. □

The following result is a noteworthy consequence of Theorem 4.2.

Corollary 4.3. *For $1 < p \leq \infty$, one has continuous inclusions $L^p(\gamma) \subset H^1(\gamma)$ and $BMO(\gamma) \subset L^{p'}(\gamma)$, where $p' = p/(p - 1)$.*

Proof. We claim that the operator \mathcal{M}_{loc} is bounded on $L^p(\gamma)$ for $1 < p \leq \infty$. Deferring momentarily the proof of this claim, we complete the proof of the corollary. Suppose that f is in $L^p(\gamma)$. Then $\mathcal{M}_{\text{loc}} f$ is in $L^1(\gamma)$, because

$$\|\mathcal{M}_{\text{loc}} f\|_1 \leq \|\mathcal{M}_{\text{loc}} f\|_p \leq C \|f\|_p < \infty,$$

since $\gamma(\mathbb{R}^n) = 1$. Moreover, $E_+(f) \leq \| |x|^2 \|_{p'} \|f\|_p < \infty$, by Hölder's inequality. Thus $f \in H^1(\gamma)$ by Theorem 4.2. It also follows that the inclusion $L^p(\gamma) \subset H^1(\gamma)$ is continuous, and by duality we get the continuous inclusion $BMO(\gamma) \subset L^{p'}(\gamma)$.

It remains to prove the claim. We shall use again the covering $\{B_j\}$ from the proof of Theorem 3.3. First we observe that the inequality

$$(4.7) \quad \|\mathcal{M}_{\text{loc}} g\|_p \leq C \|g\|_p$$

holds when $\text{supp } g \subset B_j$, with a constant C independent of j . Indeed, \mathcal{M}_{loc} is bounded on $L^p(\lambda)$, and $\mathcal{M}_{\text{loc}} g$ is supported in the ball $4B_j$, where the Gaussian measure is essentially proportional to $d\lambda$.

Given a function $f \in L^p(\gamma)$, we write it as a sum $f = \sum f_j$ with $\text{supp } f_j \subset B_j$ and with the sets $\{f_j \neq 0\}$ pairwise disjoint. We can then apply (4.7) to each f_j and sum. \square

ACKNOWLEDGEMENT

The authors thank J. Dziubański for a useful discussion about Lemma 2.1.

REFERENCES

- [1] A. Carbonaro, G. Mauceri and S. Meda, H^1 and BMO on certain measured metric spaces, *Ann. Sc. Norm. Super. Pisa Cl. Sci.* (5) **8** (2009), no. 3, 543–582. MR2581426 (2010j:42025)
- [2] A. Carbonaro, G. Mauceri and S. Meda, H^1 and BMO for certain locally doubling metric measure spaces of finite measure, *Colloq. Math.* **118** (2010), 13–41. MR2600517 (2011b:42034)
- [3] A. Carbonaro, G. Mauceri and S. Meda, Comparison of spaces of Hardy type for the Ornstein-Uhlenbeck operator, *Potential Anal.* **33** (2010), 85–105. MR2644215 (2011k:42045)
- [4] D.-C. Chang, S. G. Krantz and E. M. Stein, H^p theory on a smooth domain in \mathbb{R}^n and elliptic boundary value problems, *J. Funct. Anal.* **114** (1993), 286–347. MR1223705 (94j:46032)
- [5] R. Coifman, Y. Meyer and E.M. Stein, Some new function spaces and their applications to harmonic analysis, *J. Funct. Anal.* **62** (1985), 304–335. MR791851 (86i:46029)
- [6] R. R. Coifman and G. Weiss, Extensions of Hardy spaces and their use in analysis, *Bull. Amer. Math. Soc.* **83** (1977), 569–645. MR0447954 (56:6264)
- [7] O. Dragicevic and A. Volberg, Bellman functions and dimensionless estimates of Littlewood-Paley type, *J. Oper. Theory* **56** (2006), 167–198. MR2261616 (2007g:42023)
- [8] E. B. Fabes, C. Gutiérrez and R. Scotto, Weak type estimates for the Riesz transforms associated with the Gaussian measure, *Rev. Mat. Iberoamericana* **10** (1994), 229–281. MR1286476 (95m:42023)
- [9] L. Forzani and R. Scotto, The higher order Riesz transforms for Gaussian measure need not be weak type $(1,1)$, *Studia Math.* **131** (1998), 205–214. MR1644460 (99h:42034)
- [10] J. García-Cuerva, G. Mauceri, P. Sjögren and J.L. Torrea, Higher order Riesz operators for the Ornstein-Uhlenbeck semigroup, *Potential Anal.* **10** (1999), 379–407. MR1698617 (2000j:42024)
- [11] J. García-Cuerva, G. Mauceri, P. Sjögren and J.L. Torrea, Spectral multipliers for the Ornstein-Uhlenbeck semigroup, *J. Anal. Math.* **78** (1999), 281–305. MR1714425 (2000m:42015)
- [12] J. García-Cuerva, G. Mauceri, S. Meda, P. Sjögren and J. L. Torrea, Functional calculus for the Ornstein-Uhlenbeck operator, *J. Funct. Anal.* **183** (2001), no. 2, 413–450. MR1844213 (2002d:47023)
- [13] J. García-Cuerva, G. Mauceri, S. Meda, P. Sjögren and J.L. Torrea, Maximal operators for the Ornstein-Uhlenbeck semigroup, *J. London Math. Soc.* **67** (2003), 219–234. MR1942422 (2003k:47048)
- [14] R.F. Gundy, Sur les transformations de Riesz pour le semigroupe d’Ornstein-Uhlenbeck, *C. R. Acad. Sci. Paris Sci. Ser. I Math.* **303** (1986), 967–970. MR877182 (88c:60108)
- [15] C. E. Gutiérrez, C. Segovia and J. L. Torrea, On higher order Riesz transforms for Gaussian measures, *J. Fourier Anal. Appl.* **2** (1996), 583–596. MR1423529 (98m:42017)
- [16] C. E. Gutiérrez and W. Urbina, Estimates for the maximal operator of the Ornstein-Uhlenbeck semigroup, *Proc. Amer. Math. Soc.* **113** (1991), no. 1, 99–104. MR1068123 (92a:42023)
- [17] C. E. Gutiérrez, On the Riesz transforms for Gaussian measures, *J. Funct. Anal.* **120** (1994), 107–134. MR1262249 (95c:35013)
- [18] W. Hebisch, G. Mauceri and S. Meda, Holomorphy of spectral multipliers of the Ornstein-Uhlenbeck operator, *J. Funct. Anal.* **210** (2004), 101–124. MR2052115 (2005g:47024)
- [19] L. Liu and D. Yang, Characterizations of BMO associated with Gauss measures via commutators of local fractional integrals, *Israel J. Math.* **180** (2010), 285–315. MR2735067 (2011k:42049)
- [20] L. Liu and D. Yang, BLO spaces associated with the Ornstein-Uhlenbeck operator, *Bull. Sci. Math.* **132** (2008), 633–649. MR2474485 (2010c:42045)

- [21] J. Maas, J. van Neerven and P. Portal, Conical square functions and non-tangential maximal functions with respect to the Gaussian measure, *Publ. Mat.* **55** (2011), no. 2, 313–341. MR2839445
- [22] J. Maas, J. van Neerven and P. Portal, Whitney coverings and the tent spaces $T^{1,q}(\gamma)$ for the Gaussian measure, [arXiv:1002.4911v1](https://arxiv.org/abs/1002.4911v1) [math.FA], to appear in *Ark. Mat.*
- [23] R. A. Macías and C. Segovia, A decomposition into atoms of distributions on spaces of homogeneous type, *Adv. in Math.* **33** (1979), 271–309. MR546296 (81c:32017b)
- [24] G. Mauceri and S. Meda, BMO and H^1 for the Ornstein–Uhlenbeck operator, *J. Funct. Anal.* **252** (2007), 278–313. MR2357358 (2008m:42024)
- [25] G. Mauceri, S. Meda and P. Sjögren, Sharp estimates for the Ornstein–Uhlenbeck operator, *Ann. Sc. Norm. Sup. Pisa, Classe di Scienze, Serie V*, **3** (2004), n. 3, 447–480. MR2099246 (2005j:47042)
- [26] G. Mauceri, S. Meda and P. Sjögren, Endpoint estimates for first-order Riesz transforms associated to the Ornstein-Uhlenbeck operator, *Rev. Mat. Iberoam.* **28** (2012), no. 1, 77–91. MR2904131
- [27] T. Menárguez, S. Pérez and F. Soria, The Mehler maximal function: a geometric proof of the weak type 1, *J. London Math. Soc. (2)* **61** (2000), 846–856. MR1766109 (2001i:42029)
- [28] P. A. Meyer, Transformations de Riesz pour le lois Gaussiennes, *Springer Lecture Notes in Mathematics* **1059** (1984), 179–193. MR770960 (86i:60150)
- [29] B. Muckenhoupt, Hermite conjugate expansions, *Trans. Amer. Math. Soc.* **139** (1969), 243–260. MR0249918 (40:3159)
- [30] S. Pérez, The local part and the strong type for operators related to the Gauss measure, *J. Geom. Anal.* **11** (2001), no. 3, 491–507. MR1857854 (2002h:42027)
- [31] S. Pérez and F. Soria, Operators associated with the Ornstein-Uhlenbeck semigroup, *J. London Math. Soc.* **61** (2000), 857–871. MR1766110 (2001i:42030)
- [32] G. Pisier, Riesz transforms: a simpler analytic proof of P. A. Meyer’s inequality, *Springer Lecture Notes in Mathematics* **1321** (1988), 485–501. MR960544 (89m:60178)
- [33] P. Sjögren, On the maximal function for the Mehler kernel, in *Harmonic Analysis, Cortona, 1982*, *Springer Lecture Notes in Mathematics* **992** (1983), 73–82. MR729346 (85j:35031)
- [34] E.M. Stein, *Harmonic Analysis. Real variable methods, orthogonality and oscillatory integrals*, Princeton Math. Series No. **43**, Princeton, N.J., 1993. MR1232192 (95c:42002)
- [35] W. Urbina, On singular integrals with respect to the Gaussian measure, *Ann. Sc. Norm. Sup. Pisa, Classe di Scienze, Serie IV*, **17** (1990), no. 4, 531–567. MR1093708 (92d:42010)

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI GENOVA, VIA DODECANESO 35, 16146 GENOVA, ITALIA

E-mail address: `mauceri@dima.unige.it`

DIPARTIMENTO DI MATEMATICA E APPLICAZIONI, UNIVERSITÀ DI MILANO-BICOCCA, VIA R. COZZI 53, 20125 MILANO, ITALY

E-mail address: `stefano.meda@unimib.it`

MATHEMATICAL SCIENCES, UNIVERSITY OF GOTHENBURG, BOX 100, S-405 30 GOTHENBURG, SWEDEN — AND — MATHEMATICAL SCIENCES, CHALMERS UNIVERSITY OF TECHNOLOGY, SE-412 96 GOTHENBURG, SWEDEN

E-mail address: `peters@chalmers.se`