

DESCENT OF RESTRICTED FLAT MITTAG–LEFFLER MODULES AND GENERALIZED VECTOR BUNDLES

SERGIO ESTRADA, PEDRO GUIL ASENSIO, AND JAN TRLIFAJ

(Communicated by Birge Huisgen-Zimmermann)

ABSTRACT. A basic question for any property of quasi-coherent sheaves on a scheme X is whether the property is local, that is, it can be defined using any open affine covering of X . Locality follows from the descent of the corresponding module property: for (infinite dimensional) vector bundles and Drinfeld vector bundles, it was previously proved by Kaplansky’s technique of dévissage. Since vector bundles coincide with \aleph_0 -restricted Drinfeld vector bundles, a question arose as to whether locality holds for κ -restricted Drinfeld vector bundles for each infinite cardinal κ . We give a positive answer here by replacing the dévissage with its recent refinement involving \mathcal{C} -filtrations and the Hill Lemma.

1. INTRODUCTION

In [4, §2] Drinfeld considered generalizations of the notion of a finite dimensional vector bundle on a scheme to the infinite dimensional setting. One of the options suggested replaced finitely generated projective modules by the flat Mittag–Leffler ones, leading thus to what was later called a Drinfeld vector bundle in [7]. Mittag–Leffler modules have been studied for decades, starting from the classic works of Grothendieck, Raynaud and Gruson [16] until recent contributions in [1], [10] et al.

Another ingredient of the theory is due to Quillen. Following [15], one can compute morphisms between two objects A and B of the derived category of the category $\mathcal{Q}\mathbf{co}(X)$ of quasi-coherent sheaves on a scheme X . First, one introduces a model category structure on $\mathbf{C}(\mathcal{Q}\mathbf{co}(X))$ (= the category of unbounded chain complexes on $\mathcal{Q}\mathbf{co}(X)$). Morphisms between A and B are then computed as the $\mathbf{C}(\mathcal{Q}\mathbf{co}(X))$ -morphisms between cofibrant and fibrant replacements of A and B modulo chain homotopy. Much later, Hovey and Gillespie [11], [8] developed a machinery for transferring the computation to $\mathbf{C}(\mathcal{Q}\mathbf{co}(X))$ using complete cotorsion pairs.

Received by the editors October 14, 2011 and, in revised form, September 24, 2012.

2010 *Mathematics Subject Classification*. Primary 14F05, 16D40; Secondary 03E35, 13D07, 18E15, 55N30.

Key words and phrases. Descent, restricted Drinfeld vector bundle, Mittag–Leffler module, \mathcal{C} -filtration, Whitehead module.

The first author was partially supported by DGI MTM2008-03339, by the Fundación Seneca and by the Junta de Andalucía, Consejería de Economía, Innovación y Ciencia and FEDER funds.

The third author was supported by GAČR 201/09/0816 and MSM 0021620839.

A test question for the Drinfeld options is to what extent they fit the Quillen–Hovey theory. While the answer for the general setting of arbitrary flat Mittag–Leffler modules is negative, there is a positive answer in case we admit only filtrations by Mittag–Leffler modules of restricted size [7]. This yields the notion of a κ –restricted Drinfeld vector bundle for each infinite cardinal $\kappa \geq \aleph_0$. In fact, the \aleph_0 –restricted Drinfeld vector bundles are exactly the (infinite dimensional) vector bundles from [4, §2, Definition], that is, those quasi–coherent sheaves all of whose modules of sections are projective, but not necessarily finitely generated.

While defining the notion of a restricted Drinfeld vector bundle, one faces the question of whether the notion is local, i.e., it can be defined using any open affine covering of the scheme. The question is known to have a positive answer in the two extreme cases: for all Drinfeld vector bundles and for all (infinite dimensional) vector bundles. In [16, Seconde partie] it was proved that locality follows from the descent of the corresponding module property, that of a flat Mittag–Leffler module and a projective module, respectively. Raynaud and Gruson proved the descent of projectivity using Kaplansky’s technique of dévissage and the fact that a module M is projective if and only if M is both flat Mittag–Leffler and a direct sum of countably generated modules (cf. [4, §2]). However, whether the notion of a κ –restricted Drinfeld vector bundle is also local for $\kappa > \aleph_0$ remained open [7]. Our main result here gives a positive answer:

Theorem 1.1. *Let κ be an infinite cardinal. Then the notion of a κ –restricted Drinfeld vector bundle is local. That is, if X is any scheme, V is a set of open affine sets of X such that $X = \bigcup_{v \in V} v$, and \mathcal{M} is a quasi–coherent sheaf on X such that $\mathcal{M}(v)$ is a κ –restricted flat Mittag–Leffler $\mathcal{R}(v)$ –module for all $v \in V$, then \mathcal{M} is a κ –restricted Drinfeld vector bundle on X .*

Our proof is also based on proving the descent, this time for restricted flat Mittag–Leffler modules, but it replaces Kaplansky’s dévissage by a more subtle technique for dealing with filtrations due to Hill. We also noticed that the κ –restricted flat Mittag–Leffler modules for $\kappa > \aleph_0$ are related to another interesting class of modules, the Whitehead ones. From a result of Eklof and Shelah [6] we infer that there exist local PIDs R_κ such that the κ –restricted Mittag–Leffler R_κ –modules coincide with the Whitehead R_κ –modules under the assumption of Gödel’s Axiom of Constructibility (see Example 2.3(ii)). So in this particular case the κ –restricted Drinfeld vector bundles over the affine scheme $\text{Spec}(R_\kappa)$ are exactly the Whitehead vector bundles.

2. PRELIMINARIES

Let R be a ring. Following [16], we call a (right R –) module M *Mittag–Leffler* provided that the canonical map $M \otimes_R \prod_{i \in I} M_i \rightarrow \prod_{i \in I} M \otimes_R M_i$ is monic for each family of left R –modules $(M_i \mid i \in I)$.

As in [4, §2], we will primarily be interested in flat Mittag–Leffler modules, that is, the Mittag–Leffler modules such that the functor $M \otimes_R -$ is exact. These modules were characterized in [16, Seconde partie, 2.2.2] as the modules M such that each finite subset of M is contained in a countably generated pure and projective submodule of M . In fact, they can also be characterized without referring to purity, as the \aleph_1 –projective modules [10]. For example, if R is a PID, then a module M is flat Mittag–Leffler iff each countably generated submodule of M is free.

The corresponding notion for quasi-coherent sheaves is as follows (see [7]): a quasi-coherent sheaf \mathcal{F} on a scheme X is a *Drinfeld vector bundle* provided that for each open affine set u , the $\mathcal{R}(u)$ -module of sections $\mathcal{M}(u)$ is flat and Mittag–Leffler.

We also recall [4, §2, Definition] that a quasi-coherent sheaf \mathcal{F} on a scheme X is an (infinite dimensional) *vector bundle* provided that for each open affine set u , the $\mathcal{R}(u)$ -module $\mathcal{M}(u)$ is projective.

A module M is a *Whitehead module* provided that $\text{Ext}_R^1(M, R) = 0$ [5]. A quasi-coherent sheaf \mathcal{F} on a scheme X is a *Whitehead vector bundle* provided that for each open affine set u , the $\mathcal{R}(u)$ -module $\mathcal{M}(u)$ is Whitehead.

Of course, each projective module is Whitehead and flat Mittag–Leffler. The relations between the classes of all Whitehead and flat Mittag–Leffler modules depend on the structure of the ring R . For example, if R is a Dedekind domain which is not R -complete, then all Whitehead modules are flat Mittag–Leffler because they are torsion-free (= flat) and each of their finite rank submodules is projective; cf. the proof of [5, XII.1.2]. But if R is an R -complete Dedekind domain (i.e., R is a complete DVR), then R is cotorsion [9, 4.4.4], so the Whitehead modules coincide with the flat ones and the inclusion of the classes is reversed.

In [7] it was shown that in general, the class of all Drinfeld vector bundles does not fit the Quillen–Hovey setting for computing cohomology via model category structures. Various restricted versions of the notion were suggested there to fix the problem. In order to define them, we recall the notion of a \mathcal{C} -filtration.

Let \mathcal{C} be a class of modules. A module M is \mathcal{C} -filtered provided there exist an ordinal σ and a chain of submodules of M ,

$$0 = M_0 \subseteq \cdots \subseteq M_\alpha \subseteq M_{\alpha+1} \subseteq \cdots \subseteq M_\sigma = M,$$

which is continuous (i.e., $M_\alpha = \bigcup_{\beta < \alpha} M_\beta$ for all limit ordinals $\alpha \leq \sigma$), and for each $\alpha < \sigma$, the module $M_{\alpha+1}/M_\alpha$ is isomorphic to an element of \mathcal{C} .

The chain $(M_\alpha \mid \alpha \leq \sigma)$ is called a \mathcal{C} -filtration of M . Notice that for $\sigma = 2$, M_2 is just the extension of the module $M_1 \in \mathcal{C}$ by the module $M_2/M_1 \in \mathcal{C}$. Also, the class of all \mathcal{C} -filtered modules includes arbitrary direct sums of copies of modules in \mathcal{C} .

A class of modules \mathcal{A} is *closed under transfinite extensions* provided that $M \in \mathcal{A}$ whenever M is \mathcal{A} -filtered. In this case \mathcal{A} is closed under extensions and arbitrary direct sums.

For example, if $\mathcal{A} = {}^\perp \mathcal{B}$ for a class \mathcal{B} , then \mathcal{A} is closed under transfinite extensions by the Eklof Lemma [9, 3.1.2]. Here, we define ${}^\perp \mathcal{B} = \text{Ker Ext}_R^1(-, \mathcal{B}) = \{M \in \text{Mod-}R \mid \text{Ext}_R^1(M, N) = 0 \text{ for all } N \in \mathcal{B}\}$. Similarly the class \mathcal{B}^\perp is defined; a pair of classes $(\mathcal{A}, \mathcal{B})$ is a *cotorsion pair* provided that $\mathcal{A} = {}^\perp \mathcal{B}$ and $\mathcal{B} = \mathcal{A}^\perp$.

In particular, the classes of all projective and flat modules are closed under transfinite extensions and direct summands because they are the first components of cotorsion pairs. The class of all flat Mittag–Leffler modules is also closed under transfinite extensions and direct summands, but it is not of the form ${}^\perp \mathcal{B}$ for any class of modules \mathcal{B} in general (see [1] and [10]).

Given a class \mathcal{A} of modules closed under transfinite extensions and an infinite cardinal κ , we let $\mathcal{A}^{\leq \kappa}$ denote the class of all $\leq \kappa$ -presented modules from \mathcal{A} , and $\mathcal{A}(\kappa)$ the class of all $\mathcal{A}^{\leq \kappa}$ -filtered modules. If $M \in \mathcal{A}(\kappa)$, then M is called a

κ -restricted module in \mathcal{A} . We have the chain

$$\mathcal{A}(\aleph_0) \subseteq \mathcal{A}(\aleph_1) \subseteq \cdots \subseteq \mathcal{A}(\kappa) \subseteq \mathcal{A}(\kappa^+) \subseteq \cdots \subseteq \mathcal{A} = \bigcup_{\aleph_0 \leq \kappa} \mathcal{A}(\kappa).$$

Example 2.1. 1. If \mathcal{A} is the class of all projective modules over any ring R , then the chain above is constant because $\mathcal{A}(\aleph_0) = \mathcal{A}$ by a classical result of Kaplansky [12].

2. If \mathcal{A} is the class of all flat modules over a ring R , then the chain stabilizes at $\text{card}R + \aleph_0$, that is, $\mathcal{A}(\kappa) = \mathcal{A}$ for each infinite cardinal $\kappa \geq \text{card}R$. This follows from Enochs’ solution of the Flat Cover Conjecture [3].

3. Let \mathcal{A} be the class of all flat Mittag–Leffler modules over any ring R . If κ is an infinite cardinal, then the modules in the class $\mathcal{A}(\kappa)$ are the κ -restricted flat Mittag–Leffler modules. A quasi-coherent sheaf \mathcal{F} on a scheme X is called a κ -restricted Drinfeld vector bundle provided that for each open affine set u , the $\mathcal{R}(u)$ -module of sections $\mathcal{M}(u)$ is κ -restricted flat Mittag–Leffler; see [7, p. 1423].

If R is right perfect, then \mathcal{A} is just the class of all projective modules, so we are in case 1. But if R is a non-right perfect ring, then the chain above is strictly increasing starting from $\text{card}R + \aleph_0$, because for each infinite cardinal $\kappa \geq \text{card}R$, there exists a module $M_{\kappa^+} \in \mathcal{A}^{\leq \kappa^+} \setminus \mathcal{A}(\kappa)$ (see [10, Theorem 6.10]).

Returning to the general case, we observe that the classes $\mathcal{A}(\kappa)$ are often the first components of complete cotorsion pairs. Here, a cotorsion pair $(\mathcal{A}, \mathcal{B})$ is said to be *complete* provided that for each module M there is a short exact sequence $0 \rightarrow M \rightarrow B \rightarrow A \rightarrow 0$ with $A \in \mathcal{A}$ and $B \in \mathcal{B}$.

Lemma 2.2. *Let R be a ring and κ be an infinite cardinal. Let \mathcal{A} be a class of modules closed under transfinite extensions and direct summands, and such that $R \in \mathcal{A}$. Then $(\mathcal{A}(\kappa), (\mathcal{A}^{\leq \kappa})^\perp)$ is a complete cotorsion pair.*

Proof. First, we observe that $\mathcal{A}(\kappa)^\perp = (\mathcal{A}^{\leq \kappa})^\perp$ by the Eklof Lemma [9, 3.1.2]. Let $\mathcal{E} = {}^\perp((\mathcal{A}^{\leq \kappa})^\perp)$. Since $\mathcal{A}^{\leq \kappa}$ has a representative set \mathcal{S} of objects and contains the regular module R , [9, 3.2.4] implies that \mathcal{E} consists of all direct summands of elements of $\mathcal{A}(\kappa)$. In particular, $\mathcal{E} \subseteq \mathcal{A}$ by assumption. Moreover, the Kaplansky theorem for cotorsion pairs [9, 4.2.11] shows that each element of \mathcal{E} is $\mathcal{E}^{\leq \kappa}$ -filtered, hence $\mathcal{A}^{\leq \kappa}$ -filtered, and $\mathcal{E} = \mathcal{A}(\kappa)$. Finally, since $(\mathcal{A}^{\leq \kappa})^\perp = \mathcal{S}^\perp$, the cotorsion pair $(\mathcal{A}(\kappa), (\mathcal{A}^{\leq \kappa})^\perp)$ is complete by [9, 3.2.1]. □

Moreover, if $K \in \mathcal{A}^{\leq \kappa}$ whenever $A \in \mathcal{A}^{\leq \kappa}$ and there is a short exact sequence $0 \rightarrow K \rightarrow P \rightarrow A \rightarrow 0$ with $P \leq \kappa$ -generated projective, then the cotorsion pair in Lemma 2.2 is hereditary; that is, $\text{Ext}_R^i(A, B) = 0$ for all $i \geq 2$, $A \in \mathcal{A}(\kappa)$, and $B \in (\mathcal{A}^{\leq \kappa})^\perp$. This is the case when:

- (i) \mathcal{A} = the class of all flat Mittag–Leffler modules over any ring by [1], and
- (ii) \mathcal{A} = the class of all flat modules over any right κ -noetherian ring (where a ring is κ -noetherian provided each right ideal of R is $\leq \kappa$ -generated; cf. [9, p. 138]).

Finally, let us consider the relations between restricted Mittag–Leffler modules and the better known classes of projective and Whitehead modules:

Example 2.3. (i) By the characterization of flat Mittag–Leffler modules in [16] mentioned above, all countably generated flat Mittag–Leffler modules are projective. So the notion of an \aleph_0 -restricted flat Mittag–Leffler module coincides with that of a projective module, and (infinite dimensional) vector bundles are exactly the \aleph_0 -restricted Drinfeld vector bundles.

(ii) Let κ be an infinite cardinal. By [6, Theorem 6] there exists a local, but not complete, PID $R = R_\kappa$ of cardinality 2^κ such that for each $\leq \kappa$ -generated module M , M is Whitehead iff M is flat Mittag–Leffler. By the Eklof Lemma [9, 3.1.2], it follows that each κ -restricted flat Mittag–Leffler module is Whitehead. If we assume $V = L$ (Gödel’s Axiom of Constructibility), then the converse holds as well: if M is Whitehead, then M is κ -restricted flat Mittag–Leffler (for M of cardinality $\lambda = \kappa^+ = 2^\kappa$, we use [5, XII.1.10]; the case of $\lambda > \kappa^+$ then follows by the proof of [9, 10.1.5] for \mathcal{A} = the class of all Whitehead modules). So in this case the κ -restricted Drinfeld vector bundles over the affine scheme $X = \text{Spec}(R_\kappa)$ are exactly the Whitehead vector bundles over X .

3. LOCALITY FOR INDUCED QUASI–COHERENT SHEAVES

Let \mathfrak{P} be a property of modules. For each commutative ring R , we let \mathfrak{P}_R denote the class of all R -modules satisfying \mathfrak{P} . Throughout, we will assume that \mathfrak{P} is *compatible with ring direct products* in the following sense: if $n < \omega$, $R = \prod_{i < n} R_i$ is a ring direct product, and $M_i \in \mathfrak{P}_{R_i}$ for each $i < n$, then $M = \prod_{i < n} M_i \in \mathfrak{P}_R$.

Definition 3.1. For a scheme X , we define the quasi-coherent sheaf *induced by* \mathfrak{P} on X (or a \mathfrak{P} -*quasi-coherent sheaf on* X) as the quasi-coherent sheaf \mathcal{M} such that for each open affine set u of X , the $\mathcal{R}(u)$ -module of sections $\mathcal{M}(u)$ satisfies \mathfrak{P} (that is, $\mathcal{M}(u) \in \mathfrak{P}_{\mathcal{R}(u)}$).

Example 3.2. Let \mathfrak{P} be the property of being a projective, flat, flat Mittag–Leffler, and κ -restricted flat Mittag–Leffler module (where κ is an infinite cardinal). Then \mathfrak{P}_R is the class of all projective, flat, flat Mittag–Leffler, and κ -restricted flat Mittag–Leffler R -modules (and \mathfrak{P} is obviously compatible with ring direct products). Moreover, \mathfrak{P} -quasi-coherent sheaves on a scheme X are exactly the (infinite dimensional) vector bundles, flat quasi-coherent sheaves, Drinfeld vector bundles, and κ -restricted Drinfeld vector bundles on X , respectively.

Definition 3.3. The notion of a \mathfrak{P} -quasi-coherent sheaf is *local* in case for each open affine covering $X = \bigcup_{v \in V} v$ of X and each quasi-coherent sheaf \mathcal{M} on X , \mathcal{M} is \mathfrak{P} -quasi-coherent provided that $\mathcal{M}(v) \in \mathfrak{P}_{\mathcal{R}(v)}$ for all $v \in V$.

So locality means that the property of being a \mathfrak{P} -quasi-coherent sheaf can be tested on any open affine covering of X .

We present a useful classic tool for proving locality based on ascent and descent of the corresponding module property (cf. [16, Seconde partie, §3.1] and [13, §23.18]):

Definition 3.4. Let \mathcal{R} be a class of commutative rings and \mathfrak{P} a property of modules compatible with ring direct products.

- (1) \mathfrak{P} is said to *ascend* in \mathcal{R} , provided that $M \otimes_R S \in \mathfrak{P}_S$ whenever $\varphi : R \rightarrow S$ is a flat homomorphism of rings in \mathcal{R} and $M \in \mathfrak{P}_R$.
- (2) \mathfrak{P} is said to *descend* in \mathcal{R} , provided that $M \in \mathfrak{P}_R$ whenever $\varphi : R \rightarrow S$ is a faithfully flat homomorphism of rings in \mathcal{R} and $M \otimes_R S \in \mathfrak{P}_S$.

If \mathfrak{P} ascends and descends in \mathcal{R} , then \mathfrak{P} is called the *ascent–descent property*, or the *AD-property*, in \mathcal{R} .

Lemma 3.5. *Let \mathcal{R} be a class of commutative rings and \mathfrak{P} be an AD-property in \mathcal{R} . Then the notion of a \mathfrak{P} -quasi-coherent sheaf is local.*

Proof. Let $X = \bigcup_{v \in V} v$ be an open affine covering of X such that $\mathcal{M}(v) \in \mathfrak{P}_{\mathcal{R}(v)}$ for all $v \in V$. Let u be an arbitrary open affine set of X . Then there exists a standard open covering $u = \bigcup_{j < n} u_j$ such that for each $j < n$ there exists $v_j \in V$ such that u_j is a standard open in v_j by [13, 21.11.6], so $u_j = D(f_j) = \text{Spec}(\mathcal{R}(v_j)_{f_j})$ for some $f_j \in \mathcal{R}(v_j)$. In particular, $\mathcal{M}(u_j) \cong \mathcal{M}(v_j) \otimes_{\mathcal{R}(v_j)} \mathcal{R}(v_j)_{f_j}$; hence $\mathcal{M}(u_j) \in \mathfrak{P}_{\mathcal{R}(u_j)}$ since \mathfrak{P} ascends in \mathcal{R} . The compatibility of \mathfrak{P} with ring direct products gives $\prod_{j < n} \mathcal{M}(u_j) \in \mathfrak{P}_{\prod_{j < n} \mathcal{R}(u_j)}$. Since the canonical morphism $\psi : \mathcal{R}(u) \rightarrow \prod_{j < n} \mathcal{R}(u_j)$ is faithfully flat, we conclude that $\mathcal{M}(u) \in \mathfrak{P}_{\mathcal{R}(u)}$ since \mathfrak{P} descends in \mathcal{R} . \square

While the ascent is usually easy to prove, proving the descent is more involved.

However, it is easy to see that being a flat module is an AD-property. Also, the property of being a flat Mittag-Leffler module is known to be an AD-property: the ascent is easy to prove, while the descent follows from the next lemma from [16, Seconde partie, §5.2]. Since there is an error in its original proof in [16] (cf. [14, Introduction]), we present a different proof here following [14, §9] (see also [13, 7.87.1]):

Lemma 3.6. *Let $\varphi : R \rightarrow S$ be a faithfully flat homomorphism of commutative rings, and M be a flat R -module such that $M \otimes_R S$ is a Mittag-Leffler S -module. Then M is a Mittag-Leffler R -module.*

Proof. First, we note that since M is flat, the property of being a Mittag-Leffler module can be restated in terms of the Mittag-Leffler condition for certain inverse systems: expressing M as a direct limit of a direct system $(F_i, f_{ij} \mid i \leq j \in I)$ of finitely presented free modules, $M = \varinjlim_{i \in I} F_i$, and applying the $\text{Hom}_R(-, R)$ functor, we obtain the inverse system $(\text{Hom}_R(F_i, R), \text{Hom}_R(f_{ij}, R) \mid i \leq j \in I)$. Then M is a Mittag-Leffler R -module iff this inverse system satisfies the Mittag-Leffler condition.

We have $M \otimes_R S = \varinjlim_{i \in I} F_i \otimes_R S$, and $M \otimes_R S$ is a Mittag-Leffler S -module by assumption. The Mittag-Leffler condition says that for each $i \in I$ the family $(\text{ImHom}_S(f_{ij} \otimes_R S, S) \mid i \leq j \in I)$ of S -submodules of $\text{Hom}_S(F_i \otimes_R S, S)$ stabilizes starting from some $j \geq i$. Since F_i is finitely presented and free, there is a natural isomorphism

$$\text{Hom}_S(F_i \otimes_R S, S) \cong \text{Hom}_R(F_i, R) \otimes_R S.$$

Since φ is faithfully flat, for each $i \in I$ the family $(\text{ImHom}_R(f_{ij}, R) \mid i \leq j \in I)$ of R -submodules of $\text{Hom}_R(F_i, R)$ stabilizes starting from some $j \geq i$. As mentioned above, the latter just says that M is a Mittag-Leffler R -module. \square

4. RESTRICTING AD-PROPERTIES OF MODULES

In order to treat the restricted Drinfeld vector bundles, we will need to transfer the unrestricted version of Lemma 3.5 to a restricted one. The following result (known as the Hill Lemma and proved e.g. in [9, Theorem 4.2.6]) will be our tool for refining the technique of dévissage from \mathcal{C} -direct sums of modules to the more general setting of \mathcal{C} -filtered modules:

Lemma 4.1. *Let R be a ring, λ a regular infinite cardinal, and \mathcal{C} a class of $< \lambda$ -presented modules. Let P be a module with a \mathcal{C} -filtration $\mathcal{P} = (P_\alpha \mid \alpha \leq \sigma)$. Then there is a family \mathcal{F} consisting of submodules of P such that*

- (H1) $\mathcal{P} \subseteq \mathcal{F}$,
- (H2) \mathcal{F} is closed under arbitrary sums and intersections,

- (H3) N'/N is \mathcal{P} -filtered for all $N, N' \in \mathcal{F}$ such that $N \subseteq N'$, and
 (H4) if $N \in \mathcal{F}$ and T is a subset of P of cardinality $< \lambda$, then there exists $N' \in \mathcal{F}$ such that $N \cup T \subseteq N'$ and N'/N is $< \lambda$ -presented.

For the restricted version, we will use the following.

Theorem 4.2. *Let \mathcal{R} be a class of commutative rings and \mathfrak{P} be an AD-property in \mathcal{R} . Assume moreover that for each $R \in \mathcal{R}$, the class \mathfrak{P}_R is closed under pure submodules and transfinite extensions, and contains R .*

Let κ be an infinite cardinal such that for each $R \in \mathcal{R}$, each $M \in \mathfrak{P}_R$, and each subset $A \subseteq M$ of cardinality $\leq \kappa$, there exists a $\leq \kappa$ -presented pure submodule $N \subseteq M$ containing A .

Let \mathfrak{R} be the restricted property of modules defined by $\mathfrak{R}_R = \mathfrak{P}_R(\kappa)$ for each $R \in \mathcal{R}$. Then \mathfrak{R} is an AD-property in \mathcal{R} .

Proof. Clearly, \mathfrak{R} is compatible with ring direct products because \mathfrak{P} is such.

Next, we prove that \mathfrak{R} ascends in \mathcal{R} . If $M \in \mathfrak{P}_R(\kappa)$, then M is $(\mathfrak{P}_R)^{\leq \kappa}$ -filtered. Since S is a flat R -module, $M \otimes_R S$ is \mathcal{C} -filtered, where \mathcal{C} is the class of all modules of the form $N \otimes_R S$ where $N \in (\mathfrak{P}_R)^{\leq \kappa}$. Since \mathfrak{P} ascends in \mathcal{R} , we have $N \otimes_R S \in \mathfrak{P}_S$. Since N is $\leq \kappa$ -presented as an R -module, so is $N \otimes_R S$ as an S -module. Thus $M \otimes_R S$ is $(\mathfrak{P}_S)^{\leq \kappa}$ -filtered and \mathfrak{R} ascends in \mathcal{R} .

In order to prove that \mathfrak{R} descends in \mathcal{R} , let M be an R -module such that $P = M \otimes_R S \in \mathfrak{P}_S(\kappa)$, so there exists a $(\mathfrak{P}_S)^{\leq \kappa}$ -filtration $\mathcal{P} = (P_\alpha \mid \alpha \leq \sigma)$ of P . Since \mathfrak{P} descends in \mathcal{R} , we have $M \in \mathfrak{P}_R$.

Consider the family \mathcal{F} corresponding to the filtration \mathcal{P} by Lemma 4.1 (for the regular infinite cardinal $\lambda = \kappa^+$). Notice that Condition (H3) of Lemma 4.1 yields $\mathcal{F} \subseteq \mathfrak{P}_S(\kappa)$ and $P/N \in \mathfrak{P}_S(\kappa)$ for each $N \in \mathcal{F}$.

We will use the family \mathcal{F} to construct a $(\mathfrak{P}_R)^{\leq \kappa}$ -filtration $(M_\beta \mid \beta \leq \tau)$ of M by induction on β as follows:

We let $M_0 = 0$ and assume that M_β is constructed so that M_β is a pure submodule of M such that $N = M_\beta \otimes_R S \in \mathcal{F}$ and $M/M_\beta \in \mathfrak{P}_R$.

Assume there exists $x \in M \setminus M_\beta$. Since $M/M_\beta \in \mathfrak{P}_R$, the assumption on \mathfrak{P}_R yields a $\leq \kappa$ -presented submodule $U_1 = Q_1/M_\beta$ of M/M_β containing $x + M_\beta$ such that U_1 is pure in M/M_β (and hence Q_1 is pure in M by [9, Lemma 1.2.17(b)]). Then $N = N_0 \subseteq Q_1 \otimes_R S$.

Condition (H4) of Lemma 4.1 provides us with $N_1 \in \mathcal{F}$ such that $Q_1 \otimes_R S \subseteq N_1$ and N_1/N_0 is $\leq \kappa$ -presented. Again, the assumption on \mathfrak{P}_R gives a $\leq \kappa$ -presented submodule $U_1 \subseteq U_2 = Q_2/M_\beta$ of M/M_β such that U_2 is pure in M/M_β (and hence Q_2 is pure in M) and $N_1 \subseteq Q_2 \otimes_R S$. Proceeding similarly, we obtain a sequence of pure R -submodules $M_\beta = Q_0 \subseteq Q_1 \subseteq Q_2 \subseteq \dots$ of M and a sequence $N = N_0 \subseteq N_1 \subseteq N_2 \subseteq \dots$ of elements of \mathcal{F} .

Let $M_{\beta+1} = \bigcup_{i < \omega} Q_i$ and $N' = \bigcup_{i < \omega} N_i$. Then $M_{\beta+1}$ is a pure submodule of M , and $N' \in \mathcal{F}$ by condition (H2) of Lemma 4.1. Moreover, $M_{\beta+1}/M_\beta = \bigcup_{i < \omega} U_i$ is a $\leq \kappa$ -presented pure submodule of M/M_β , hence $M_{\beta+1}/M_\beta \in (\mathfrak{P}_R)^{\leq \kappa}$ by the assumption on \mathfrak{P}_R . Moreover, $x \in M_{\beta+1}$ and $M_{\beta+1} \otimes_R S = N'$ by construction.

Since $(M/M_{\beta+1}) \otimes_R S \cong P/N' \in \mathfrak{P}_S(\kappa)$, the descent of \mathfrak{P} in \mathcal{R} gives $M/M_{\beta+1} \in \mathfrak{P}_R$.

If β is a limit ordinal, we let $M_\beta = \bigcup_{\gamma < \beta} M_\gamma$. Then M_β is pure in P and $M_\beta \otimes_R S = \bigcup_{\gamma < \beta} (M_\gamma \otimes_R S) \in \mathcal{F}$ by condition (H2) of Lemma 4.1. Again, $(M/M_\beta) \otimes_R S \cong P/(M_\beta \otimes_R S) \in \mathcal{F}$, so the descent of \mathfrak{P} implies that $M/M_\beta \in \mathfrak{P}_R$.

By construction, there exists an ordinal τ such that $M_\tau = M$; hence $(M_\beta \mid \beta \leq \tau)$ is the desired $(\mathfrak{P}_R)^{\leq \kappa}$ -filtration of M . \square

To complete the picture, we recall several known properties of flat Mittag-Leffler modules:

Lemma 4.3. *Let R be a ring. Then the class of all flat Mittag-Leffler modules is closed under pure submodules.*

Let κ be an infinite cardinal. Then each $\leq \kappa$ -generated flat Mittag-Leffler module is $\leq \kappa$ -presented. Moreover, each subset of cardinality $\leq \kappa$ in a flat Mittag-Leffler module M is contained in a pure and $\leq \kappa$ -presented submodule N of M .

Proof. The first claim is [16, Seconde partie, 2.1.6]. For the second, see [2, Proposition 3.8] and [7, Lemma 2.7(ii)]. \square

Now, we can prove our main result:

Proof of Theorem 1.1. Let \mathfrak{P} denote the property of being a flat Mittag-Leffler module. By Lemmas 3.6 and 4.3, \mathfrak{P} satisfies the assumptions of Theorem 4.2 for $\mathcal{R} =$ the class of all commutative rings. Thus the property \mathfrak{R} of being a κ -restricted flat Mittag-Leffler module is an AD-property, and the claim follows by Lemma 3.5. \square

REFERENCES

- [1] Lidia Angeleri Hügel and Dolores Herbera, *Mittag-Leffler conditions on modules*, Indiana Univ. Math. J. **57** (2008), no. 5, 2459–2517, DOI 10.1512/iumj.2008.57.3325. MR2463975 (2009i:16002)
- [2] Silvana Bazzoni and Dolores Herbera, *Cotorsion pairs generated by modules of bounded projective dimension*, Israel J. Math. **174** (2009), 119–160, DOI 10.1007/s11856-009-0106-x. MR2581211 (2011a:16018)
- [3] L. Bican, R. El Bashir, and E. Enochs, *All modules have flat covers*, Bull. London Math. Soc. **33** (2001), no. 4, 385–390, DOI 10.1017/S0024609301008104. MR1832549 (2002e:16002)
- [4] Vladimir Drinfeld, *Infinite-dimensional vector bundles in algebraic geometry: an introduction*, The unity of mathematics, Progr. Math., vol. 244, Birkhäuser Boston, Boston, MA, 2006, pp. 263–304, DOI 10.1007/0-8176-4467-9.7. MR2181808 (2007d:14038)
- [5] Paul C. Eklof and Alan H. Mekler, *Almost free modules*, Set-theoretic methods, Revised edition, North-Holland Mathematical Library, vol. 65, North-Holland Publishing Co., Amsterdam, 2002. MR1914985 (2003e:20002)
- [6] Paul C. Eklof and Saharon Shelah, *Whitehead modules over large principal ideal domains*, Forum Math. **14** (2002), no. 3, 477–482, DOI 10.1515/form.2002.021. MR1899295 (2003f:13014)
- [7] Sergio Estrada, Pedro A. Guil Asensio, Mike Prest, and Jan Trlifaj, *Model category structures arising from Drinfeld vector bundles*, Adv. Math. **231** (2012), no. 3-4, 1417–1438, DOI 10.1016/j.aim.2012.06.011. MR2964610
- [8] James Gillespie, *Kaplansky classes and derived categories*, Math. Z. **257** (2007), no. 4, 811–843, DOI 10.1007/s00209-007-0148-x. MR2342555 (2009e:55030)
- [9] Rüdiger Göbel and Jan Trlifaj, *Approximations and endomorphism algebras of modules*, de Gruyter Expositions in Mathematics, vol. 41, Walter de Gruyter GmbH & Co. KG, Berlin, 2006. MR2251271 (2007m:16007)
- [10] Dolores Herbera and Jan Trlifaj, *Almost free modules and Mittag-Leffler conditions*, Adv. Math. **229** (2012), no. 6, 3436–3467, DOI 10.1016/j.aim.2012.02.013. MR2900444
- [11] Mark Hovey, *Cotorsion pairs, model category structures, and representation theory*, Math. Z. **241** (2002), no. 3, 553–592, DOI 10.1007/s00209-002-0431-9. MR1938704 (2003m:55027)
- [12] Irving Kaplansky, *Projective modules*, Ann. of Math (2) **68** (1958), 372–377. MR0100017 (20 #6453)
- [13] J. de Jong et al., *The Stacks Project*, Version 118f702, <http://math.columbia.edu/algebraic-geometry/stacks-git/book.pdf>.

- [14] A. Perry, *Faithfully flat descent for projectivity of modules*, arXiv:1011.0038v1.
- [15] Daniel G. Quillen, *Homotopical algebra*, Lecture Notes in Mathematics, No. 43, Springer-Verlag, Berlin, 1967. MR0223432 (36 #6480)
- [16] Michel Raynaud and Laurent Gruson, *Critères de platitude et de projectivité. Techniques de “platification” d’un module* (French), Invent. Math. **13** (1971), 1–89. MR0308104 (46 #7219)

DEPARTAMENTO DE MATEMÁTICA APLICADA, UNIVERSIDAD DE MURCIA, CAMPUS DEL ESPINARDO 30100, MURCIA, SPAIN

E-mail address: `sestrada@um.es`

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE MURCIA, CAMPUS DEL ESPINARDO 30100, MURCIA, SPAIN

E-mail address: `paguil@um.es`

FACULTY OF MATHEMATICS AND PHYSICS, DEPARTMENT OF ALGEBRA, CHARLES UNIVERSITY, SOKOLOVSKÁ 83, 186 75 PRAGUE 8, CZECH REPUBLIC

E-mail address: `trlifaj@karlin.mff.cuni.cz`