

## INTERPRETING WEAK KÖNIG'S LEMMA USING THE ARITHMETIZED COMPLETENESS THEOREM

TIN LOK WONG

(Communicated by Mirna Džamonja)

ABSTRACT. We present a previously unpublished proof of the conservativity of  $\text{WKL}_0$  over  $\text{IS}_1$  using the Arithmetized Completeness Theorem, which, in particular, constitutes an  $\omega$ -interpretation of  $\text{WKL}_0$  in  $\text{IS}_1$ . We also show that  $\text{WKL}_0^*$  is interpretable in  $\text{ID}_0 + \text{exp}$ .

$\text{IS}_1$  and  $\text{WKL}_0$  are prominent theories in the study of arithmetic and the foundations of mathematics [3, 8]. The intimate connections between these theories can best be seen from a fundamental and influential theorem of Leo Harrington's which states that every countable model of  $\text{IS}_1$  expands to a model of  $\text{WKL}_0$ . This theorem, in particular, implies  $\text{WKL}_0$  is conservative over  $\text{IS}_1$ .

In this short note, we demonstrate how (several strengthenings due independently to Hájek [2] and Avigad [1] of) Harrington's theorem can be established simply by putting together a number of standard facts in the literature. While the usual proof of Harrington's theorem is based on tree forcing [8, Section IX.2], our approach uses instead a version of the Arithmetized Completeness Theorem, which can be proved by a forcing argument in which the conditions are binary strings of bounded lengths.

The notation in this paper is more or less standard. Unless otherwise stated, 'definable' means 'parametrically definable'. We assume familiarity with Hájek–Pudlák [3] and some acquaintance with Simpson [8]. Nevertheless, let us first briefly review some basic definitions and facts.

The language for first-order arithmetic  $\mathcal{L}_1$  has symbols  $0, 1, +, \times, \leq$ . A *bounded* quantifier is one of the form  $Qx \leq t$ , where  $Q \in \{\forall, \exists\}$  and  $t$  is a term not involving  $x$ . An  $\mathcal{L}_1$  formula is  $\Delta_0$  if all the quantifiers it contains are bounded. Formulas of the form  $\exists \bar{v} \varphi(\bar{v}, \bar{x})$  where  $\varphi \in \Delta_0$  are called  $\Sigma_1$ . A formula  $\theta$  is  $\Delta_1$  if, modulo logical equivalence, both  $\theta$  and  $\neg\theta$  are  $\Sigma_1$ . The closure of  $\Sigma_1$  under Boolean operations and bounded quantification is denoted  $\Delta_0(\Sigma_1)$ . Axiomatize  $\text{IS}_1$  by the theory of the non-negative parts of discretely ordered rings (commonly referred to as  $\text{PA}^-$ ) and the induction scheme for  $\Sigma_1$  formulas. Define  $\text{ID}_0$  similarly. The theory  $\text{BS}_1$  consists of  $\text{ID}_0$  and the  $\Sigma_1$  *collection scheme*, which asserts that

$$\forall \bar{z} \forall a (\forall x \leq a \exists y \varphi(x, y, \bar{z}) \rightarrow \exists b \forall x \leq a \exists y \leq b \varphi(x, y, \bar{z})),$$

---

Received by the editors July 10, 2015 and, in revised form, October 2, 2015 and November 6, 2015.

2010 *Mathematics Subject Classification*. Primary 03C62, 03F25, 03H15.

Part of this paper was presented at the Logic Colloquium in Vienna, Austria, in July 2014. The author was financially supported by the Austrian Science Fund (FWF) project P24654-N25 while this research was carried out.

whenever  $\varphi \in \Sigma_1$ . We denote by  $\text{exp}$  a fixed  $\mathcal{L}_I$  sentence expressing the totality of  $x \mapsto 2^x$  over  $\text{ID}_0$ . It is commonly known [3, Sections I.1(c), IV.1(f), and V.1(a)] that  $\text{IS}_1 \vdash \text{BS}_1 + \text{exp}$  and  $\text{BS}_1 \not\vdash \text{exp}$ . Within  $\text{ID}_0 + \text{exp}$ , the usual operations on sequences and syntactical objects are well behaved [3, Section V.3].

The language for second-order arithmetic  $\mathcal{L}_{II}$  has a first-order sort for numbers and a second-order sort for sets of numbers. We write  $\mathcal{L}_{II}$  structures as pairs  $(M, \mathcal{X}^c)$ , where  $M$  is the universe for the first-order sort, and  $\mathcal{X}^c$  is the universe for the second-order sort. The language  $\mathcal{L}_{II}$  has a copy of  $\mathcal{L}_I$  on the first-order sort, and a membership relation linking the two sorts. All  $\mathcal{L}_{II}$  structures are required to satisfy the axiom of extensionality. The definitions of  $\Sigma_0^0, \Sigma_1^0, \Delta_1^0, \text{IS}_0^0, \text{IS}_1^0, \dots$  are analogous to those of  $\Delta_0, \Sigma_1, \Delta_1, \text{ID}_0, \text{IS}_1, \dots$ , except that we now allow the appearance of second-order variables. *Weak König's Lemma* asserts that every unbounded 0–1 tree has an unbounded branch. The theory  $\text{WKL}_0^*$  consists of  $\text{IS}_0^0 + \text{exp}$ , the  $\Delta_1^0$  comprehension scheme, and Weak König's Lemma. Define  $\text{WKL}_0 = \text{WKL}_0^* + \text{IS}_1^0$ . The following theorem shows that  $\text{WKL}_0$  is *conservative* over  $\text{IS}_1$ , i.e., every  $\mathcal{L}_I$  sentence provable in  $\text{WKL}_0$  is already provable in  $\text{IS}_1$ .

**Theorem 1** (Hájek, Avigad). *Every  $M \models \text{IS}_1$  expands to  $(M, \mathcal{X}^c) \models \text{WKL}_0$ .*

*Proof.* First, it is well known [3, Corollary I.4.34(2)] that  $\text{IS}_1$  proves the consistency of  $\text{ID}_0 + \text{exp}$ . Second, the Low Arithmetized Completeness Theorem in Hájek–Pudlák [3, Theorem I.4.27] tells us that, provably in  $\text{IS}_1$ , every  $\Delta_1$ -definable consistent theory has a definable model all of whose  $\Sigma_1^0$  properties are  $\Delta_0(\Sigma_1)$ -definable. These two facts together yield  $K \models \text{ID}_0 + \text{exp}$  properly end extending  $M$  such that whenever  $\bar{c} \in K$  and  $\theta$  is an  $\mathcal{L}_I$  formula,

$$(*) \quad \{\bar{x} \in M : K \models \theta(\bar{v}, \bar{x}, \bar{c}) \text{ for some } \bar{v} \in M\} \text{ is } \Delta_0(\Sigma_1)\text{-definable in } M.$$

As is usual in Arithmetized Completeness Theorem constructions (for example, that in Paris [6, p. 254]), the embedding  $M \rightarrow K$  hinges on the observation that every  $a \in M$  is represented by a term  $\bar{a}$  in the  $\mathcal{L}_I$  in  $M$ . The image of this embedding is closed downwards because  $\text{ID}_0 + \text{exp} \vdash \forall x \leq \bar{a} \bigvee_{i \leq a} x = i$  for all  $a$ , provably in  $\text{IS}_1$ .

Let us say that an element  $c \in K$  codes a subset  $A \subseteq M$  if

$$A = \{i \in M : \text{the } i\text{th digit in the binary expansion of } c \text{ is } 1\}.$$

Denote the collection of all subsets of  $M$  coded in  $K$  by  $\text{Cod}(K/M)$ . A standard overspill argument [9, Theorem 4.8] shows  $(M, \text{Cod}(K/M)) \models \text{WKL}_0^*$ . For the sake of completeness, we include a sketch of this argument here. Consider  $\Delta_1^0$  comprehension. If  $A \subseteq M$  is  $\Delta_1^0$ -definable in  $(M, \text{Cod}(K/M))$ , then its  $\Delta_1^0$ -ness overflows into  $K \setminus M$ , producing a bound which enables  $K$  to see  $A$  in a  $\Delta_0$  way. Bounded  $\Delta_0$  comprehension, available in  $K$  from  $\text{ID}_0 + \text{exp}$ , then gives what we want. Next, if  $B$  is an unbounded 0–1 tree in  $(M, \text{Cod}(K/M))$ , then its code contains a node whose length is in  $K \setminus M$  by overspill, and the predecessors of such a node trace an unbounded branch in  $B$ . This shows Weak König's Lemma in  $(M, \text{Cod}(K/M))$ .

It remains to prove  $(M, \text{Cod}(K/M)) \models \text{IS}_1^0$ . Notice for each  $\varphi \in \Sigma_0^0$ , we can find  $\psi \in \Delta_0$ , simply by replacing set parameters by their codes, such that if  $\bar{A} \in \text{Cod}(K/M)$  coded by  $\bar{c} \in K$ , then for all  $\bar{x} \in M$ ,

$$(M, \text{Cod}(K/M)) \models \varphi(\bar{x}, \bar{A}) \iff K \models \psi(\bar{x}, \bar{c}).$$

Hence, every  $\Sigma_1^0$  property of  $(M, \text{Cod}(K/M))$  translates to a  $\Delta_0(\Sigma_1)$  property of  $M$  by  $(*)$ . Since  $\text{IS}_1$  proves the induction scheme for  $\Delta_0(\Sigma_1)$  formulas [3, Lemma I.2.14], this implies  $(M, \text{Cod}(K/M)) \models \text{IS}_1^0$ , as required.  $\square$

A similar proof was independently discovered by F. Félix Lara-Martín [personal communication]. It solves the second part of Problem 1 in Paris [6]. Analogous arguments for higher levels of the arithmetic hierarchy can be found in Paris’s paper. The original proofs by Hájek and Avigad both go via an  $\omega$ -interpretation of  $\text{WKL}_0$  in  $\text{I}\Sigma_1$ . The reader may consult Section III.1(a) of Hájek–Pudlák [3] for background information about interpretations.

**Definition.** An interpretation of an  $\mathcal{L}_{\text{II}}$  theory in an  $\mathcal{L}_1$  theory is an  $\omega$ -interpretation if its restriction to the first-order sort is the identity interpretation.

Essentially, an  $\omega$ -interpretation of an  $\mathcal{L}_{\text{II}}$  theory  $T$  in an  $\mathcal{L}_1$  theory  $T_0$  is a uniform recipe for expanding every  $M \models T_0$  to  $(M, \mathcal{X}) \models T$ . Hájek’s  $\omega$ -interpretation [2, Section 3] employs a notion of ‘very low’ sets with respect to which Weak König’s Lemma is true. Avigad’s  $\omega$ -interpretation [1] involves a formalization of the usual forcing proof. While the former of these is slightly cleaner, the latter has the advantage of being applicable also to general models  $(M, \mathcal{X}) \models \text{I}\Sigma_1^0$ , not only the ones in which  $\mathcal{X}$  is finite. Upon closer inspection, one sees that our proof is uniform enough to give rise to an  $\omega$ -interpretation too. Notice our  $\omega$ -interpretation, unlike theirs, is iteration-free.

**Theorem 2** (Hájek, Avigad). *There is an  $\omega$ -interpretation of  $\text{WKL}_0$  in  $\text{I}\Sigma_1$ .*

*Proof.* By looking into the proof of the Low Arithmetized Completeness Theorem, the reader can verify that the definition (of the elementary diagram) of  $K$  in our proof of Theorem 1 does not depend on  $M$ . Since every element of  $K$  codes a set in  $\text{Cod}(K/M)$ , we can use the definition of  $K$  as the interpretation of the second-order sort. The membership relation is interpreted accordingly. (We do not have a choice on how the other symbols are interpreted.)  $\square$

As shown by Simpson and Smith [9, Theorem 4.6], an analogue of Theorem 1 holds at the  $\text{B}\Sigma_1 + \text{exp}$  level: every countable model  $M \models \text{B}\Sigma_1 + \text{exp}$  expands to  $(M, \mathcal{X}) \models \text{WKL}_0^*$ . It is not known whether the countability condition can be omitted here. In particular, it is not known whether there is an  $\omega$ -interpretation of  $\text{WKL}_0^*$  in  $\text{B}\Sigma_1 + \text{exp}$ . What we can prove here is that  $\text{WKL}_0^*$  is interpretable in  $\text{B}\Sigma_1 + \text{exp}$ . Since  $\text{B}\Sigma_1 + \text{exp}$  is interpretable in  $\text{I}\Delta_0 + \text{exp}$  [2, Theorem 2.4], this is equivalent to interpretability in  $\text{I}\Delta_0 + \text{exp}$ . This answers Question (5) in Hájek [2].

**Corollary 3.** *There is an interpretation of  $\text{WKL}_0^*$  in  $\text{I}\Delta_0 + \text{exp}$ .*

*Proof.* We only describe how to make a model of  $\text{WKL}_0^*$  from a model of  $\text{I}\Delta_0 + \text{exp}$ . Modulo the finite axiomatizability of  $\text{I}\Sigma_1$  [3, Theorem I.2.52], it is straightforward to check that this description gives rise to an interpretation.

Take any  $M \models \text{I}\Delta_0 + \text{exp}$ . If  $M \models \text{I}\Sigma_1$ , then apply Theorem 2. Suppose  $M \not\models \text{I}\Sigma_1$ . Follow Kołodziejczyk–Yokoyama [5, Lemma 9] to find a parameter-free  $\Sigma_1$ -definable proper cut  $I$  of  $M$  that is closed under  $x \mapsto 2^x$ . Then  $(I, \text{Cod}(M/I)) \models \text{WKL}_0^*$  as in our proof of Theorem 1. This is the model we want.  $\square$

Having known the conservativity of a stronger theory  $T$  over a weaker theory  $T_0$ , the natural question is then whether  $T$  helps make proofs of theorems of  $T_0$  significantly shorter. Ignjatovic [4, Chapter 3] observed that  $\omega$ -interpretations can be utilized to obtain negative answers to such questions.

**Definition.** Let  $T_0$  be (an axiomatization of) a theory, and let  $T$  be an extension of  $T_0$ , possibly in a bigger language. We say  $T$  has *at most polynomial speed-up* over  $T_0$  if there exists  $n \in \mathbb{N}$  such that for every proof from  $T$  with  $\ell$  symbols of a theorem of  $T_0$ , there exists a proof from  $T_0$  with at most  $\ell^n + n$  symbols of the same conclusion.

It does not matter whether we mean sequence-proofs or tree-proofs here because these two notions are polynomially related to each other [7, Theorem 4.1]. With sequence-proofs in mind, it is easy to see that for every finitely axiomatized  $\mathcal{L}_\Pi$  theory  $T$ , if one has an  $\omega$ -interpretation of  $T$  in an  $\mathcal{L}_I$  theory  $T_0$ , then  $T$  has at most polynomial speed-up over  $T_0$ .

**Corollary 4** (Hájek, Avigad).  $\text{WKL}_0$  has at most polynomial speed-up over  $\text{I}\Sigma_1$ .

*Proof.* Since  $\text{WKL}_0$  is finitely axiomatizable [8, Lemma VIII.2.10], this follows directly from Theorem 2.  $\square$

We do not know whether  $\text{WKL}_0^*$  has at most polynomial speed-up over  $\text{I}\Delta_0 + \text{exp}$ .

#### ACKNOWLEDGEMENTS

The author would like to thank Ali Enayat for his encouragement, and for pointing out the connections with interpretations and speed-ups. He is also grateful to Enayat, the anonymous referee, and the editor for suggesting a number of improvements in the presentation of this paper.

#### REFERENCES

- [1] Jeremy Avigad, *Formalizing forcing arguments in subsystems of second-order arithmetic*, Ann. Pure Appl. Logic **82** (1996), no. 2, 165–191, DOI 10.1016/0168-0072(96)00003-6. MR1419805 (97k:03067)
- [2] Petr Hájek, *Interpretability and fragments of arithmetic*, Arithmetic, proof theory, and computational complexity (Prague, 1991), Oxford Logic Guides, vol. 23, Oxford Univ. Press, New York, 1993, pp. 185–196. MR1236462 (94f:03066)
- [3] Petr Hájek and Pavel Pudlák, *Metamathematics of first-order arithmetic*, Perspectives in Mathematical Logic, Springer-Verlag, Berlin, 1993. MR1219738 (94d:03001)
- [4] Aleksandar Djordje Ignjatovic, *Fragments of first and second-order arithmetic and length of proofs*, ProQuest LLC, Ann Arbor, MI, 1990. Thesis (Ph.D.)—University of California, Berkeley. MR2685328
- [5] Leszek Aleksander Kołodziejczyk and Keita Yokoyama, *Categorical characterizations of the natural numbers require primitive recursion*, Ann. Pure Appl. Logic **166** (2015), no. 2, 219–231, DOI 10.1016/j.apal.2014.10.003. MR3281817
- [6] J. B. Paris, *Some conservation results for fragments of arithmetic*, Model theory and arithmetic (Paris, 1979), Lecture Notes in Math., vol. 890, Springer, Berlin-New York, 1981, pp. 251–262. MR645006 (83f:03060)
- [7] Pavel Pudlák, *The lengths of proofs*, Handbook of proof theory, Stud. Logic Found. Math., vol. 137, North-Holland, Amsterdam, 1998, pp. 547–637, DOI 10.1016/S0049-237X(98)80023-2. MR1640332 (99i:03073)
- [8] Stephen G. Simpson, *Subsystems of second order arithmetic*, 2nd ed., Perspectives in Logic, Cambridge University Press, Cambridge; Association for Symbolic Logic, Poughkeepsie, NY, 2009. MR2517689 (2010e:03073)
- [9] Stephen G. Simpson and Rick L. Smith, *Factorization of polynomials and  $\Sigma_1^0$  induction*, Ann. Pure Appl. Logic **31** (1986), no. 2-3, 289–306, DOI 10.1016/0168-0072(86)90074-6. Special issue: second Southeast Asian logic conference (Bangkok, 1984). MR854297 (88b:03093)

KURT GÖDEL RESEARCH CENTER FOR MATHEMATICAL LOGIC, UNIVERSITY OF VIENNA, AUSTRIA  
*E-mail address:* tin.lok.wong@univie.ac.at