

## HOLOMORPHIC ISOMETRIES OF THE COMPLEX UNIT BALL INTO IRREDUCIBLE BOUNDED SYMMETRIC DOMAINS

NGAIMING MOK

(Communicated by Lei Ni)

**ABSTRACT.** Let  $\Omega \subset S$  be an irreducible bounded symmetric domain of rank  $\geq 2$  embedded as an open subset in its dual Hermitian symmetric manifold  $S$  of the compact type. Write  $c_1(S) = (p+2)\delta$ ,  $\delta \in H^2(S, \mathbb{Z}) \cong \mathbb{Z}$  being the positive generator. We prove that there exists a nonstandard holomorphic embedding of the  $(p+1)$ -dimensional complex unit ball  $B^{p+1}$  into  $\Omega$  which is isometric with respect to canonical Kähler-Einstein metrics  $g$  resp.  $h$  normalized so that minimal disks are of constant Gaussian curvature  $-2$ . We construct such holomorphic isometries using varieties of minimal rational tangents (VMRTs). We also prove that  $n \leq p+1$  for any holomorphic isometry  $f : (B^n, g) \rightarrow (\Omega, h)$ . Our proofs rely on an extension theorem for holomorphic isometries of Mok (2012), the asymptotic behavior due to Klembeck (1978) of standard complete Kähler metrics on strictly pseudoconvex domains, and the fine structure of boundaries of bounded symmetric domains in their Harish-Chandra realizations due to Wolf (1972).

On a bounded domain  $U \Subset \mathbb{C}^m$  on the Euclidean space we denote by  $ds_U^2$  the Bergman metric. Let  $D \Subset \mathbb{C}^n$  and  $\Omega \Subset \mathbb{C}^N$  be bounded domains on Euclidean spaces. In Mok [11] (2012) we studied germs of holomorphic isometries between these domains. As a special case of [11, Theorem 2.2.1] (cf. also Remarks after the proof of the theorem), we proved

**Theorem 1** (Mok [11]). *Let  $\lambda > 0$  and  $f : (D, \lambda ds_D^2; x_0) \rightarrow (\Omega, ds_\Omega^2; y_0)$  be a holomorphic isometry with respect to the Bergman metric up to a normalizing constant. Assume that the Bergman metrics on  $D$  and  $\Omega$  are complete, that the Bergman kernel  $K_D(z, w)$  on  $D$  extends to a rational function in  $(z, \bar{w})$ , and that analogously the Bergman kernel  $K_D(\xi, \zeta)$  extends to a rational function in  $(\xi, \bar{\zeta})$ . Then,  $f$  extends to a proper holomorphic isometric embedding  $F : (D, \lambda ds_D^2) \rightarrow (\Omega, ds_\Omega^2)$ . Moreover,  $\text{Graph}(f) \subset D \times \Omega$  extends to an affine-algebraic subvariety  $V \subset \mathbb{C}^n \times \mathbb{C}^N$ .*

In particular, the extension theorem above applies to the case where  $D \Subset \mathbb{C}^n$  and  $\Omega \Subset \mathbb{C}^N$  are bounded symmetric domains in their Harish-Chandra realizations. When  $D$  is also irreducible and of rank  $\geq 2$ , Clozel-Ullmo [2] (2003) observed that the proof of Hermitian metric rigidity in Mok [8], [9] already yields the total geodesy of  $f$ . Thus, nonstandard (i.e., not totally geodesic) holomorphic isometries from an irreducible bounded symmetric domain  $D$  into a (not necessarily irreducible) bounded symmetric domain  $\Omega$  may exist only in the case where  $D$  is of rank 1, i.e., biholomorphic to the complex unit ball  $B^n$  of dimension  $n \geq 1$ .

---

Received by the editors December 28, 2015.

2010 *Mathematics Subject Classification.* Primary 53C35, 53C55.

This research was partially supported by the GRF 7046/10 of the HKRGC, Hong Kong.

In Mok [11] we constructed examples of nonstandard holomorphic isometries of the Poincaré disk into polydisks and into the Siegel upper half-plane of genus 3. It was unknown whether there exist nonstandard holomorphic isometries of the complex unit ball of dimension  $\geq 2$  into bounded symmetric domains, and the question was raised in Mok [10].

In this article we prove an existence theorem on nonstandard holomorphic isometric embeddings from the complex unit ball of a certain specific dimension into a given irreducible bounded symmetric domain of rank  $\geq 2$ , as follows.

**Main Theorem.** *Let  $\Omega \Subset \mathbb{C}^N$  be an irreducible bounded symmetric domain of rank  $\geq 2$  and denote by  $S$  the irreducible Hermitian symmetric manifold of the compact type dual to  $\Omega$ . Denoting by  $\delta \in H^2(S, \mathbb{Z}) \cong \mathbb{Z}$  the positive generator of the second integral cohomology group of  $S$ , we write  $c_1(S) = (p+2)\delta$ . Then, there exists a non-standard proper holomorphic isometric embedding  $F : (B^{p+1}, ds_{B^{p+1}}^2) \hookrightarrow (\Omega, ds_\Omega^2)$ .*

## 1. PRELIMINARIES AND PROOF OF THE MAIN THEOREM

Denote by  $\Omega \subset S$  the Borel embedding of  $\Omega$  as an open subset of its dual Hermitian symmetric manifold  $S$  of the compact type. For instance, in the case of type-I domains  $D_{p,q}^I = \{Z \in M(p, q; \mathbb{C}) : I_q - \overline{Z}^t Z > 0\}$ , where  $M(p, q; \mathbb{C})$  denotes the complex vector space of  $p$ -by- $q$  matrices with complex coefficients, we have the Borel embedding  $D_{p,q}^I \subset G(p, q)$ ,  $G(p, q)$  being the Grassmannian of complex  $q$ -planes of  $W \cong \mathbb{C}^{p+q}$ . The construction of holomorphic isometric embeddings of the complex unit ball in the Main Theorem for irreducible bounded symmetric domains  $\Omega$  is related to geometric concepts on the compact dual  $S$ , which is a Fano manifold of Picard number 1 uniruled by projective lines, viz., rational curves  $C$  representing the generator of  $H^2(X, \mathbb{Z}) \cong \mathbb{Z}$ , which are realized as projective lines when  $S$  is canonically embedded into the projective space  $\mathbb{P}(H^0(S, \mathcal{O}(1))^*)$ ,  $\mathcal{O}(1)$  being the positive generator of the Picard group  $\text{Pic}(S) \cong \mathbb{Z}$ . In a general theory of projective uniruled manifolds, we have defined the notion of varieties of minimal rational tangents (VMRTs); cf. Hwang-Mok [6] (1999). In the case of a projective submanifold  $X$  uniruled by projective lines, as is the case of  $S \hookrightarrow \mathbb{P}(H^0(S, \mathcal{O}(1))^*)$ , the VMRT at a general point  $x \in X$  is a smooth projective submanifold  $\mathcal{C}_x(X) \subset \mathbb{P}(T_x(X))$  consisting of projectivizations of nonzero vectors  $\alpha \in T_x(X)$  tangent to projective lines  $\ell$  passing through  $x$ . Here by a general point we mean a point  $x \in X$  through which every minimal rational curve passing through  $x$  is free, i.e.,  $T_X|_\ell$  is a direct sum of holomorphic line bundles of degree  $\geq 0$ . In our case of irreducible Hermitian symmetric manifolds of the compact type  $S$  all points  $x \in S$  are equivalent under  $\text{Aut}(S)$ , and  $\mathcal{C}_x(S) \subset \mathbb{P}T_x(S)$  is itself a Hermitian symmetric manifold of the compact type, either of rank 2 embedded by the minimal embedding, or of rank 1, thus biholomorphic to  $\mathbb{P}^n$ , and embedded either by the minimal embedding (i.e., by  $\mathcal{O}(1)$ ) or by the Veronese embedding (i.e., by  $\mathcal{O}(2)$ ). For instance, in the case of  $G(p, q)$ , we have  $\mathcal{C}_x(S) \cong \mathbb{P}^{p-1} \times \mathbb{P}^{q-1} \subset \mathbb{P}(\mathbb{C}^p \otimes \mathbb{C}^q) \cong \mathbb{P}^{pq-1}$  given by the Segre embedding. In the cases where  $p = 1$  or  $q = 1$ ,  $S$  is the projective space and  $\mathcal{C}_x(S) = \mathbb{P}T_x(S)$ . The cases where  $\mathcal{C}_x(S) \subset \mathbb{P}T_x(X)$  are given by the Veronese embedding of projective spaces correspond precisely to type-III Hermitian symmetric manifolds of the compact type, viz., where  $S$  is the Lagrangian Grassmannian consisting of Lagrangian vector subspaces in a complex symplectic vector space.

Write  $G_0$  for the identity component of the holomorphic isometry group of  $(\Omega, ds_\Omega^2)$ . We have  $\Omega = G_0/K$ , where  $K \subset G_0$  is the isotropy subgroup at a base point  $0 \in \Omega$ . We denote by  $G^{\mathbb{C}}$  the identity component of  $\text{Aut}(S)$  and by  $\mathfrak{g}^{\mathbb{C}}$  its Lie algebra.  $G^{\mathbb{C}}$  contains  $G_0$  as a noncompact real form.  $G^{\mathbb{C}}$  acts transitively on  $S$  and we have  $S = G^{\mathbb{C}}/P$  as a rational homogeneous manifold, where  $P \subset G^{\mathbb{C}}$  is a maximal parabolic subgroup. The Borel embedding  $\beta : \Omega \hookrightarrow S$  is given by  $\beta(gK) = gP \in G^{\mathbb{C}}/P \cong S$ . In the embedding  $\Omega$  is contained in the orbit of  $0 = eK \in \Omega$  under  $M^+ = \exp(\mathfrak{m}^+)$ , where  $\mathfrak{g}^{\mathbb{C}} = \mathfrak{m}^+ \oplus \mathfrak{k}^{\mathbb{C}} \oplus \mathfrak{m}^-$  is the Harish-Chandra decomposition in standard notation. Here  $\mathfrak{k}^{\mathbb{C}}$  is the complexification of the Lie algebra  $\mathfrak{k}$  of the isotropy subgroup  $K \subset G_0$ ,  $\mathfrak{k}^{\mathbb{C}} \oplus \mathfrak{m}^-$  is the Lie algebra  $\mathfrak{p}$  of  $P \subset G^{\mathbb{C}}$ , and hence  $\mathfrak{m}^+ \subset \mathfrak{g}^{\mathbb{C}}$  can be identified with  $T_0(S) = T_0(G^{\mathbb{C}}/P) = \mathfrak{g}^{\mathbb{C}}/\mathfrak{p} \cong \mathfrak{m}^+$ . The Harish-Chandra realization is given by  $\Omega = G_0 \cdot 0 \in M^+ \cdot 0 \cong \mathfrak{m}^+ \cong \mathbb{C}^N$ . Here  $M^+ \cdot 0$  is the  $M^+$ -orbit of  $0 = eP$ . We note that  $M^+ \cdot 0 \cong \mathfrak{m}^+$  is a Zariski open subset of  $S = G^{\mathbb{C}} \cdot 0$ , and we have the inclusions  $\Omega \Subset \mathbb{C}^N \subset S$  incorporating both the Harish-Chandra realization and the Borel embedding. For details we refer the reader to Wolf [16] (1972).

In our proof of the Main Theorem we will make use of algebro-geometric objects to derive a holomorphic isometry by taking the limit of certain “algebraic” subsets of  $\Omega$  under a one-parameter family of automorphisms in  $\text{Aut}(\Omega)$ . More precisely, we will rely on the one hand on VMRTs of the Hermitian symmetric manifold  $S$  dual to  $\Omega$  and on the other hand on asymptotic properties of restrictions of canonical Kähler metrics to strictly pseudoconvex submanifolds exiting  $\Omega$  along regular boundary points. For the latter half we have the following well-known statement about the geometry of strictly pseudoconvex domains, which goes back to Klembeck [7] (1978); cf. also Cheng-Yau [1] (1980).

**Proposition 1** (Klembeck [7]). *Let  $U \subset \mathbb{C}^n$  be a Euclidean domain,  $\rho$  be a smooth real function on  $U$  and  $b$  be a point on  $U$ . Suppose  $\rho(b) = 0$  and  $d\rho(x) \neq 0$  for any  $x \in U$ . Assume furthermore that  $\rho$  is strictly plurisubharmonic on  $U$ , i.e.,  $\sqrt{-1}\partial\bar{\partial}\rho > 0$  on  $U$ . Let  $U' \subset U$  be the open subset defined by  $\rho < 0$ , and  $s$  be the Kähler metric on  $U'$  for which the Kähler form is given by  $\omega = \sqrt{-1}\partial\bar{\partial}(-\log(-\rho))$ . Then  $(U', s)$  is asymptotically of constant holomorphic sectional curvature  $-2$  at  $b$ , i.e., at a point  $x \in U'$  choosing  $\epsilon(x) \geq 0$  to be the smallest nonpositive number such that holomorphic sectional curvatures of  $(U', s)$  at  $x$  are bounded between  $-2 - \epsilon(x)$  and  $-2 + \epsilon(x)$ ; then  $\epsilon(x) \rightarrow 0$  as  $x \in U'$  approaches  $b \in \partial U' \cap U$ .*

We note that our convention for a Kähler metric  $g$  is given in local holomorphic coordinates by  $g = 2\text{Re}(\sum g_{\alpha\bar{\beta}}(z)dz^\alpha \otimes d\bar{z}^\beta)$  (with a factor of 2) with Kähler form  $\omega_g = \sqrt{-1}\sum g_{\alpha\bar{\beta}}(z)dz^\alpha \wedge d\bar{z}^\beta$ , which explains why the asymptotic holomorphic sectional curvature is  $-2$  in place of  $-4$  as in [7].

We are now ready to give a proof of the Main Theorem.

*Proof of the Main Theorem.* We will construct an explicit example of a bona fide holomorphic isometric embedding  $\Phi : (B^{p+1}, ds_{B^{p+1}}^2) \hookrightarrow (\Omega, ds_\Omega^2)$ . The ensuing discussion involves the fine structure of bounded symmetric domains, for which we refer the reader to Wolf [16] for details. Let  $\Omega \Subset \mathbb{C}^N \subset S$  be the inclusions incorporating both the Borel embedding of  $\Omega$  into  $S$ , the compact dual manifold of  $\Omega$ , and the Harish-Chandra realization  $\Omega \Subset \mathbb{C}^N$  of  $\Omega$ , where  $\mathbb{C}^N$  is the orbit of

$0 = eP$  under  $M^+ = \exp(\mathfrak{m}^+)$  as described above. Let  $\mathcal{C}_0 \subset \mathbb{P}T_0(\Omega)$  be the VMRT of  $S$  at  $0 \in \Omega \subset S$ . ( $\mathcal{C}_0$  is equivalently the variety of tangents to minimal disks on  $(\Omega, ds_\Omega^2)$ .) For  $x \in S$  define

$$\mathcal{V}_x = \bigcup \{ \ell : \ell \text{ is a minimal rational curve on } S \text{ through } x \},$$

and we call  $\mathcal{V}_x$  the variety swept out by minimal rational curves passing through  $x$ . For  $x \in S$  we define  $V_x := \mathcal{V}_x \cap \Omega \subset \Omega$ . When  $x \in \Omega$ ,  $V_x \subset \Omega$  is a  $(p+1)$ -dimensional subvariety which is smooth except for an isolated singularity at  $0 \in V$  in the case where  $\text{rank}(\Omega) := r \geq 2$ . Write  $K \subset \text{Aut}_0(\Omega)$  for the isotropy subgroup at  $0 \in \Omega$ . By the Polydisk Theorem, there exists a maximal polydisk  $P \cong \Delta^r$  embedded in  $\Omega$  as a totally geodesic complex submanifold. Moreover, all maximal polydisks on  $\Omega$  are equivalent to each other under  $\text{Aut}_0(\Omega)$ , and we have furthermore

$$\Omega = \bigcup \{ \gamma P : \gamma \in K \}.$$

For  $\eta \in T_0(\Omega)$ ,  $\eta$  is equivalent to some  $\xi = \text{diag}(\eta_1, \dots, \eta_r) \in T_0(P)$  under  $K$ -action. Moreover, there is an injective group homomorphism  $\text{Aut}(P) \hookrightarrow \text{Aut}_0(\Omega)$  for the full group  $\text{Aut}(P)$  of automorphisms of  $P$  (which contains the permutations of Cartesian factors). Thus, for  $\eta \neq 0$ ,  $[\eta] \in \mathcal{C}_0(S)$  if and only if the Euclidean disk  $\mathbb{C}\eta \cap \Omega$  is of radius 1, and  $\mathcal{C}_0(S)$  is a single orbit under the action of  $K$  on  $\mathbb{P}T_0(S)$ . We have

$$V_0 = \mathcal{V}_0 \cap B^N; \text{ hence } \partial V_0 \subset \partial B^N \text{ is strictly pseudoconvex.}$$

Choose the minimal disk  $D = \Delta \times \{0\} \subset V$ . From  $\text{Aut}(D) \hookrightarrow \text{Aut}(\Omega)$  we get a one-parameter family of transvections  $\Phi = \{ \varphi_t : -1 < t < 1 \} \subset \text{Aut}_0(\Omega)$ ,  $\varphi_t|_D \in \text{Aut}(D)$ ,  $\varphi_t(z, 0) = \left( \frac{z+t}{1+tz}, 0 \right)$ . For  $t \in (-1, 1)$  and the point  $\varphi_t(0) = (t, 0) \in D = \Delta \times \{0\} \subset \Omega$ , we sometimes write  $\mathcal{V}_t$  for  $\mathcal{V}_{(t,0)}$ , etc. Here Harish-Chandra coordinates on  $\Omega$ , which are unique up to unitary transformations, are chosen such that a maximal polydisk is given by  $P = \{ (z_1, \dots, z_r; 0, \dots, 0) : |z_i| < 1 \text{ for } 1 \leq i \leq r \}$ . In what follows, for a subset  $E \subset \mathbb{C}^N$  and  $y \in \mathbb{C}^n$  we write  $E+y := \{ x+y : x \in E \}$ . Since  $\text{Aut}(S)$  preserves the set of minimal rational curves on  $S$ , for any  $\gamma \in \text{Aut}(S)$  we have  $\mathcal{C}_{\gamma(x)}(S) = [d\gamma(x)](\mathcal{C}_x(S))$ , where for a linear isomorphism  $\lambda : E \rightarrow E'$  between two finite-dimensional vector spaces we denote by  $[\lambda] : \mathbb{P}(E) \rightarrow \mathbb{P}(E')$  the induced projective linear isomorphism. In particular, for  $\gamma = \exp(\eta)$ ,  $\eta \in \mathfrak{m}^+$ , belonging to the vector group of translations  $M^+ = \exp(\mathfrak{m}^+)$ , we have  $\mathcal{V}_\eta \cap \mathbb{C}^N = (\mathcal{V}_0 \cap \mathbb{C}^N) + \eta$ . Now for  $t \in (-1, 1)$ ,  $\mathcal{V}_t = \mathcal{V}_{(t,0)}$  is uniquely determined by the variety of minimal rational tangents  $\mathcal{C}_t(S) \subset \mathbb{P}T_t(S)$ . Thus  $\mathcal{V}_t = \gamma(\mathcal{V}_0)$  for any  $\gamma \in \text{Aut}(S)$  such that  $\gamma(0) = (t, 0)$ . In particular, we may take  $\gamma = \exp(\eta)$ , where  $\eta = \varphi_t(0) = (t, 0) \in \Omega \cong \mathbb{C}^N \cong \mathfrak{m}^+$ , i.e.,  $\gamma|_{\mathbb{C}^N}$  is the Euclidean translation by  $(t, 0)$ . As a consequence,  $V_t = \mathcal{V}_t \cap \Omega = (\mathcal{V}_0 + (t, 0)) \cap \Omega$ . Letting  $t \rightarrow -1$  and writing  $q$  for  $(-1, 0) \in \partial D \subset \partial \Omega$ ,  $V_t$  converges to  $V_q$  as subvarieties in  $\Omega$  as  $t \rightarrow -1$ . Any point  $x$  on  $V_q$  is the limit of points  $x_t \in V_t$  as  $t \rightarrow -1$ . More explicitly,  $\mathcal{V}_q = \mathcal{V}_0 + q$ , so that  $x = x_0 + q$  for some  $x_0 \in \mathcal{V}_0$ , and it suffices to take  $x_t = x_0 + (t, 0)$  which lies on  $\Omega$  for  $t$  sufficiently close to  $-1$ , so that  $x_t \in (\mathcal{V}_0 + (t, 0)) \cap \Omega = \mathcal{V}_t \cap \Omega = V_t$ .

Denote by  $h$  the canonical Kähler-Einstein metric on  $\Omega$  such that minimal disks are of Gaussian curvature  $-2$ . Then, for any  $x \in V_q$ , the germ of Kähler manifold  $(V_q, h|_{V_q}; x)$  is the limit of  $(V_t, h|_{V_t}; x_t)$  for some  $x_t \in V_t$  converging to  $x$  as  $t \rightarrow -1$ . The latter is equivalent under the action of  $\Phi$  to  $(V_0, h|_{V_0}; \varphi_t^{-1}(x_t))$ . Note that  $\varphi_t^{-1}(0) = (-t, 0) \rightarrow (1, 0) := p$  as  $t \rightarrow -1$ .

On  $\Omega$  we denote by  $d(\cdot, \cdot)$  the distance function with respect to the complete Kähler-Einstein metric  $h$ , which is invariant under  $\text{Aut}(\Omega)$ . Observe that

$$d(\varphi_t^{-1}(x_t), \varphi_t^{-1}(0)) = d(x_t, 0)$$

is bounded. For  $x \in V_0$  write  $\delta(x) = 1 - \|x\|$ , where  $\|\cdot\|$  stands for the Euclidean norm, and denote by  $g_e$  the Euclidean metric. From the strict pseudoconvexity of  $V_0$  at the boundary point  $p \in \partial V_0$ , we know that  $g|_{V_0} \geq \frac{C}{\delta} g_e|_{V_0}$  for some constant  $C > 0$ , i.e.,  $g_e|_{V_0} \leq \frac{\delta}{C} g|_{V_0}$ . It follows that  $\|\varphi_t^{-1}(x_t) - \varphi_t^{-1}(0)\| \rightarrow 0$  as  $t \rightarrow -1$ . Since  $\varphi_t^{-1}(0)$  converges to  $p = (1, 0)$  as  $t \rightarrow -1$ , it follows that, as  $t \rightarrow -1$ , the points  $\varphi_t^{-1}(x_t)$  on  $\Omega$  converge in  $\mathbb{C}^N$  to  $p$  too.

By Proposition 1,  $(V_0, h|_{V_0})$  is asymptotically of constant holomorphic sectional curvature  $-2$  as points approach  $p \in \partial V$ . Thus,  $V_q$  is of constant holomorphic sectional curvature  $-2$  at  $x \in V$ . Since  $x \in V$  is arbitrary,  $V \subset \Omega$  is of constant holomorphic sectional curvature  $-2$ . It follows that  $(V_q, h|_{V_q})$  is locally holomorphically isometric to  $(B^{p+1}, g)$ . From the extension theorem of Mok [11] on holomorphic isometries (Theorem 1 here),  $V_q \subset \Omega$  is actually the image of a *bona fide* holomorphic isometric embedding  $F : (B^{p+1}, g) \hookrightarrow (\Omega, h)$ . Writing  $Z := F(B^{p+1}) \subset \Omega$ , for any  $x \in Z$  and any vector  $\xi \in T_x(Z)$  of unit length with respect to  $(\Omega, h)$ , denoting by  $R$  the curvature tensor with respect to specified Kähler metrics, we have

$$R_{\xi\bar{\xi}\xi\bar{\xi}}(Z, h|_Z) = R_{\xi\bar{\xi}\xi\bar{\xi}}(\Omega, h) - \|\sigma(\xi, \xi)\|^2,$$

where  $\sigma$  denotes the second fundamental form of  $(Z, h|_Z) \hookrightarrow (\Omega, h)$  and  $\|\cdot\|$  denotes the norm measured with respect to metrics induced from  $h$ . As is well known,  $[\xi] \in \mathcal{C}_x(S)$  if and only if  $R_{\xi\bar{\xi}\xi\bar{\xi}}(\Omega, h) = -2$  (cf. Mok [9, Appendix III, Proposition 1, p. 242ff]). Since  $(Z, h|_Z)$  is of constant holomorphic sectional curvature  $-2$  it follows that  $\sigma(\xi, \xi) = 0$  if and only if  $\xi$  is a minimal rational tangent on  $S$ . Since  $\text{rank}(S) \geq 2$ , we have  $\mathcal{C}_x(S) \subsetneq \mathbb{P}T_x(S)$ , so that  $\sigma(\xi, \xi) \neq 0$  for a generic nonzero vector  $\xi \in T_x(\Omega)$ , hence  $(Z, h|_Z) \hookrightarrow (\Omega, h)$  is a fortiori nonstandard.

For the proof of the Main Theorem it remains to show that  $F : (B^{p+1}, ds_{B^{p+1}}^2) \hookrightarrow (\Omega, ds_\Omega^2)$ , i.e.,  $F : B^{p+1} \hookrightarrow \Omega$  is also a holomorphic isometry with respect to the Bergman metric, viz.,  $F^*(ds_\Omega^2) = ds_{B^{p+1}}^2$ . For any bounded homogeneous domain  $U \Subset \mathbb{C}^m$ , the Bergman metric  $ds_U^2$  is of constant Ricci curvature  $-1$  (cf. Mok [9, p. 59], proof of Proposition 3). To deduce  $F^*(ds_\Omega^2) = ds_{B^{p+1}}^2$  from the statement  $F^*h = g$  it suffices to check that  $g = \nu ds_{B^{p+1}}^2, h = \nu ds_\Omega^2$  for the same constant  $\nu$ . Equivalently, it suffices to check that the Kähler-Einstein manifolds  $(B^{p+1}, g)$  and  $(\Omega, h)$  have the same Kähler-Einstein constants.

For the complex unit ball we have  $\text{Ric}(B^{p+1}, g) = -(2 + p)$  since  $(B^{p+1}, g)$  is of constant holomorphic sectional curvature  $-2$ , while holomorphic bisectional curvatures  $R_{\alpha\bar{\alpha}\beta\bar{\beta}}(g) = -1$  whenever  $\alpha$  and  $\beta$  are orthogonal unit vectors (with respect to  $g$ ) at some  $x \in B^{p+1}$ .

As to  $(\Omega, h)$  we consider the dual Hermitian symmetric manifold  $S$  of the compact type. Let  $\mathfrak{g}_0 = \mathfrak{k} \oplus \sqrt{-1}\mathfrak{m}$  be the Cartan decomposition of the Lie algebra  $\mathfrak{g}_0$  of  $G_0$ . Let  $G_c \subset G^\mathbb{C}$  be the compact real form with Lie algebra  $\mathfrak{g}_c = \mathfrak{k} \oplus \mathfrak{m}$ . Let  $h_c$  be the  $G_c$ -invariant Kähler-Einstein metric on  $S = G_c/K$  such that  $h_c$  agrees with  $h_0$  at  $0 = eP \in G^\mathbb{C}/P \cong S$ . Then  $(\Omega, h)$  and  $(S, h_c)$  constitute a dual pair of Hermitian symmetric manifolds such that the curvature tensors are opposite at 0. Thus, at 0 we have  $R_{i\bar{j}k\bar{\ell}}(h) = -R_{i\bar{j}k\bar{\ell}}(h_c)$ . (For more details on dual symmetric spaces we refer the reader to Helgason [4].) Let  $\alpha \in T_0(S)$  be a unit minimal rational tangent

and  $\ell \subset S$  be a minimal rational curve on  $S$  passing through 0 such that  $T_0(\ell) = \mathbb{C}\alpha$ . We have the Grothendieck decomposition  $T_S|_\ell \cong \mathcal{O}(2) \oplus \mathcal{O}(1)^p \oplus \mathcal{O}^q$ , where the direct summand  $\mathcal{O}(2) = T_\ell$ , and the direct sum  $\mathcal{O}(2) \oplus \mathcal{O}(1)^p \subset T_S|_\ell$  is uniquely determined. At 0, writing  $P_\alpha := (\mathcal{O}(2) \oplus \mathcal{O}(1)^p)_0$ , we have  $T_{[\alpha]}(\mathcal{C}_0(S)) \equiv P_\alpha/\mathbb{C}\alpha$ . We have the eigenspace decomposition at 0 of  $(S, h_c)$  for the Hermitian bilinear form  $H_\alpha(\xi, \eta) = R_{\alpha\bar{\alpha}\xi\bar{\eta}}(h_c)$  given by  $T_0(\Omega) = \mathbb{C}\alpha \oplus \mathcal{H}_\alpha \oplus \mathcal{N}_\alpha$ , where  $\mathbb{C}\alpha$  resp.  $\mathcal{H}_\alpha$  resp.  $\mathcal{N}_\alpha$  is the eigenspace of  $H_\alpha$  corresponding to the eigenvalues 2 resp. 1 resp. 0. Moreover, we have  $P_\alpha = \mathbb{C}\alpha \oplus \mathcal{H}_\alpha$ . By duality and computing at  $0 \in \Omega$  we see that the Ricci constant of  $(\Omega, h)$  is equal to  $-(2+p)$ . (We refer the reader to Hwang-Mok [6] for more details regarding minimal rational curves, and to Mok [9] for the eigenspace decomposition of  $H_\alpha$  on Hermitian symmetric manifolds.) Thus  $(B^{p+1}, g)$  and  $(\Omega, h)$  have the same Kähler-Einstein constant  $-(2+p)$ , and it follows that  $F^*ds_\Omega^2 = ds_{B^{p+1}}^2$  for the Bergman metrics  $ds_\Omega^2$  and  $ds_{B^{p+1}}^2$ , as desired. The proof of the Main Theorem is complete.  $\square$

## 2. UPPER BOUNDS ON DIMENSIONS OF COMPLEX UNIT BALLS ISOMETRICALLY EMBEDDABLE INTO AN IRREDUCIBLE BOUNDED SYMMETRIC DOMAIN

From our Main Theorem, which is an existence result on holomorphic isometries of complex unit balls  $B^n$  into irreducible bounded symmetric domains, it is natural to ask whether the dimension  $n = p + 1$  is the maximal possible dimension for holomorphic isometries. Maintaining the notation  $g$  resp.  $h$  for the canonical Kähler-Einstein metric on the complex unit ball  $B^n$  resp.  $\Omega$  such that minimal disks are of constant Gaussian curvature  $-2$ , we have the following estimate on the dimension of the complex unit ball  $B^n$ .

**Theorem 2.** *Let  $\Omega \subset S$  be the Borel embedding of an irreducible bounded symmetric domain  $\Omega$  into its dual Hermitian symmetric manifold  $S$  of the compact type, where  $\text{Pic}(S) \cong \mathbb{Z}$ , generated by the positive line bundle  $\mathcal{O}(1)$ . Let  $g$  resp.  $h$  be the canonical Kähler-Einstein metric on  $B^n$  resp.  $\Omega$  normalized so that minimal disks on  $B^n$  resp.  $\Omega$  are of constant Gaussian curvature  $-2$ . Let  $p = p(\Omega)$  be the nonnegative integer such that  $K_S^{-1} \cong \mathcal{O}(p+2)$ . Let  $n \geq 1$  and  $F : (B^n, g) \rightarrow (\Omega, h)$  be a holomorphic isometry (which is necessarily a proper holomorphic isometric embedding). Then  $n \leq p + 1$ .*

We need some preparation for the proof of Theorem 2 to understand the implication of the hypothesis  $F^*g = h$  on the boundary behavior of the map  $F$ . For an irreducible bounded symmetric domain  $\Omega \Subset \mathbb{C}^N$  in its Harish-Chandra realization, from the fine structure on  $\Omega$  (cf. Wolf [16]), the boundary  $\partial\Omega$  decomposes into exactly  $r$  orbits under the action of the identity component  $G_0$  of  $\text{Aut}(\Omega)$ . In what follows we will be dealing with semi-analytic sets  $A \subset \mathbb{C}^m$ , and a point on  $a \in A$  is said to be smooth if and only if there is some neighborhood  $U$  of  $a$  such that  $A \cap U \subset U$  is a real-analytic submanifold. We will denote by  $\text{Reg}(A) \subset A$  the set of smooth points of  $A$ . Thus  $\text{Reg}(A) \subset \mathbb{C}^m$  is a locally closed real-analytic submanifold. Since  $\Omega$  is irreducible and of rank  $r$ , we have a decomposition  $\partial\Omega = E_1 \cup \dots \cup E_r$  into  $r$  orbits under the action of  $G_0$ , where  $E_1 \subset \partial\Omega$  is the unique open orbit,  $E_r$  is the Shilov boundary, and  $E_k$  lies in the closure of  $E_\ell$  whenever  $k \leq \ell$ . By Mok-Ng [13, Lemma 2.2.3] the set  $\partial\Omega \subset \mathbb{C}^n$  is smooth at  $b \in \partial\Omega$  if and only if  $b$  lies on the unique open  $G_0$ -orbit  $E_1 \subset \partial\Omega$ , so that  $\text{Reg}(\Omega) = E_1$ .

Regarding the Bergman kernel on bounded symmetric domains we have the following lemma from Faraut-Korányi [3, pp. 76-77, especially Eqns. (3.4) and (3.9)].

**Lemma 1.** *Let  $\Omega \Subset \mathbb{C}^n$  be a bounded symmetric domain in its Harish-Chandra realization. Write  $K_\Omega(z, w)$  for the Bergman kernel of  $\Omega$ . Then, there exists a polynomial  $Q_\Omega(z, w)$  holomorphic in  $z$  and anti-holomorphic in  $w$  such that  $K_\Omega(z, w) = \frac{1}{Q_\Omega(z, w)}$ . More precisely,  $Q(z, w) = A_\Omega H(z, w)^m$ , where  $H(z, w)$  is some polynomial in  $(z_1, \dots, z_n; \overline{w_1}, \dots, \overline{w_n})$  invariant under  $K$ ,  $A_\Omega > 0$  is a constant and  $m$  is a positive integer, with the following property. Let  $P \cong \Delta^r$  be a maximal polydisk on  $\Omega$  passing through 0, identified as  $\Delta^r \times \{0\}$  in Harish-Chandra coordinates. Then,  $z = (z_1, \dots, z_r; 0) \in P$  and we have*

$$H(z, z) = (1 - |z_1|^2) \times \dots \times (1 - |z_r|^2) .$$

From Lemma 1, writing  $\rho(z) = -H(z, z)$ , the Kähler metric with Kähler form  $\omega_h = \sqrt{-1} \partial \bar{\partial} (-\log(-\rho))$  is a multiple of the Bergman metric. Denoting by  $P \subset \Omega$  the maximal polydisk as in Lemma 1, restricting to the minimal disk  $D = \Delta \times \{0\} \subset P \subset \Omega$  this gives the Poincaré disk with Gaussian curvature  $-2$ , and hence  $\omega_h$  is precisely the Kähler form of  $h$  as in the statement of Theorem 2.

For  $x \in \mathbb{C}^N$ , we consider  $d\rho(x) \in T_x^*(\mathbb{C}^N)$ . For the proof of Theorem 2 we will need the following lemma concerning  $d\rho$  on  $\partial\Omega$ .

**Lemma 2.**  *$E_1 \subset \partial\Omega$  is precisely the subset of  $\partial\Omega$  where  $d\rho$  is nonzero.*

*Proof.* To see this, note first of all that at a point  $p = (1, 0, \dots, 0; 0) \in \partial P \subset \mathbb{C}^r \times \{0\} := W$ , which lies on  $E_1$ , we have by direct computation using the formula in Lemma 1 that  $d\rho(p)|_{T_p(W)} \neq 0$ , so that, a fortiori we have  $d\rho(p) \neq 0$ . It follows that  $d\rho(\gamma(p)) \neq 0$  for any  $\gamma \in G_0$ , hence  $d\rho$  is nonzero on  $E_1$ . To prove the claim in the above we argue by contradiction. Suppose there exists a point  $q \in E_s$  for some  $s \geq 2$  such that  $d\rho(q) \neq 0$ . Since any open neighborhood  $U$  of  $q$  on  $\partial\Omega$  meets  $E_k$  for  $1 \leq k \leq s$ , for the sake of convenience and without loss of generality we may take  $s = 2$ .

Since  $d\rho(q) \neq 0$  by assumption, the level set  $Q_0 := \{\rho = 0\}$ , which is a real-analytic subset of  $\mathbb{C}^N$ , is smooth at  $q$ . Denote by  $Q$  the irreducible component of  $Q_0$  containing  $q \in E_2$ . We have checked that  $E_1 \subset Q_0$ , and  $q \in Q$  by assumption, hence  $E_2 = G_0 \cdot q$  also lies on  $Q$ , by  $G_0$ -invariance of  $Q$ . Since  $E_2 \subset \overline{E_1}$ , we have  $E_1 \cup E_2 \subset Q$ . On the other hand, assuming without loss of generality that  $q$  belongs to the closure  $\overline{P}$  of the maximal polydisk  $P$  and, restricting to the complex vector subspace  $W \subset \mathbb{C}^N$  which contains  $P$  as an open set, we may take  $q \in \partial\Delta \times \partial\Delta \times \Delta^{r-2} \times \{0\} \subset \overline{\Omega}$ . Since  $Q \supset E_1$  we have  $Q \supset \partial\Delta \times \Delta^{r-1} \times \{0\}$ , hence  $Q \supset \partial\Delta \times \mathbb{C}^{r-1} \times \{0\}$  by the real-analyticity of  $Q$ , showing that  $Q - \overline{\Omega} \neq \emptyset$ . More precisely, for any open neighborhood  $U$  of  $q$  in  $Q$ ,  $U - \overline{\Omega} \neq \emptyset$ . From this we are going to get a contradiction, for topological reasons.

To this end note first of all that  $\partial\Omega$  is homeomorphic to the unit sphere  $S^{2N-1}$ . In fact, by the Hermann Convexity Theorem (Hermann [5]),  $\partial\Omega$  is the unit sphere with respect to some Banach norm on the complex vector space  $\mathbb{C}^N$ , and, denoting by  $\|\cdot\|$  the Euclidean norm on  $\mathbb{C}^N$ , the map  $\Phi : \partial\Omega \rightarrow S^{2N-1}$  defined by  $\Phi(x) = \frac{x}{\|x\|}$  is a homeomorphism of  $\partial\Omega$  onto the unit sphere  $S^{2N-1}$ . Consequently,  $\partial\Omega$  is a topological manifold. Let  $U_0$  be a neighborhood of  $q$  in  $\partial\Omega$ ,  $U_0 \subset E_1 \cup E_2$ , such that  $U_0$  is homeomorphic to  $\mathbb{R}^{2n-1}$ . The inclusion  $i : U_0 \hookrightarrow Q$  gives a continuous and injective map from  $U_0$  into the smooth part  $\text{Reg}(Q)$  of  $Q$ . Shrinking  $U_0$  if

necessary we may assume that  $U$  lies on a Euclidean coordinate chart. By Brouwer's Invariance of Domain Theorem in Algebraic Topology (cf. Spanier [15, p. 199, Theorem 6]),  $i$  is open, giving a homeomorphism of  $U_0$  onto an open neighborhood  $U = i(U_0)$  of  $q$  in  $Q$ . But by the last paragraph we have on the other hand  $U - \bar{\Omega} \neq \emptyset$ , a plain contradiction, showing that  $s \geq 2$  cannot happen. In other words  $s = 1$  and  $E_1 \subset \partial\Omega$  is precisely the locus over which  $d\rho \neq 0$ , as desired.  $\square$

For the proof of Theorem 2 we will furthermore need to know the structure of the unique open orbit  $E_1 \subset \partial\Omega$  as can be deduced from Wolf [16]. We refer to Mok-Tsai [14] for the notion of characteristic subspaces and to Mok [9] for geometric formulations in terms of curvature of the canonical Kähler-Einstein metric.

Let  $\Omega \Subset \mathbb{C}^N \subset S$  be an irreducible bounded symmetric domain in its Harish-Chandra realization as a Euclidean domain and in its embedding as an open subset of the dual Hermitian symmetric manifold  $S$  of the compact type. Let  $\alpha \in T_0(\Omega)$  be a minimal rational tangent and write  $T_0(\Omega) = \mathbb{C}\alpha \oplus \mathcal{H}_\alpha \oplus \mathcal{N}_\alpha$  for the eigenspace decomposition with respect to the Hermitian bilinear form  $H_\alpha(\xi, \eta) := R_{\alpha\bar{\alpha}\xi\bar{\eta}}(\Omega, h)$  corresponding to the eigenvalues  $-2$ ,  $-1$  and  $0$  (as given in the proof of Theorem 2). Then, there exists a unique totally geodesic complex submanifold  $\Omega'_\alpha \subset \Omega$  such that  $T_0(\Omega'_\alpha) = \mathcal{N}_\alpha$ .  $\Omega'_\alpha \subset \Omega$  is a characteristic subspace of maximal dimension in the sense of Mok-Tsai [14]. The complex submanifolds  $\Omega'_\alpha$  are all equivalent to each other under the action of the isotropy subgroup  $K \subset G_0$ . Moreover, denoting by  $(\Delta, g)$  the Poincaré disk with constant Gaussian curvature  $-2$  and by  $(\Omega', h')$  the bounded symmetric domain isometric to  $(\Omega'_\alpha, h|_{\Omega'_\alpha})$  for any minimal rational tangent  $\alpha \in T_0(\Omega)$ , by  $D_\alpha$  the minimal disk on  $\Omega$  passing through  $0$  such that  $T_0(D_\alpha) = \mathbb{C}\alpha$ , we have a holomorphic isometric embedding  $\Lambda_\alpha : (\Delta, g) \times (\Omega', h') \hookrightarrow \Omega$  onto a totally geodesic complex submanifold  $\Sigma \subset \Omega$  such that  $\Lambda_\alpha(0, 0) = 0$  and such that  $T_0(\Sigma_\alpha) = \mathbb{C}\alpha \oplus \mathcal{N}_\alpha = T_0(D_\alpha) \oplus T_0(\Omega'_\alpha)$ . We have  $\text{Reg}(\partial\Sigma_\alpha) \supset \Lambda_\alpha(\partial\Delta \times \Omega')$ . For any  $\theta \in \mathbb{R}$ , we have  $\Lambda_\alpha(\{e^{i\theta}\} \times \Omega') := \Omega'_{\alpha, \theta} \subset E_1$ . The full statement of the lemma below on the structure of  $\text{Reg}(\partial\Omega)$  is not strictly necessary for the proof of Theorem 2, but it is useful to put things in perspective as the proof will relate the construction in the Main Theorem to the formulation and proof of Theorem 2.

**Lemma 3.** *The real analytic manifold  $E_1 = \text{Reg}(\partial\Omega)$  decomposes into the disjoint union of complex manifolds of the form  $\Omega'_{\alpha, \theta}$ . Moreover, for  $t \in \Omega'_{\alpha, \theta} \subset E_1$ ,  $T_t(\Omega'_{\alpha, \theta})$  is precisely the 0-eigenspace of the Levi form  $\sqrt{-1}\partial\bar{\partial}\rho$  restricted to the complex tangent space  $T_t^{1,0}(\partial\Omega)$ .*

Here the complex tangent space  $T_t^{1,0}(\partial\Omega) \subset T^{1,0}(\mathbb{C}^N) \cong \mathbb{C}^N$  is the complex hyperplane  $E_t$  such that, denoting by  $T^{\mathbb{R}}(\cdot)$  the real tangent bundle,  $\text{Re}(E_t) \subset T^{\mathbb{R}}(\mathbb{C}^N) \cong \mathbb{R}^{2N}$  is the  $J$ -invariant real 2-dimensional vector subspace defined by  $\text{Re}(E_t) = T_t^{\mathbb{R}}(\partial\Omega) \cap JT_t^{\mathbb{R}}(\partial\Omega)$ , where  $J$  is the  $J$ -operator of the standard complex structure on  $\mathbb{C}^N$ . (For a point  $x \in \mathbb{C}^N$  we make no distinction between  $T_x(\mathbb{C}^N)$  and  $T_x^{1,0}(\mathbb{C}^N)$ .) The restriction of the Levi form (which depends on the choice of a defining function) to the complex tangent space  $T_t^{1,0}(\partial\Omega)$  is uniquely determined up to a positive multiplicative constant. Since  $\bar{\Omega} \subset \mathbb{C}^N$  is convex (Hermann [5]),  $E_1 = \text{Reg}(\partial\Omega)$  is in particular weakly pseudoconvex, so that the Levi form  $\sqrt{-1}\partial\bar{\partial}\rho$  is positive semi-definite on  $T_t^{1,0}(\partial\Omega)$ .

*Proof of Lemma 3.* The first statement is well known (cf. Wolf [16]). As to the second statement, since  $\rho|_{\Omega'_{\alpha,\theta}} \equiv 0$ , clearly  $T_t^{1,0}(\Omega'_{\alpha,\theta})$  lies on the 0-eigenspace  $L_t$  of the restriction of  $\sqrt{-1}\partial\bar{\partial}\rho$  to  $T_t^{1,0}(\partial\Omega)$ . Since  $G_0$  acts transitively on  $E_1$  the complex dimension of  $L_t$  is independent of  $t \in E_1$ . By the construction in the Main Theorem, writing  $\mathcal{V}_0 \subset S$  for the variety swept out by minimal rational curves passing through  $0 \in \Omega \Subset \mathbb{C}^N$ ,  $\mathcal{V}_0 \cap \partial\Omega = \mathcal{V}_0 \cap E_1 = \mathcal{V}_0 \cap \partial B^N$  is a smooth strictly pseudoconvex real hypersurface of  $\mathcal{V}_0$ . Thus for  $t \in \mathcal{V}_0 \cap E_1$ ,  $\sqrt{-1}\partial\bar{\partial}\rho$  is positive definite on  $T_t^{1,0}(\partial(\mathcal{V}_0 \cap \Omega)) = T_t^{1,0}(\mathcal{V}_0 \cap E_1)$ . Now  $T_t^{1,0}(\mathcal{V}_0 \cap E_1)$  is a complex hyperplane in  $T_t^{1,0}(\mathcal{V}_0) \cong \mathbb{C}^{p+1}$ . Thus,  $T_t^{1,0}(\mathcal{V}_0 \cap E_1) \subset T_t^{1,0}(\partial\Omega)$  is a complex  $p$ -dimensional vector subspace on which the Levi form of  $\rho$  is positive definite, so that  $T_t^{1,0}(\mathcal{V}_0 \cap E_1) \cap T_t^{1,0}(\Omega'_{\alpha,\theta}) = 0$ . But  $\dim_{\mathbb{C}}(\Omega'_{\alpha,\theta}) = \dim_{\mathbb{C}}(\mathcal{N}_{\alpha}) = q$ , and  $p + q = N - 1 = \dim_{\mathbb{C}}(T_t^{1,0}(\Omega'_{\alpha,\theta}))$ . It follows that the 0-eigenspace  $L_t$  is exactly the same as  $T_t^{1,0}(\Omega'_{\alpha,\theta})$ , as asserted.  $\square$

We are now ready to give a proof of Theorem 2, as follows.

*Proof of Theorem 2.* Let  $F : (B^n, g) \rightarrow (\Omega, h)$  be a holomorphic isometry. By Mok [11] (cf. Theorem 1 here),  $F$  is a proper holomorphic embedding, and  $\text{Graph}(F) \subset \mathbb{C}^n \times \mathbb{C}^N$  extends to an affine-algebraic subvariety. In particular, at a general point  $b \in \partial B^n$ , there exists an open neighborhood  $U$  of  $b$  in  $\mathbb{C}^n$  such that  $F|_{U \cap \partial B^n}$  extends to a holomorphic embedding  $F^{\sharp} : U \rightarrow \mathbb{C}^N$  with the property that  $F^{\sharp}(U \cap \partial B^n) \subset \partial\Omega$ . Recall that we have a decomposition  $\partial\Omega = E_1 \cup \dots \cup E_r$  into  $r$  orbits under the action of  $G_0$ . Taking  $U \cap \partial B^n$  to be connected there is a smallest positive integer  $s$ ,  $1 \leq s \leq r$  such that  $F^{\sharp}(U \cap \partial B^n) \subset \overline{E_s}$ . For this choice of  $s$  there is some (possibly empty) real-analytic subvariety  $A \subsetneq U \cap \partial B^n$  such that  $F^{\sharp}(\partial B^n - A) \subset E_s$ , and we will simply say that  $F^{\sharp}$  (or just  $F$ ) exits  $\partial\Omega$  along the stratum  $E_s$ .

We claim that the stratum  $E_s$  where  $F^{\sharp}$  exits  $\partial\Omega$  must necessarily be  $E_1$ . Suppose  $F^{\sharp}$  exits  $\partial\Omega$  along  $E_s$  for some  $s \geq 2$ . Let  $b \in U \cap \partial B^n$  be a general point and  $\Gamma$  be a connected component of the intersection of  $U$  with an affine line  $L$  passing through  $b$ . Then  $(\Gamma, g|_{\Gamma}) \hookrightarrow (B^n, g)$  is a totally geodesic Hermitian Riemann surface of constant Gaussian curvature  $-2$ . On the other hand,  $d((F^{\sharp})^*\rho)$  must vanish on  $T_b(\Gamma)$  by Lemma 2, so that  $(F^{\sharp})^*\rho|_{\Gamma}$  must vanish exactly to the order  $m$  for some  $m \geq 2$  at  $b \in \Gamma \cap \partial B^n$ . By a direct computation  $(\Gamma, F^*h|_{\Gamma})$  is asymptotically of Gaussian curvature  $\frac{-2}{m}$ , a plain contradiction. As a consequence, we have proven that  $F^{\sharp}$  exits along  $E_1 = \text{Reg}(\partial\Omega)$ .

To prove Theorem 2 it suffices now to consider the situation where  $b \in \partial B^n$  is a general point,  $U$  is a neighborhood of  $b$  in  $\mathbb{C}^n$ ,  $F^{\sharp}$  is a holomorphic embedding on  $U$ ,  $H := F^{\sharp}(U \cap \partial B^n) = F^{\sharp}(U) \cap E_1 \subset F^{\sharp}(U) := Z$ . Here  $H$ , being the image of the strictly pseudoconvex hypersurface  $U \cap \partial B^n \subset U$  under the embedding  $F^{\sharp}$ , must be a smooth strictly pseudoconvex hypersurface in  $Z$ . Thus, restricting to  $T_t^{1,0}(Z \cap E_1)$  for  $t = F^{\sharp}(b)$ ,  $b \in U \cap \partial B^n$ , the Levi form  $\sqrt{-1}\partial\bar{\partial}\rho$  must be positive definite. By Lemma 3, the kernel  $L_t$  of  $\sqrt{-1}\partial\bar{\partial}\rho$  on  $T_t^{1,0}(E_1)$  is of dimension exactly equal to  $q$ , and we conclude that  $(n - 1) + q \leq N - 1$ , hence  $n \leq N - q = p + 1$ , as desired.  $\square$

In the proof of Theorem 2 it was established that  $F^*h = g$  implies that  $\overline{F(B^n)} \cap \text{Reg}(\partial\Omega) \neq \emptyset$ . We note that when  $F$  is assumed to be a holomorphic isometry up

to some normalizing constant, the converse of the above statement also holds true. For the proof of the converse statement we need the following lemma.

**Lemma 4** (Mok [12, Lemma 7]). *Let  $W \subset \mathbb{C}^N$  be a connected open subset such that  $W \cap \partial\Omega \subset \text{Reg}(\partial\Omega)$ . Suppose  $\Lambda \subset W$  is a holomorphic curve such that  $\Lambda \cap \text{Reg}(\partial\Omega) \neq \emptyset$ . Then, for a general point  $t \in \Lambda \cap \partial\Omega$ , we have  $T_t^{1,0}(\Lambda) \cap T_t^{1,0}(\partial\Omega) = 0$ , where  $T_t^{1,0}(\partial\Omega)$  is the complex tangent space of  $\text{Reg}(\partial\Omega)$  at  $t$ .*

We proceed now to prove

**Theorem 3.** *Let  $n \geq 1$ ,  $\lambda > 0$ , and  $F : (B^n, \lambda g) \rightarrow (\Omega, h)$  be a holomorphic isometry such that  $\overline{F(B^n)} \cap \text{Reg}(\partial\Omega) \neq \emptyset$ . Then,  $\lambda = 1$  and  $n \leq p + 1$ .*

*Proof.* By assumption  $\overline{F(B^n)} \cap \text{Reg}(\partial\Omega) \neq \emptyset$ . Let  $b \in \partial B^n$  be a general point,  $U$  be a connected open neighborhood of  $b$  in  $\mathbb{C}^n$  such that  $F|_{U \cap B^n}$  admits an extension to a holomorphic embedding  $F^\sharp : U \rightarrow \mathbb{C}^N$ . Recall that  $\omega_g$  stands for the Kähler form of  $(B^n, g)$  and  $\omega_h$  stands for the Kähler form of  $(\Omega, h)$ . We have  $\omega_h = \sqrt{-1} \partial \bar{\partial}(-\log(-\rho))$  (where  $\rho(z) = -H(z, z)$  as in Lemma 1), so that  $F^* \omega_h = \sqrt{-1} \partial \bar{\partial}(-\log(-F^* \rho))$ , while  $\omega_g = \sqrt{-1} \partial \bar{\partial}(-\log(1 - \|z\|^2))$ .

By Lemma 4, when  $n = 1$  replacing  $b$  by some point on  $U \cap \partial B^1$  and shrinking  $U$  if necessary we may assume without loss of generality that  $(F^\sharp)^* \rho$  vanishes along  $U \cap \partial B^1$  exactly to the order 1, so that  $-F^* \rho = (1 - \|z\|^2) \mu$  on  $U$  for some smooth function  $\mu$  on  $U$  such that  $\mu|_{U \cap B^1}$  is strictly positive. For  $B^n$  of arbitrary dimension, one can slice  $U$  by complex lines passing through a general point  $b \in \partial B^n$  to conclude from Lemma 2 that  $d\rho(b)|_\Gamma \neq 0$  and hence a fortiori we have  $d\rho(b) \neq 0$ , so that we may also assume without loss of generality that  $(F^\sharp)^* \rho$  vanishes exactly to the order 1 on  $U \cap \partial B^n$ . Thus on  $U \cap B^n$ , we have also

$$-\log(-F^* \rho) = -\log(1 - \|z\|^2) + \log \mu$$

for a smooth function  $\mu$  on  $U$  which is strictly positive on  $U \cap \partial B^n$ . As a consequence, we have

$$\begin{aligned} F^*(\omega_h) &= \sqrt{-1} \partial \bar{\partial}(-\log(-F^* \rho)) \\ &= \sqrt{-1} \partial \bar{\partial}(-\log(1 - \|z\|^2)) + \sqrt{-1} \partial \bar{\partial} \mu = \omega_g + \sqrt{-1} \partial \bar{\partial} \mu. \end{aligned}$$

By assumption  $F^* \omega_h = \lambda \omega_g$ . On the other hand  $\sqrt{-1} \partial \bar{\partial} \mu$  is bounded as a  $(1, 1)$ -form on  $U \cap B^n$  by a multiple of the Euclidean Kähler form  $\beta$ , while  $\omega_g$  is bounded from below by  $\frac{A}{1 - \|z\|^2} \beta$  for some constant  $A > 0$ , from which it follows that  $\sqrt{-1} \partial \bar{\partial} \mu = o(\omega_g)$  on  $U \cap B^n$  as points approach  $\partial B^n$ . As a consequence,  $\lambda = 1$  and  $F^* \omega_g = \omega_h$ , and from Theorem 2 we conclude that  $n \leq p + 1$ , as desired.  $\square$

#### REFERENCES

- [1] Shiu Yuen Cheng and Shing Tung Yau, *On the existence of a complete Kähler metric on noncompact complex manifolds and the regularity of Fefferman's equation*, Comm. Pure Appl. Math. **33** (1980), no. 4, 507–544, DOI 10.1002/cpa.3160330404. MR575736
- [2] Laurent Clozel and Emmanuel Ullmo, *Correspondances modulaires et mesures invariantes* (French), J. Reine Angew. Math. **558** (2003), 47–83, DOI 10.1515/crll.2003.042. MR1979182
- [3] J. Faraut and A. Korányi, *Function spaces and reproducing kernels on bounded symmetric domains*, J. Funct. Anal. **88** (1990), no. 1, 64–89, DOI 10.1016/0022-1236(90)90119-6. MR1033914
- [4] Sigurdur Helgason, *Differential geometry, Lie groups, and symmetric spaces*, Pure and Applied Mathematics, vol. 80, Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London, 1978. MR514561

- [5] Robert Hermann, *Geometric aspects of potential theory in the symmetric bounded domains. II*, Math. Ann. **151** (1963), 143–149. MR0155342
- [6] Jun-Muk Hwang and Ngaiming Mok, *Varieties of minimal rational tangents on uniruled projective manifolds*, Several complex variables (Berkeley, CA, 1995), Math. Sci. Res. Inst. Publ., vol. 37, Cambridge Univ. Press, Cambridge, 1999, pp. 351–389. MR1748609
- [7] Paul F. Klembeck, *Kähler metrics of negative curvature, the Bergmann metric near the boundary, and the Kobayashi metric on smooth bounded strictly pseudoconvex sets*, Indiana Univ. Math. J. **27** (1978), no. 2, 275–282. MR0463506
- [8] Ngaiming Mok, *Uniqueness theorems of Hermitian metrics of seminegative curvature on quotients of bounded symmetric domains*, Ann. of Math. (2) **125** (1987), no. 1, 105–152, DOI 10.2307/1971290. MR873379
- [9] Ngaiming Mok, *Metric rigidity theorems on Hermitian locally symmetric manifolds*, Series in Pure Mathematics, vol. 6, World Scientific Publishing Co., Inc., Teaneck, NJ, 1989. MR1081948
- [10] Ngaiming Mok, *Geometry of holomorphic isometries and related maps between bounded domains*, Geometry and analysis. No. 2, Adv. Lect. Math. (ALM), vol. 18, Int. Press, Somerville, MA, 2011, pp. 225–270. MR2882446
- [11] Ngaiming Mok, *Extension of germs of holomorphic isometries up to normalizing constants with respect to the Bergman metric*, J. Eur. Math. Soc. (JEMS) **14** (2012), no. 5, 1617–1656, DOI 10.4171/JEMS/343. MR2966662
- [12] Ngaiming Mok, *Local holomorphic curves on a bounded symmetric domain in its Harish-Chandra realization exiting at regular points of the boundary*, Pure Appl. Math. Q. **10** (2014), no. 2, 259–288, DOI 10.4310/PAMQ.2014.v10.n2.a4. MR3267347
- [13] Ngaiming Mok and Sui Chung Ng, *Germs of measure-preserving holomorphic maps from bounded symmetric domains to their Cartesian products*, J. Reine Angew. Math. **669** (2012), 47–73. MR2980451
- [14] Ngaiming Mok and I Hsun Tsai, *Rigidity of convex realizations of irreducible bounded symmetric domains of rank  $\geq 2$* , J. Reine Angew. Math. **431** (1992), 91–122. MR1179334
- [15] E. H. Spanier, *Algebraic topology*, McGraw-Hill Book Co., New York-Toronto, Ontario-London, 1966.
- [16] Joseph A. Wolf, *Fine structure of Hermitian symmetric spaces*, Symmetric spaces (Short Courses, Washington Univ., St. Louis, Mo., 1969–1970), Dekker, New York, 1972, pp. 271–357. Pure and App. Math., Vol. 8. MR0404716

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF HONG KONG, POKFULAM ROAD, HONG KONG

*E-mail address:* nmok@hku.hk