

## ON THE REGULARITY OF VECTOR FIELDS UNDERLYING A DEGENERATE-ELLIPTIC PDE

ERIKA BATTAGLIA, STEFANO BIAGI, AND GIULIO TRALLI

(Communicated by Jeremy Tyson)

ABSTRACT. In this note we discuss the relationship, under an algebraic constant rank condition, between the regularity of the characteristic form's coefficients of a degenerate elliptic linear PDO in  $\mathbb{R}^N$  and the regularity of vector fields controlling its degeneracy. We consider both the cases where the number of vector fields is  $N$  and it is equal to the rank.

### 1. INTRODUCTION

Let  $\Omega$  be an open subset of  $\mathbb{R}^N$ , and let  $\mathcal{L}$  be a second-order linear partial differential operator of the form

$$(1) \quad \mathcal{L} = \sum_{i,j=1}^N a_{ij}(x) \partial_{x_i} \partial_{x_j} + \sum_{k=1}^N b_k(x) \partial_{x_k}, \quad \text{for } x \in \Omega.$$

Let us just assume (for now) that, for all  $x$ , the matrix  $A(x) = (a_{ij}(x))_{i,j=1}^N$  is symmetric and nonnegative definite; that is,  $\mathcal{L}$  is degenerate-elliptic. It is well-known (see e.g. [6, 22, 38]) that the existence of vector fields with suitable properties allowing us to write  $\mathcal{L}$  as a sum of squares (possibly up to first order terms) can be crucial for studying qualitative and quantitative properties for the solutions or subsolutions to  $\mathcal{L}u = 0$ : the properties of the metric space related to such vector fields have been widely investigated and successfully exploited. Moreover, motivated by the studies on certain nonlinear degenerate-elliptic equations of sub-Riemannian type and on linear subelliptic equations with nonsmooth coefficients (see [1, 11, 12, 14, 42, 45] and the monographs [7, 8, 43], with the references therein), there have been recent investigations concerning the minimal regularity assumptions for having vector fields with some Hörmander-type properties [9, 10, 24, 31–36, 40]. For these reasons we think it is worthwhile to focus on conditions under which we can guarantee the existence in  $\Omega$  of vector fields with the desired regularity just looking at the quadratic form  $A(x)$ .

Before stating our results, let us be more precise about the PDE setting we are dealing with. In [39] Phillips and Sarason showed that if  $x \mapsto A(x)$  is a  $C^2$  map, then its (symmetric, nonnegative definite) square root  $S(x)$  has locally Lipschitz-continuous entries. Then we can identify the  $N$  columns of  $S$  with  $N$  locally

---

Received by the editors January 28, 2016, and, in revised form, June 3, 2017.

2010 *Mathematics Subject Classification*. Primary 35J70, 15A23, 47A56.

*Key words and phrases*. Degenerate-elliptic operators, nonsmooth vector fields, constant rank quadratic forms.

The third author is the corresponding author.

Lipschitz-continuous vector fields  $X_1, \dots, X_N$ . Up to lower order terms we can write  $\mathcal{L}$  as the sum of their squares. The fact they are locally Lipschitz allows us to consider their integral curves and to look at the Carnot-Carathéodory distance  $d_{CC}$ . Essential tools in order to trigger some PDE methods are the  $X$ -connectivity (which makes  $d_{CC}$  an actual distance), the doubling condition for the metric balls, and Poincaré inequalities. These properties provide Sobolev embeddings (see [17, 18, 20]) and the possibility to develop the De Giorgi-Moser technique for the Hölder regularity of the solutions. In this perspective there is the study of operators, even with possibly very rough coefficients, as the  $X$ -elliptic operators in the sense of [19, 27], where the existence of Lipschitz vector fields with doubling and Poincaré properties controlling the degeneracy are required axiomatically [19, 26, 27, 44, 46] (see the subsection below; see also [25, 27, 28] for a more general condition about an  $X$ -controllable almost exponential map). On the other hand, one can ask for which families of vector fields the connectivity/doubling/Poincaré properties are satisfied. It is very well-known that such properties hold true for smooth vector fields satisfying the Hörmander condition thanks to the milestones [23, 37]. In the case of nonsmooth vector fields, it is known for vector fields in specific diagonal forms since the pioneering works [15, 16]; see also the results and the references in [43]. For nonsmooth vector fields in general form there has been a renewed interest, as we mentioned, in step- $r$  Hörmander-type conditions with minimal regularity assumptions (such as  $C^{r-1,1}$  or peculiar intrinsic regularities) in order to have doubling conditions [10, 33, 36] and Poincaré inequalities [4, 10, 13, 28, 30, 33, 36].

Let us denote, for any  $x \in \Omega$ , by  $X_1(x), \dots, X_N(x)$  the vector fields identified with the columns of the square root  $S(x)$  of  $A(x)$ . In general it is not possible to improve the Phillips-Sarason result either for the Lipschitz outcome or for the  $C^2$ -assumption, as we can trivially show respectively with the examples

$$A(x) = \begin{pmatrix} 1 & 0 \\ 0 & x^2 \end{pmatrix} \quad \text{or} \quad A(x) = \begin{pmatrix} 1 & 0 \\ 0 & |x|^{2-\epsilon} \end{pmatrix}.$$

Nevertheless, in the first section of his celebrated paper [6] Bony quickly stated that it is possible to have smooth vector fields as soon as we have a smooth  $A(x)$  with constant rank. This is our starting point. Thus, throughout the paper we assume that

$$(2) \quad \text{there exists } m \in \{1, \dots, N\} \text{ such that} \\ \text{rk}(A(x)) \equiv m \quad \text{for all } x \in \Omega.$$

Under this constant rank condition we can prove the following:

**Theorem 1.1.** *The vector fields  $X_1, \dots, X_N$  have the same regularity as  $A$ ; that is, if  $A$  satisfies (2) and  $A \in C^{k,\alpha}(\Omega)$  for some  $\alpha \in [0, 1]$  and  $k \in \mathbb{N} \cup \{0, \infty, \omega\}$ , then every  $X_j \in C^{k,\alpha}(\Omega)$ .*

In our hypotheses, the  $N$  vectors  $X_1(x), \dots, X_N(x)$  span at any  $x$  an  $m$ -dimensional subspace of  $\mathbb{R}^N$ , and they are thus linearly dependent (in the meaningful case  $m < N$ ). In the applications it might be useful to have  $m$  linearly independent vector fields and to have information about their regularity. We can consider, at any fixed  $x$ , the decomposition

$$A(x) = R(x)R^t(x),$$

for some  $N \times m$  matrix  $R(x)$  with maximal rank. We may denote by  $\tilde{X}_1(x), \dots, \tilde{X}_m(x)$  the linearly independent vector fields identified with the  $m$  columns of  $R(x)$ . The decomposition (and hence the possible choice for the vector fields at any  $x$ ) is, of course, not unique. We can prove the following:

**Theorem 1.2.** *Let  $\Omega$  be a contractible open subset of  $\mathbb{R}^N$ . Suppose  $A$  satisfies (2) and  $A \in C^k(\Omega)$  for some  $k \in \mathbb{N} \cup \{0, \infty\}$ . Then, there exist  $m$  linearly independent vector fields  $\tilde{X}_1, \dots, \tilde{X}_m \in C^k(\Omega)$  such that  $A(x) = R(x)R^t(x)$  for all  $x \in \Omega$ .*

Therefore, in the case  $k \geq 1$ , we can write  $\mathcal{L}$  as the sum of  $\tilde{X}_j^2$  up to adjusting the first order terms. Let us explicitly remark that the reason for the  $C^k$ -regularity and not for the  $C^{k,\alpha}$ -regularity in Theorem 1.2 is technical: in our arguments we exploit a differential topology tool for which we have found a clear reference just in the  $C^k$ -smoothness category; as long as it holds true also with Lipschitz/Hölder/ $C^\omega$  regularity, Theorem 1.2 applies even in those situations.

The plan of the paper is as follows. In Section 2 we prove Theorem 1.1, which will readily follow from a simple differential geometry lemma. In Section 3 we finally prove Theorem 1.2. Before doing this, let us add here some further comments on our results.

**1.1. Comments and consequences.** In this subsection we denote, with abuse of notation, by  $X_1, \dots, X_r$  the vector fields whose existence and regularity in  $\Omega$  is ensured by Theorem 1.1 ( $r = N$ ) or by Theorem 1.2 ( $r = m$ , with  $\Omega$  contractible) under the constant rank assumption (2). Let us write them as  $X_j = \sum_{i=1}^N \sigma_{ij}(x)\partial_{x_i}$  (the  $N \times r$  matrix  $\sigma(x)$  is  $S(x)$  or  $R(x)$ , respectively).

We would like at first to give the details of the statement we already mentioned by Bony in [6, p. 279], since it was the initial motivation for the present study. Suppose we have a second order operator  $\mathcal{L}$  as in (1), where the nonnegative matrix  $A(x)$  and the first order terms  $(b_h(x))_{h=1}^N$  are  $C^\infty$ -smooth functions. Then our vector fields  $X_1, \dots, X_r$  are  $C^\infty$ -smooth. We have

$$(3) \quad \mathcal{L} = \sum_{j=1}^r X_j^2 + X_0 \quad \text{in } \Omega,$$

where the  $C^\infty$ -smooth vector field  $X_0 = \sum_{h=1}^N \left( b_h(x) - \sum_{j=1}^r X_j(\sigma_{hj})(x) \right) \partial_{x_h}$ . Thus, under our assumptions, it is completely equivalent to test the Hörmander condition on  $X_1, \dots, X_r$  or on the vector fields  $A_1, \dots, A_N$  identified with the columns of  $A$ . In fact, at any  $x \in \Omega$ , we have

$$(4) \quad \text{Lie} \{A_1, \dots, A_N\}(x) = \text{Lie} \{X_1, \dots, X_r\}(x) \quad \text{and}$$

$$(5) \quad \text{Lie} \{A_1, \dots, A_N, B\}(x) = \text{Lie} \{X_1, \dots, X_r, X_0\}(x),$$

where  $B(x) = \sum_{h=1}^N (b_h(x) - \sum_{j=1}^r \partial_{x_j}(a_{jh})(x)) \partial_{x_h}$ . As is well-known, the fact that such vector spaces coincide for every  $x$  with the whole  $\mathbb{R}^N$  is a sufficient condition for the hypoellipticity of  $\mathcal{L}$  in  $\Omega$ . In case of  $C^\omega$ -coefficients, the condition coming from (5) is even necessary for the hypoellipticity [38] (see also [3, 5] for discussions and applications concerning connectivity properties for  $C^\infty$ -hypoellipticity).

Since we have been careful with the regularity properties, we can be more precise. Suppose the nonnegative matrix  $A$  to be  $C^k$ , and the first order terms  $(b_h)_{h=1}^N \in C^{k-1}$ , for some  $\mathbb{N} \ni k \geq 1$ . Then we have (3) with  $X_1, \dots, X_r \in C^k$ ,

and  $X_0 \in C^{k-1}$ . Moreover, if we define for  $s \in \mathbb{N}$  the vector space  $\text{Lie}_s\{Y_j\} = \text{span}\{Y_j, [Y_{j_1}, Y_{j_2}], \dots, [Y_{j_1}, [Y_{j_2}, [\dots, [Y_{j_{s-1}}, Y_{j_s}], \dots]]\}$ , for any  $x \in \Omega$  we have

$$(6) \quad \text{Lie}_s\{A_1, \dots, A_N\}(x) = \text{Lie}_s\{X_1, \dots, X_r\}(x) \quad \text{for all } s \leq k + 1 \quad \text{and}$$

$$(7) \quad \text{Lie}_s\{A_1, \dots, A_N, B\}(x) = \text{Lie}_s\{X_1, \dots, X_r, X_0\}(x) \quad \text{for all } s \leq k.$$

*Remark 1.3.* For the reader’s convenience we provide here an outline of the proof of (6) and (7). This will give a fortiori a proof of (4) and (5).

In the notation we are using, for any  $j \in \{1, \dots, N\}$  and  $l \in \{1, \dots, r\}$ , we have  $A_j(x) = \sum_{i=1}^N a_{ij}(x)\partial_{x_i}$ ,  $X_l(x) = \sum_{i=1}^N \sigma_{il}(x)\partial_{x_i}$ . We recall that the existence of the  $C^k$ -matrix valued function  $\sigma(\cdot)$  globally defined in  $\Omega$  with rank  $m$  comes from our results. Since we have  $A(x) = \sigma(x)\sigma^t(x)$ , we can write  $A_j(x) = \sum_{l=1}^r \sigma_{jl}(x)X_l(x)$  for every  $j$ . This yields in particular (6) with  $s = 1$ , which reads as

$$\text{span}\{A_1(x), \dots, A_N(x)\} = \text{span}\{X_1(x), \dots, X_r(x)\} \quad \text{for all } x,$$

since both vector spaces are  $m$ -dimensional. Let us now prove (6) with  $s = 2$ , which means we have to prove for all  $x \in \Omega$  that

$$\text{span}\{A_j(x), [A_j, A_h](x), j, h = 1, \dots, N\} = \text{span}\{X_l(x), [X_l, X_q](x), l, q = 1, \dots, r\}.$$

The inclusion  $\subseteq$  is trivial since

$$\begin{aligned} [A_j, A_h](x) &= \sum_{l,q=1}^r \sigma_{jl}(x)\sigma_{hq}(x)[X_l, X_q](x) + \sum_{l,q=1}^r \sigma_{jl}(x)X_l(\sigma_{hq})(x)X_q(x) \\ &\quad - \sum_{l,q=1}^r \sigma_{hq}(x)X_q(\sigma_{jl})(x)X_l(x) \quad \text{for all } x \in \Omega. \end{aligned}$$

To deal with the opposite inclusion, we can fix any  $x_0 \in \Omega$  and find an open neighborhood of  $x_0$  where  $\text{span}\{A_1(x), \dots, A_N(x)\} = \text{span}\{A_{j_1}(x), \dots, A_{j_m}(x)\}$  and  $\text{span}\{X_1(x), \dots, X_r(x)\} = \text{span}\{X_{l_1}(x), \dots, X_{l_m}(x)\}$  (recall that  $r$  can be either  $m$  or  $N$ ). Thus, in such a neighborhood, we can write  $A_{j_p}(x) = \sum_{i=1}^m \hat{\sigma}_{pi}(x)X_{l_i}(x)$  for all  $p \in \{1, \dots, m\}$  for some  $\hat{\sigma}(\cdot) \in C^k$ . By construction  $\hat{\sigma}(x)$  is invertible and the inverse has to be  $C^k$ . Hence, we have an open neighborhood  $U_{x_0} \subset \Omega$  of  $x_0$  such that  $X_{l_i}(x) = \sum_{p=1}^m c_{ip}(x)A_{j_p}(x)$  for all  $i \in \{1, \dots, m\}$  and  $x \in U_{x_0}$ , where  $c(\cdot) \in C^k(U_{x_0})$ . This is enough to prove that  $\text{Lie}_2\{X_l\}(x) = \text{Lie}_2\{X_{l_i}\}(x) \subseteq \text{Lie}_2\{A_{j_p}\}(x) \subseteq \text{Lie}_2\{A_j\}(x)$  for all  $x \in U_{x_0}$ . The arbitrariness of  $x_0$  completes the proof of the ( $s = 2$ )-case in (6). We can proceed in the same way for higher  $s$ , until the coefficients are differentiable (that is,  $s \leq k + 1$ ). The proof of (7) is completely analogous since  $X_0(x) = B(x) + \sum_{j=1}^N \sum_{l=1}^r \partial_{x_j}(\sigma_{jl})(x)X_l(x)$  for all  $x \in \Omega$ .

The identifications in (6) and (7) might be useful (in the operative sense) in order to check quantitative properties of the distances related to the vector fields under the assumptions of Hörmander condition or involutive properties. The quantitative properties we are referring to are related to the analysis of the exponential map and almost-exponential maps carried out in [10, 33–35, 37].

Finally we would like to mention that, under our constant rank assumption, every operator  $\mathcal{L}$  in (1) can be seen as an  $X$ -elliptic operator as long as we can control the first order term given by  $X_0$ . Let us clarify this point while recalling the definition in [19]. Suppose  $A$  to be  $C^1$  and the first order terms  $b_k$ ’s to be

measurable. We can write, besides (3), the operator as  $\mathcal{L} = \operatorname{div}(A\nabla) + B$ . By the definition of our vector fields, we get

$$\langle A(x)\xi, \xi \rangle = \|\sigma^t(x)\xi\|^2 = \sum_{j=1}^r \langle X_j(x), \xi \rangle^2 \quad \forall \xi \in \mathbb{R}^N \text{ and } \forall x \in \Omega,$$

which gives at one time the conditions (1.2) and (1.5) in [19]. If we want  $\mathcal{L}$  to be  $X$ -elliptic in  $\Omega$ , we are thus left to satisfy the condition (1.4) in [19]; i.e. there has to exist a measurable function  $\gamma$  such that  $\langle B(x), \xi \rangle^2 \leq \gamma^2(x) \sum_{j=1}^r \langle X_j(x), \xi \rangle^2$ . Comparing  $B(x)$  and  $X_0(x)$  we have

$$\langle B(x) - X_0(x), \xi \rangle^2 \leq rN^2 \max_{jl} |\partial_{x_j}(\sigma_{jl})(x)|^2 \sum_{j=1}^r \langle X_j(x), \xi \rangle^2.$$

Hence, according to the definition in [19],  $\mathcal{L}$  is uniformly  $X$ -elliptic in  $\Omega$  if there exists a function  $\gamma_0$  such that

$$\langle X_0(x), \xi \rangle^2 \leq \gamma_0^2(x) \sum_{j=1}^r \langle X_j(x), \xi \rangle^2 \quad \forall \xi \in \mathbb{R}^N \text{ and } \forall x \in \Omega.$$

The results proved in [19, 46] for such operators require the metric, doubling, and Poincaré properties we mentioned in the Introduction, as well as some integrability conditions for the function  $\gamma$ . In the case of the vector fields considered in Theorem 1.1, we can assume  $A$  to be locally Lipschitz in  $\Omega$  and the conditions we have just seen regarding the  $X$ -ellipticity are meant to be satisfied almost everywhere.

## 2. THE SQUARED CASE

Theorem 1.1 concerns the regularity of the square root  $S(x)$  of  $A(x)$ . The square root of a nonnegative definite matrix is easily defined via diagonalization. One could think, in a naive way, to look separately at the regularity of the eigenvalues and of the eigenvectors as functions of  $x$ . Unfortunately, there are well-known examples of smooth matrices for which it is not possible to find an even continuous choice of eigenvectors (see e.g. [41]). In this direction, let us give here the example (often used in the PDE-setting we deal with) of the Heisenberg Laplacian in  $\mathbb{R}^3$  for which

$$(8) \quad A(x_1, x_2, x_3) = \begin{pmatrix} 1 & 0 & x_2 \\ 0 & 1 & -x_1 \\ x_2 & -x_1 & x_1^2 + x_2^2 \end{pmatrix}.$$

We note that, for such  $A$ , the condition (2) is satisfied with  $m = 2$ : in fact the eigenvalues of  $A(x_1, x_2, x_3)$  are  $1 + x_1^2 + x_2^2, 1, 0$ . For  $x_1^2 + x_2^2 \neq 0$ , the unit eigenvectors (up to a sign choice) are respectively

$$v_{1+r}(x) = \frac{(x_2, -x_1, x_1^2 + x_2^2)}{\sqrt{x_1^2 + x_2^2} \sqrt{1 + x_1^2 + x_2^2}}, \quad v_1(x) = \frac{(x_1, x_2, 0)}{\sqrt{x_1^2 + x_2^2}}, \quad \text{and} \quad v_0(x) = \frac{(-x_2, x_1, 1)}{\sqrt{1 + x_1^2 + x_2^2}}.$$

Thus, for any choice of an orthonormal basis of eigenvectors at a point  $(0, 0, x_3)$ ,  $v_{1+r}$  and  $v_1$  cannot be continuous at that point. Nonetheless, denoting by  $\Lambda$  and  $V$  the eigenvalue and eigenvector matrices, one can check that the square root  $S(x) = V(x)\sqrt{\Lambda(x)}V^t(x)$  is real-analytic in  $\mathbb{R}^3$  (see (10)), as it is also a consequence of our result. In order to give a proof of Theorem 1.1, let us fix some notation.

Let us fix the natural numbers  $N \geq m \geq 1$ . Denoting by  $\text{Sym}_N(\mathbb{R})$  the real vector space of all  $N \times N$  symmetric matrices with real coefficients, let us put

$$S_N^+(m) = \{A \in \text{Sym}_N(\mathbb{R}) : A \geq 0 \text{ and } \text{rk}(A) = m\}.$$

Let us consider the smooth (real-analytic) map

$$q : S_N^+(m) \rightarrow S_N^+(m), \quad q(M) = M^2.$$

The square root function is, of course, the global inverse of  $q$ . In our notation, for any  $x \in \Omega$ , the vector fields  $X_1(x), \dots, X_N(x)$  are identified with the columns of  $q^{-1}(A(x))$ . The key assumption (2) ensures the well-position of this composition, for all  $x \in \Omega$ . The proof of Theorem 1.1 will thus be a straight consequence of the following:

**Lemma 2.1.** *The function  $q$  is a  $C^\omega$ -diffeomorphism.*

Let us first recall that  $S_N^+(m)$  is an embedded smooth submanifold of  $\text{Sym}_N(\mathbb{R}) \cong \mathbb{R}^{N(N+1)/2}$ , with dimension  $d_N(m) = Nm - \frac{m(m-1)}{2}$ . As a matter of fact, the  $C^\omega$ -differentiable structure of  $S_N^+(m)$  locally around a matrix  $M_0$  (with a nonvanishing  $m \times m$  minor) is basically given by being the zero set of the minors of order  $(m + 1)$  (in the case  $m < N$ ; if  $m = N$ , it is just an open set of  $\text{Sym}_N(\mathbb{R})$ ).

*Proof of Lemma 2.1.* By implicit function theorem, it is enough to show that the differential of  $q$  at any point  $M \in S_N^+(m)$  is an isomorphism between  $T_M(S_N^+(m))$  and  $T_{M^2}(S_N^+(m))$ . Without loss of generality we can consider  $M$  in the diagonal form  $\text{diag}(\lambda_1, \dots, \lambda_m, 0, \dots, 0)$ , for positive  $\lambda_1, \dots, \lambda_m$ . For such diagonal matrices it is easy to describe the tangent space. The map

$$\Psi : \text{Sym}_N(\mathbb{R}) \rightarrow \mathbb{R}^{\frac{N(N+1)}{2} - d_N(m)}, \quad \Psi(A) = \left( \det(a_{ij})_{\substack{i \in \{1, \dots, m\} \cup \{p\} \\ j \in \{1, \dots, m\} \cup \{q\}}} \right)_{\substack{p, q = m+1, \dots, N \\ p \leq q}}$$

can be used as a local defining function for  $S_N^+(m)$  around  $M$ . Hence, one has

$$(9) \quad T_M(S_N^+(m)) = \ker(d_M\Psi) = \{H \in \text{Sym}_N(\mathbb{R}) : h_{ij} = 0 \text{ for all } i, j = m + 1, \dots, N\}.$$

For all  $H \in T_M(S_N^+(m))$ , we have

$$d_Mq(H) = \frac{d}{dt}\Big|_{t=0} (M + tH)^2 = MH + HM.$$

In order to show that  $d_Mq$  is an isomorphism, we may just prove the injectivity. If  $d_Mq(H) = 0$  for some  $H$ , then for every  $j = 1, \dots, N$  we have

$$0 = d_Mq(H) e_j = M(He_j) + H(Me_j).$$

When  $1 \leq j \leq m$ , from the above identity we infer that  $He_j$  is an eigenvector for  $M$  with corresponding eigenvalue  $(-\lambda_j)$ : since  $M$  is nonnegative definite, this implies that  $He_j = 0$  for all  $j = 1, \dots, m$ . On the other hand, if  $j \in \{m + 1, \dots, N\}$ , we get that  $He_j$  is in the kernel of  $M$  which is spanned by  $e_{m+1}, \dots, e_N$ . Since  $H \in T_M(S_N^+(m))$  and recalling (9), we immediately deduce that  $He_j = 0$  even for all  $j = m + 1, \dots, N$ . This proves that  $H = 0$  and the lemma.  $\square$

After having proved Theorem 1.1, let us give here a slight generalization of the last lemma. As a matter of fact, all the  $p$ th-powers of a matrix give a  $C^\omega$ -diffeomorphism if restricted to  $S_N^+(m)$ . Even for the sake of more generalization, we

could consider a real-analytic function  $f : \mathbb{R} \rightarrow \mathbb{R}$  satisfying the following properties:

$f(0) = 0$ ,  $f \not\equiv 0$ , and the derivatives of  $f$  of every order are nonnegative.

Under these assumptions, Pringsheim’s theorem ensures that the Maclaurin series of  $f$  converges to  $f$  itself for all  $x \in \mathbb{R}$ , that is,

$$f(x) = \sum_{n=1}^{\infty} \frac{f^{(n)}(0)}{n!} x^n, \quad \text{for all } x \in \mathbb{R}.$$

Moreover, we have that  $f$  is (strictly) monotonically increasing on  $I := [0, +\infty[$ , and it restricts to a bijection from  $I$  to itself. Let us now define the real-analytic map

$$q_f : S_N^+(m) \rightarrow S_N^+(m), \quad q_f(M) = \sum_{n=1}^{\infty} \frac{f^{(n)}(0)}{n!} M^n.$$

If  $M \in S_N^+(m)$  and  $P$  is an orthogonal  $N \times N$  matrix such that  $P^t M P$  is the diagonal matrix  $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_m, 0, \dots, 0)$ , one can write  $q_f(A)$  as

$$q_f(M) = P q_f(\Lambda) P^t = P \text{diag}(f(\lambda_1), \dots, f(\lambda_m), 0, \dots, 0) P^t.$$

It is then easy to recognize that  $q_f$  is globally invertible (for all  $m = 1, \dots, N$ ).

**Lemma 2.2.** *The map  $q_f$  is a real-analytic diffeomorphism.*

*Proof.* As in the proof of Lemma 2.1, we have to show that the differential of  $q_f$  at any point  $M \in S_N^+(m)$  is an isomorphism between  $T_M(S_N^+(m))$  and  $T_{q_f(M)}(S_N^+(m))$ : without loss of generality we can assume that  $M = \text{diag}(\lambda_1, \dots, \lambda_m, 0, \dots, 0)$ , for positive  $\lambda_1, \dots, \lambda_m$ . For  $H \in T_M(S_N^+(m))$  we have

$$\begin{aligned} d_M(q_f)(H) &= \frac{d}{dt} \Big|_{t=0} \sum_{n=1}^{\infty} \frac{f^{(n)}(0)}{n!} (M + tH)^n \\ &= \sum_{n=1}^{\infty} \frac{f^{(n)}(0)}{n!} (M^{n-1}H + M^{n-2}HM + \dots + HM^{n-1}), \end{aligned}$$

by the uniform convergence of the power series. Let us prove the injectivity of  $d_M(q_f)$ . To this aim, let  $d_M(q_f)(H) = 0$  for some  $H \in T_M(S_N^+(m))$ . For  $j \in \{1, \dots, m\}$ , we have

$$0 = d_M(q_f)(H)e_j = \sum_{n=1}^{\infty} \frac{f^{(n)}(0)}{n!} \left[ \left( \sum_{k=0}^{n-2} \lambda_j^k M^{n-1-k} \right) \cdot (He_j) + \lambda_j^{n-1} \cdot (He_j) \right].$$

If we now define, for all  $n \in \mathbb{N}$ ,  $B_n := \sum_{k=0}^{n-2} \lambda_j^k M^{n-1-k} \in S_N^+(m)$ , then it is immediate to see that the series  $\sum_{n=1}^{\infty} \frac{f^{(n)}(0)}{n!} B_n$  is absolutely convergent on the Banach space  $\text{Sym}_N(\mathbb{R})$ , and thus it converges to a matrix  $B_0$  which is actually (symmetric and) nonnegative definite. We can then write the above equality in the following way:

$$0 = B_0(He_j) + \left( \sum_{n=1}^{\infty} \frac{f^{(n)}(0)}{n!} \lambda_j^{n-1} \right) (He_j) = B_0(He_j) + \frac{f(\lambda_j)}{\lambda_j} (He_j).$$

It follows that  $He_j$  is an eigenvector of  $B_0 \geq 0$  with eigenvalue  $(-f(\lambda_j)/\lambda_j) < 0$ , and thus  $He_j = 0$  for all  $j = 1, \dots, m$ . On the other hand, for  $j \in \{m + 1, \dots, N\}$ ,

the vector  $e_j$  is in the kernel of  $M$ . By setting

$$M_0 := \sum_{n=1}^{\infty} \frac{f^{(n)}(0)}{n!} M^{n-1} = \text{diag}(f(\lambda_1)/\lambda_1, \dots, f(\lambda_m)/\lambda_m, f'(0), \dots, f'(0)),$$

we get that  $0 = d_M(q_f)(H)e_j = M_0(He_j)$ ; i.e. the vector  $He_j$  is in the kernel of  $M_0$ . There are two cases. If  $f'(0) \neq 0$ , this gives immediately  $He_j = 0$  for all  $j = m+1, \dots, N$ . If  $f'(0) = 0$ , then the vectors  $He_{m+1}, \dots, He_N$  must be included in the span $\{e_{m+1}, \dots, e_N\}$ : since  $H \in T_M(S_N^+(m))$ , we obtain again  $He_j = 0$  for all  $j = m+1, \dots, N$ . In both cases we have  $H = 0$ , and the proof is complete.  $\square$

### 3. THE RECTANGULAR CASE

Let us go back to the example of the quadratic form  $A(x)$  in (8) for the Heisenberg Laplacian. In the previous section we dealt with its square root

(10)

$$S(x) = \begin{pmatrix} \frac{1}{x_1^2+x_2^2} \left( x_1^2 + \frac{x_2^2}{\sqrt{1+x_1^2+x_2^2}} \right) & \frac{x_1 x_2}{x_1^2+x_2^2} \left( 1 - \frac{1}{\sqrt{1+x_1^2+x_2^2}} \right) & \frac{x_2}{\sqrt{1+x_1^2+x_2^2}} \\ \frac{x_1 x_2}{x_1^2+x_2^2} \left( 1 - \frac{1}{\sqrt{1+x_1^2+x_2^2}} \right) & \frac{1}{x_1^2+x_2^2} \left( x_2^2 + \frac{x_1^2}{\sqrt{1+x_1^2+x_2^2}} \right) & -\frac{x_1}{\sqrt{1+x_1^2+x_2^2}} \\ \frac{x_2}{\sqrt{1+x_1^2+x_2^2}} & -\frac{x_1}{\sqrt{1+x_1^2+x_2^2}} & \frac{x_1^2+x_2^2}{\sqrt{1+x_1^2+x_2^2}} \end{pmatrix}.$$

The entries of  $S(x)$  have nothing to do with the usual Heisenberg vector fields  $\partial_{x_1} + x_2 \partial_{x_3}$ ,  $\partial_{x_2} - x_1 \partial_{x_3}$ , which are a couple of smooth and linearly independent vector fields defined globally in  $\mathbb{R}^3$ . In this section we thus want to deal with the existence and the regularity of a *global* decomposition  $A = RR^t$ , which in the Heisenberg example can be seen, for example, as

$$R(x) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ x_2 & -x_1 \end{pmatrix}.$$

Let us fix some notation. Fix two natural numbers  $N \geq m \geq 1$ . Denoting by  $M_{N \times m}(\mathbb{R})$  the vector space of the  $N \times m$  matrices with real coefficients, let us define

$$\Omega_N(m) = \{M \in M_{N \times m}(\mathbb{R}) : \text{rk}(M) = m\},$$

which is an open subset of  $M_{N \times m}(\mathbb{R}) \cong \mathbb{R}^{Nm}$ . Let us restate Theorem 1.2 in this notation.

**Theorem 1.2.** *Let  $\Omega \subseteq \mathbb{R}^N$  be a contractible open set and let  $A : \Omega \rightarrow S_N^+(m)$  be a map of class  $C^k$  (for some  $k \in \mathbb{N} \cup \{0, \infty\}$ ). Then there exists a map  $R : \Omega \rightarrow \Omega_N(m)$  of class  $C^k$  such that*

$$(11) \quad A(x) = R(x)R(x)^t \quad \text{for all } x \in \Omega.$$

In order to prove this theorem, we are going to exploit a smooth identification between  $S_N^+(m)$  and the quotient of  $M_{N \times m}(\mathbb{R})$  by the action of the orthogonal group. This is probably a known fact, but, since we want to use some specific properties of this isomorphism and of the quotient manifold, we will briefly show the construction for the sake of completeness. After that, we are going to invoke a homotopy lifting property for the related principal bundle, which will allow us to prove the desired theorem.



Let us denote by  $O(m)$  the group of the  $m \times m$  orthogonal matrices with real coefficients, which is a Lie group of dimension  $m(m - 1)/2$ . There is a standard real-analytic (right-)action of  $O(m)$  on the manifold  $\Omega_N(m)$ , that is,

$$\rho : \Omega_N(m) \times O(m) \rightarrow \Omega_N(m), \quad \rho(M, O) = MO.$$

It is not difficult to see that this action  $\rho$  is free and proper; i.e. (see also [29])

- if there exist  $M \in \Omega_N(m)$  and  $O \in O(m)$  such that  $MO = M$ , then  $O = \text{Id}_m$ ;
- the map  $\Omega_N(m) \times O(m) \ni (M, O) \mapsto (MO, M) \in \Omega_N(m) \times \Omega_N(m)$  is proper (by compactness of  $O(m)$ ).

Therefore the orbit space

$$V_N(m) = \Omega_N(m)/O(m)$$

is a topological manifold of dimension  $\dim(\Omega_N(m)) - \dim(O(m)) = d_N(m) = \dim(S_N^+(m))$ . Moreover it has a unique smoothly differentiable structure with the property that the quotient map  $\pi : \Omega_N(m) \rightarrow V_N(m)$  is a smooth submersion (see e.g. [29, Theorem 21.10]). Let us now consider the real-analytic map  $\Theta$  defined as follows:

$$\Theta : \Omega_N(m) \rightarrow S_N^+(m), \quad \Theta(M) = MM^t.$$

Since  $\Theta$  is constant on each orbit of the action  $\rho$ , it defines a smooth map  $\bar{\Theta}$  on the quotient  $V_N(m)$ :

$$\bar{\Theta} : V_N(m) \rightarrow S_N^+(m), \quad \bar{\Theta}([M]) = \Theta(M) = MM^t.$$

In other words,  $\bar{\Theta}$  is the unique (smooth) map such that the following diagram commutes:

$$\begin{array}{ccc} \Omega_N(m) & \xrightarrow{\quad \Theta \quad} & S_N^+(m) \\ \pi \downarrow & \nearrow \bar{\Theta} & \\ V_N(m) & & \end{array}$$

Since it is easy to see (e.g. via diagonalization) that  $\Theta$  is surjective, then  $\bar{\Theta}$  is surjective as well. Concerning the injectivity of  $\bar{\Theta}$ , it is a straightforward consequence of the following lemma.

**Lemma 3.1.** *Let  $M, N \in \Omega_N(m)$  with  $MM^t = NN^t$ . Then there exists  $O \in O(m)$  such that  $M = NO$ .*

*Proof.* Considering  $A = MM^t = NN^t \in S_N^+(m)$ , let  $\lambda_1, \dots, \lambda_m$  be its strictly positive eigenvalues. Let us take an orthonormal basis  $\{x_1, \dots, x_N\}$  of  $\mathbb{R}^N$  of eigenvectors of  $A$  such that  $Ax_i = \lambda_i x_i$  for  $i \in \{1, \dots, m\}$ . Let us define two more orthonormal bases  $\mathcal{B}_1 = \{u_1, \dots, u_m\}$  and  $\mathcal{B}_2 = \{v_1, \dots, v_m\}$  of  $\mathbb{R}^m$  as follows:

$$u_i = \frac{1}{\sqrt{\lambda_i}}(M^t x_i), \quad v_i = \frac{1}{\sqrt{\lambda_i}}(N^t x_i) \quad \text{for all } i = 1, \dots, m.$$

Let us make  $O$  the orthogonal  $m \times m$  matrix bringing  $\mathcal{B}_1$  into  $\mathcal{B}_2$ . For any  $i \in \{1, \dots, m\}$ , we get

$$Mu_i = \frac{1}{\sqrt{\lambda_i}}(MM^t)x_i = \frac{1}{\sqrt{\lambda_i}}(NN^t)x_i = Nv_i = N(Ou_i).$$

This ends the proof. □

Hence  $\bar{\Theta}$  is a smooth bijection from  $V_N(m)$  to  $S_N^+(m)$ . We want to see that it is actually a smooth diffeomorphism.

**Lemma 3.2.**  *$V_N(m)$  and  $S_N^+(m)$  are smoothly diffeomorphic via  $\bar{\Theta}$ .*

*Proof.* We have to show that the differential  $d_{[M]}\bar{\Theta}$  at any point  $[M] \in V_N(m)$  is nonsingular. Since the  $V_N(m)$  and  $S_N^+(m)$  have the same dimension, it is enough to show the surjectivity of  $d_{[M]}\bar{\Theta}$  which is equivalent ( $\pi$  being a submersion) to the surjectivity of  $d_M\Theta : T_M(\Omega_N(m)) \rightarrow T_{\Theta(M)}(S_N^+(m))$  for any  $M \in \Omega_N(m)$ .

We claim that, without loss of generality, we can just consider  $M \in \Omega_N(m)$  with  $\Theta(M) = \text{diag}(\lambda_1, \dots, \lambda_m, 0, \dots, 0)$ . Let us convince ourselves of this claim. Fix  $M_0 \in \Omega_N(m)$  and  $A = \Theta(M_0)$  and take an orthogonal  $N \times N$  matrix  $U$  such that

$$U^t A U = (U^t M_0) \cdot (U^t M_0)^t = \Lambda = \text{diag}(\lambda_1, \dots, \lambda_m, 0, \dots, 0),$$

for some positive  $\lambda_1, \dots, \lambda_m$ . Denote by  $L_U$  the automorphism of  $M_{N \times m}(\mathbb{R})$  defined by  $L_U(M) = U^t M$ , and by  $C_U$  the automorphism of  $\text{Sym}_N(\mathbb{R})$  defined by  $C_U(A) = U A U^t$ . For all  $M \in \Omega_N(m)$  we have by construction that  $(C_U \circ \Theta \circ L_U)(M) = \Theta(M)$ . We thus get

$$d_{M_0}\Theta = C_U \circ d_{(U^t M_0)}\Theta \circ L_U$$

on  $T_{M_0}(\Omega_N(m)) = M_{N \times m}(\mathbb{R})$ . By our choice of  $U$ , the matrix  $U^t M \in \Omega_N(m)$  is such that  $\Theta(U^t M) = \Lambda$ , and the claim is proved.

We can now fix  $M \in \Omega_N(m)$  with  $\Theta(M) = \text{diag}(\lambda_1, \dots, \lambda_m, 0, \dots, 0)$ , and look for the surjectivity of  $d_M\Theta$ . This assumption implies that  $M_{ij} = 0$  for all  $i = m + 1, \dots, N$  and  $j = 1, \dots, m$ . For every  $H \in M_{N \times m}(\mathbb{R}) \equiv T_M(\Omega_N(m))$ , we get

$$d_M\Theta(H) = \frac{d}{dt}\Big|_{t=0} \Theta(M + tH) = M \cdot H^t + H \cdot M^t.$$

If we denote, for fixed  $i \in \{1, \dots, N\}$  and  $j \in \{1, \dots, m\}$ , by  $E^{(ij)}$  the standard  $(i, j)$ -elementary matrix of the canonical basis of  $M_{N \times m}(\mathbb{R})$ , we have  $(M(E^{(ij)})^t)_{hk} = M_{hj}\delta_{ki}$  and  $(E^{(ij)}M^t)_{hk} = M_{kj}\delta_{hi}$ . We can thus verify that, for every  $i \in \{1, \dots, N\}$  and  $j \in \{1, \dots, m\}$ ,

$$d_M\Theta(E^{(ij)}) = (M(E^{(ij)})^t) + (E^{(ij)}M^t) = \sum_{l=1}^m M_{lj}(\tilde{E}^{(li)} + \tilde{E}^{(il)})$$

where, for  $r, s \in \{1, \dots, N\}$ ,  $\tilde{E}^{(rs)}$  denotes the standard  $(r, s)$ -elementary matrix of the canonical basis of  $M_N(\mathbb{R})$ . It is now quite easy to see that the image of  $d_M\Theta$ , which is generated by the  $Nm$  vectors  $d_M\Theta(E^{(11)}), \dots, d_M\Theta(E^{(Nm)})$ , spans the whole  $T_{\Theta(M)}(S_N^+(m))$  (by recalling (9)). The proof is complete.  $\square$

Therefore, we can think of the  $C^k$  map  $A$  in Theorem 1.2 as a map which takes value in the quotient manifold  $V_N(m)$ . Our aim is somehow to extract a map taking value in  $\Omega_N(m)$ . We can do that thanks to the structure of a principal  $O(m)$ -bundle. Let us recall (we also refer to e.g. [21, Chapter IV] and [47, Chapter II]) that a principal  $G$ -bundle is a (locally trivial) smooth fiber bundle with typical fiber a Lie group  $G$ , for which the action preserves the fibers and there is an equivariance property between the action and the bundle charts. Moreover, if  $M$  is a smooth manifold and  $G$  is a Lie group acting smoothly, freely, and properly on  $M$ , then the quadruple  $(M, M \rightarrow M/G, M/G, G)$  is a principal  $G$ -bundle (see, for example, [29, Exercise 21-6]). That is why the quadruple  $\mathcal{F}_m = (\Omega_N(m), \pi, V_N(m), O(m))$  is a principal  $O(m)$ -bundle.

The main ingredient for our purposes is the following differential topology tool for principal  $G$ -bundles:

let  $\mathcal{F} = (E, \mathfrak{p}, M, G)$  be a principal  $G$ -bundle and let  $X$  be a manifold of class  $C^k$ ; then  $\mathcal{F}$  satisfies the following *homotopy lifting property* with respect to  $X$ :

if  $f \in C^k(X; E)$  and  $H : X \times [0, 1] \rightarrow M$  is a map of class  $C^k(X \times [0, 1])$  such that  $H(\cdot, 0) = \mathfrak{p} \circ f$ , then there exists a map  $\tilde{H} : X \times [0, 1] \rightarrow E$  of class  $C^k(X \times [0, 1])$  satisfying  $H = \mathfrak{p} \circ \tilde{H}$ .

For a proof we refer the reader to [21, Chapter III, Theorem 2.4] (for a particular class of fiber bundles), [2, Chapter III], and [47, Chapter II, Corollary 6.1]. Roughly speaking, if we have a  $C^k$ -homotopy  $H : X \times [0, 1] \rightarrow M$  for which  $H(\cdot, 0)$  admits a  $C^k$ -lift to  $E$ , then the whole homotopy  $H$  admits a  $C^k$ -lift to  $E$ . We can now complete the proof.

*Proof of Theorem 1.2.* Let us set

$$\bar{A} : \Omega \rightarrow V_N(m), \quad \bar{A}(x) = \bar{\Theta}^{-1}(A(x)).$$

Since  $\Omega$  is contractible, we can find a point  $x_0 \in \Omega$  which is a deformation retract of  $\Omega$ ; i.e. there exists a smooth map  $r : \Omega \times [0, 1] \rightarrow \Omega$  such that  $r(x, 0) = x_0$  and  $r(x, 1) = x$  for all  $x \in \Omega$ . Let us now consider the homotopy

$$H : \Omega \times [0, 1] \rightarrow V_N(m), \quad H(x, t) = \bar{A}(r(x, t)).$$

Since  $r$  is smooth and  $\bar{A}$  is  $C^k$  in  $\Omega$ , we have  $H \in C^k(\Omega \times [0, 1]; V_N(m))$ . Moreover, being  $H(\cdot, 0)$  the constant map  $\bar{A}(x_0)$ , it trivially admits a lift of class  $C^k$  from  $\Omega$  to  $\Omega_N(m)$  (namely, the constant map  $\Omega \ni x \mapsto R_0 \in \Omega_N(m)$ , where  $R_0$  is some matrix in  $\Omega_N(m)$  such that  $[R] = \bar{A}(x_0)$ ). By the homotopy lifting property there exists  $\bar{H} : \Omega \times [0, 1] \rightarrow \Omega_N(m)$  of class  $C^k$  such that  $H = \pi \circ \bar{H}$ . We can then define a map  $R$  from  $\Omega$  to  $\Omega_N(m)$  in the following way:

$$R : \Omega \rightarrow \Omega_N(m), \quad R(x) = \bar{H}(x, 1).$$

By construction the map  $R$  is of class  $C^k$  on  $\Omega$ , that is, the same regularity as  $A$ . For every  $x \in \Omega$  we also have

$$R(x)R(x)^t = \Theta(R(x)) = \bar{\Theta}(\pi \circ \bar{H}(x, 1)) = \bar{\Theta}(H(x, 1)) = \bar{\Theta}(\bar{A}(x)) = A(x),$$

which is exactly the desired identity (11). □

Let us remark explicitly that the map  $R$  (and therefore the vector fields) we have constructed depends on  $A$  in a much *stronger* way with respect to the square-root construction of Section 2. More precisely, we cannot hope to find a smooth map  $\sigma : V_N(m) \rightarrow \Omega_N(m)$  (independent of  $A$ ) in order to define  $R = \sigma(\bar{\Theta}^{-1}(A))$ . As a matter of fact, if such a  $\sigma$  exists, then it would be a smooth global section of our bundle  $\mathcal{F}_m$ . Since  $\mathcal{F}_m$  is a principal  $O(m)$ -bundle, it should be globally trivial (see e.g. [47, Theorem 4.2]). This means, in particular, that  $\Omega_N(m)$  should be diffeomorphic to the product manifold  $V_N(m) \times O(m)$ , which is a contradiction (if  $m < N$  the space  $\Omega_N(m)$  is indeed connected, whereas  $V_N(m) \times O(m)$  has at least two connected components). The case  $m = N$  is slightly different. In fact, the polar decomposition provides a global trivialization of the principal bundle  $\mathcal{F}_N = (\text{GL}_N(\mathbb{R}), \pi, V_N(N), O(N))$ , and the map  $\sqrt{\cdot} : V_N(N) \equiv S_N^+(N) \rightarrow \text{GL}_N(\mathbb{R})$  defines a smooth global section.

## REFERENCES

- [1] F. Abedin, C. E. Gutiérrez, and G. Tralli, *Harnack's inequality for a class of non-divergent equations in the Heisenberg group*, Comm Partial Differential Equations, to appear. DOI: 10.1080/03605302.2017.1384836.
- [2] R. Abraham, J. E. Marsden, and T. Ratiu, *Manifolds, tensor analysis, and applications*, 2nd ed., Applied Mathematical Sciences, vol. 75, Springer-Verlag, New York, 1988. MR960687
- [3] Kazuo Amano, *A necessary condition for hypoellipticity of degenerate elliptic-parabolic operators*, Tokyo J. Math. **2** (1979), no. 1, 111–120, DOI 10.3836/tjm/1270473562. MR541900
- [4] S. G. Basalaev, *The Poincaré inequality for  $C^{1,\alpha}$ -smooth vector fields* (Russian, with Russian summary), Sibirsk. Mat. Zh. **55** (2014), no. 2, 267–284; English transl., Sib. Math. J. **55** (2014), no. 2, 215–229. MR3237332
- [5] Erika Battaglia, Stefano Biagi, and Andrea Bonfiglioli, *The strong maximum principle and the Harnack inequality for a class of hypoelliptic non-Hörmander operators* (English, with English and French summaries), Ann. Inst. Fourier (Grenoble) **66** (2016), no. 2, 589–631. MR3477885
- [6] Jean-Michel Bony, *Principe du maximum, inégalité de Harnack et unicité du problème de Cauchy pour les opérateurs elliptiques dégénérés* (French, with English summary), Ann. Inst. Fourier (Grenoble) **19** (1969), no. fasc. 1, 277–304 xii. MR0262881
- [7] Marco Bramanti, Luca Brandolini, Ermanno Lanconelli, and Francesco Uguzzoni, *Non-divergence equations structured on Hörmander vector fields: heat kernels and Harnack inequalities*, Mem. Amer. Math. Soc. **204** (2010), no. 961, vi+123, DOI 10.1090/S0065-9266-09-00605-X. MR2604962
- [8] M. Bramanti, L. Brandolini, M. Manfredini, and M. Pedroni, *Fundamental solutions and local solvability for nonsmooth Hörmander's operators*, Mem. Amer. Math. Soc., **249** (2017), DOI 10.1090/memo/1182.
- [9] Marco Bramanti, Luca Brandolini, and Marco Pedroni, *On the lifting and approximation theorem for nonsmooth vector fields*, Indiana Univ. Math. J. **59** (2010), no. 6, 2093–2138, DOI 10.1512/iumj.2010.59.4298. MR2919750
- [10] Marco Bramanti, Luca Brandolini, and Marco Pedroni, *Basic properties of nonsmooth Hörmander's vector fields and Poincaré's inequality*, Forum Math. **25** (2013), no. 4, 703–769, DOI 10.1515/form.2011.133. MR3089748
- [11] Luca Capogna, Giovanna Citti, and Maria Manfredini, *Smoothness of Lipschitz minimal intrinsic graphs in Heisenberg groups  $\mathbb{H}^n$ ,  $n > 1$* , J. Reine Angew. Math. **648** (2010), 75–110, DOI 10.1515/CRELLE.2010.080. MR2774306
- [12] G. Citti, E. Lanconelli, and A. Montanari, *Smoothness of Lipschitz-continuous graphs with nonvanishing Levi curvature*, Acta Math. **188** (2002), no. 1, 87–128, DOI 10.1007/BF02392796. MR1947459
- [13] Giovanna Citti, Maria Manfredini, Andrea Pinamonti, and Francesco Serra Cassano, *Poincaré-type inequality for Lipschitz continuous vector fields* (English, with English and French summaries), J. Math. Pures Appl. (9) **105** (2016), no. 3, 265–292, DOI 10.1016/j.matpur.2015.09.001. MR3465805
- [14] Giovanna Citti, Andrea Pascucci, and Sergio Polidoro, *On the regularity of solutions to a nonlinear ultraparabolic equation arising in mathematical finance*, Differential Integral Equations **14** (2001), no. 6, 701–738. MR1826957
- [15] Bruno Franchi and Ermanno Lanconelli, *Une métrique associée à une classe d'opérateurs elliptiques dégénérés* (French), Conference on linear partial and pseudodifferential operators (Torino, 1982), Rend. Sem. Mat. Univ. Politec. Torino **Special Issue** (1983), 105–114 (1984). Conference on linear partial and pseudodifferential operators (Torino, 1982). MR745979
- [16] Bruno Franchi and Ermanno Lanconelli, *Hölder regularity theorem for a class of linear nonuniformly elliptic operators with measurable coefficients*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **10** (1983), no. 4, 523–541. MR753153
- [17] Bruno Franchi, Raul Serapioni, and Francesco Serra Cassano, *Approximation and imbedding theorems for weighted Sobolev spaces associated with Lipschitz continuous vector fields* (English, with Italian summary), Boll. Un. Mat. Ital. B (7) **11** (1997), no. 1, 83–117. MR1448000
- [18] Nicola Garofalo and Duy-Minh Nhieu, *Isoperimetric and Sobolev inequalities for Carnot-Carathéodory spaces and the existence of minimal surfaces*, Comm. Pure Appl. Math.

- 49** (1996), no. 10, 1081–1144, DOI 10.1002/(SICI)1097-0312(199610)49:10<1081::AID-CPA3>3.0.CO;2-A. MR1404326
- [19] Cristian E. Gutiérrez and Ermanno Lanconelli, *Maximum principle, nonhomogeneous Harnack inequality, and Liouville theorems for  $X$ -elliptic operators*, Comm. Partial Differential Equations **28** (2003), no. 11–12, 1833–1862, DOI 10.1081/PDE-120025487. MR2015404
- [20] Piotr Hajlasz and Pekka Koskela, *Sobolev met Poincaré*, Mem. Amer. Math. Soc. **145** (2000), no. 688, x+101, DOI 10.1090/memo/0688. MR1683160
- [21] Morris W. Hirsch, *Differential topology*, Graduate Texts in Mathematics, No. 33, Springer-Verlag, New York-Heidelberg, 1976. MR0448362
- [22] Lars Hörmander, *Hypoelliptic second order differential equations*, Acta Math. **119** (1967), 147–171, DOI 10.1007/BF02392081. MR0222474
- [23] David Jerison, *The Poincaré inequality for vector fields satisfying Hörmander’s condition*, Duke Math. J. **53** (1986), no. 2, 503–523, DOI 10.1215/S0012-7094-86-05329-9. MR850547
- [24] Maria Karmanova and Sergey Vodop’yanov, *Geometry of Carnot-Carathéodory spaces, differentiability, coarea and area formulas*, Analysis and mathematical physics, Trends Math., Birkhäuser, Basel, 2009, pp. 233–335, DOI 10.1007/978-3-7643-9906-1\_14. MR2724617
- [25] Alessia E. Kogoj, *A control condition for a weak Harnack inequality*, Nonlinear Anal. **75** (2012), no. 11, 4198–4204, DOI 10.1016/j.na.2012.03.008. MR2921982
- [26] Alessia E. Kogoj and Stefanie Sonner, *Attractors met  $X$ -elliptic operators*, J. Math. Anal. Appl. **420** (2014), no. 1, 407–434, DOI 10.1016/j.jmaa.2014.05.070. MR3229832
- [27] Ermanno Lanconelli and Alessia Elisabetta Kogoj,  *$X$ -elliptic operators and  $X$ -control distances*, Contributions in honor of the memory of Ennio De Giorgi (Italian), Ricerche Mat. **49** (2000), no. suppl., 223–243. MR1826225
- [28] Ermanno Lanconelli and Daniele Morbidelli, *On the Poincaré inequality for vector fields*, Ark. Mat. **38** (2000), no. 2, 327–342, DOI 10.1007/BF02384323. MR1785405
- [29] John M. Lee, *Introduction to smooth manifolds*, Graduate Texts in Mathematics, vol. 218, Springer-Verlag, New York, 2003. MR1930091
- [30] Maria Manfredini, *A note on the Poincaré inequality for Lipschitz vector fields of step two*, Proc. Amer. Math. Soc. **138** (2010), no. 2, 567–575, DOI 10.1090/S0002-9939-09-10054-0. MR2557174
- [31] Maria Manfredini, *Fundamental solutions for sum of squares of vector fields operators with  $C^{1,\alpha}$  coefficients*, Forum Math. **24** (2012), no. 5, 973–1011, DOI 10.1515/form.2011.093. MR2988569
- [32] Annamaria Montanari and Daniele Morbidelli, *Balls defined by nonsmooth vector fields and the Poincaré inequality* (English, with English and French summaries), Ann. Inst. Fourier (Grenoble) **54** (2004), no. 2, 431–452. MR2073841
- [33] Annamaria Montanari and Daniele Morbidelli, *Nonsmooth Hörmander vector fields and their control balls*, Trans. Amer. Math. Soc. **364** (2012), no. 5, 2339–2375, DOI 10.1090/S0002-9947-2011-05395-X. MR2888209
- [34] Annamaria Montanari and Daniele Morbidelli, *Almost exponential maps and integrability results for a class of horizontally regular vector fields*, Potential Anal. **38** (2013), no. 2, 611–633, DOI 10.1007/s11118-012-9289-6. MR3015367
- [35] Annamaria Montanari and Daniele Morbidelli, *Step- $s$  involutive families of vector fields, their orbits and the Poincaré inequality* (English, with English and French summaries), J. Math. Pures Appl. (9) **99** (2013), no. 4, 375–394, DOI 10.1016/j.matpur.2012.09.005. MR3035947
- [36] Annamaria Montanari and Daniele Morbidelli, *Generalized Jacobi identities and ball-box theorem for horizontally regular vector fields*, J. Geom. Anal. **24** (2014), no. 2, 687–720, DOI 10.1007/s12220-012-9351-z. MR3192293
- [37] Alexander Nagel, Elias M. Stein, and Stephen Wainger, *Balls and metrics defined by vector fields. I. Basic properties*, Acta Math. **155** (1985), no. 1–2, 103–147, DOI 10.1007/BF02392539. MR793239
- [38] O. A. Oleĭnik and E. V. Radkevič, *Second order equations with nonnegative characteristic form*, translated from the Russian by Paul C. Fife, Plenum Press, New York-London, 1973. MR0457908
- [39] R. S. Phillips and Leonard Sarason, *Elliptic-parabolic equations of the second order*, J. Math. Mech. **17** (1967/1968), 891–917. MR0219868

- [40] Franco Rampazzo and Héctor J. Sussmann, *Commutators of flow maps of nonsmooth vector fields*, J. Differential Equations **232** (2007), no. 1, 134–175, DOI 10.1016/j.jde.2006.04.016. MR2281192
- [41] Franz Rellich, *Perturbation theory of eigenvalue problems*, assisted by J. Berkowitz, with a preface by Jacob T. Schwartz, Gordon and Breach Science Publishers, New York-London-Paris, 1969. MR0240668
- [42] Cristian Rios, Eric T. Sawyer, and Richard L. Wheeden, *Regularity of subelliptic Monge-Ampère equations*, Adv. Math. **217** (2008), no. 3, 967–1026, DOI 10.1016/j.aim.2007.07.004. MR2383892
- [43] Eric T. Sawyer and Richard L. Wheeden, *Hölder continuity of weak solutions to subelliptic equations with rough coefficients*, Mem. Amer. Math. Soc. **180** (2006), no. 847, x+157, DOI 10.1090/memo/0847. MR2204824
- [44] Giulio Tralli and Francesco Uguzzoni, *Wiener criterion for  $X$ -elliptic operators*, J. Differential Equations **259** (2015), no. 11, 6510–6527, DOI 10.1016/j.jde.2015.07.031. MR3397330
- [45] G. Tralli and F. Uguzzoni, *On a non-smooth potential analysis for Hörmander-type operators*, submitted.
- [46] Francesco Uguzzoni, *Estimates of the Green function for  $X$ -elliptic operators*, Math. Ann. **361** (2015), no. 1-2, 169–190, DOI 10.1007/s00208-014-1072-0. MR3302617
- [47] Gerard Walschap, *Metric structures in differential geometry*, Graduate Texts in Mathematics, vol. 224, Springer-Verlag, New York, 2004. MR2045823

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DEGLI STUDI DI BOLOGNA, PIAZZA DI PORTA S. DONATO, 5 - 40126 BOLOGNA, ITALY

*Current address:* Dipartimento di Matematica “Tullio Levi-Civita”, Università degli Studi di Padova, Via Trieste 63, 35121 Padova, Italy

*E-mail address:* [erika.battaglia@math.unipd.it](mailto:erika.battaglia@math.unipd.it)

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DEGLI STUDI DI BOLOGNA, PIAZZA DI PORTA S. DONATO, 5 - 40126 BOLOGNA, ITALY

*E-mail address:* [stefano.biagi3@unibo.it](mailto:stefano.biagi3@unibo.it)

DIPARTIMENTO DI MATEMATICA, SAPIENZA UNIVERSITÀ DI ROMA, P.LE ALDO MORO 5, 00185 ROMA, ITALY

*E-mail address:* [tralli@mat.uniroma1.it](mailto:tralli@mat.uniroma1.it)