

*Students' Solutions Manual for*

A Discrete  
Transition  
to Advanced  
Mathematics

**Bettina Richmond**  
**Thomas Richmond**

# Contents

<b>1</b>	<b>Sets and Logic</b>	<b>1</b>
1.1	Sets . . . . .	1
1.2	Set Operations . . . . .	2
1.3	Partitions . . . . .	3
1.4	Logic and Truth Tables . . . . .	4
1.5	Quantifiers . . . . .	5
1.6	Implications . . . . .	6
<b>2</b>	<b>Proofs</b>	<b>9</b>
2.1	Proof Techniques . . . . .	9
2.2	Mathematical Induction . . . . .	10
2.3	The Pigeonhole Principle . . . . .	12
<b>3</b>	<b>Number Theory</b>	<b>15</b>
3.1	Divisibility . . . . .	15
3.2	The Euclidean Algorithm . . . . .	16
3.3	The Fundamental Theorem of Arithmetic . . . . .	17
3.4	Divisibility Tests . . . . .	18
3.5	Number Patterns . . . . .	19
<b>4</b>	<b>Combinatorics</b>	<b>23</b>
4.1	Getting from Point A to Point B . . . . .	23
4.2	The Fundamental Principle of Counting . . . . .	24
4.3	A Formula for the Binomial Coefficients . . . . .	25
4.4	Combinatorics with Indistinguishable Objects . . . . .	25
4.5	Probability . . . . .	26
<b>5</b>	<b>Relations</b>	<b>29</b>
5.1	Relations . . . . .	29
5.2	Equivalence Relations . . . . .	30
5.3	Partial Orders . . . . .	31
5.4	Quotient Spaces . . . . .	32

<b>6</b>	<b>Functions and Cardinality</b>	<b>35</b>
6.1	Functions . . . . .	35
6.2	Inverse Relations and Inverse Functions . . . . .	36
6.3	Cardinality of Infinite Sets . . . . .	37
6.4	An Order Relation on Cardinal Numbers . . . . .	38
<b>7</b>	<b>Graph Theory</b>	<b>39</b>
7.1	Graphs . . . . .	39
7.2	Matrices, Digraphs, and Relations . . . . .	40
7.3	Shortest Paths in Weighted Graphs . . . . .	41
7.4	Trees . . . . .	42
<b>8</b>	<b>Sequences</b>	<b>45</b>
8.1	Sequences . . . . .	45
8.2	Finite Differences . . . . .	46
8.3	Limits of Sequences of Real Numbers . . . . .	47
8.4	Some Convergence Properties . . . . .	48
8.5	Infinite Arithmetic . . . . .	49
8.6	Recurrence Relations . . . . .	50
<b>9</b>	<b>Fibonacci Numbers and Pascal's Triangle</b>	<b>53</b>
9.1	Pascal's Triangle . . . . .	53
9.2	The Fibonacci Numbers . . . . .	54
9.3	The Golden Ratio . . . . .	56
9.4	Fibonacci Numbers and the Golden Ratio . . . . .	56
9.5	Pascal's Triangle and the Fibonacci Numbers . . . . .	58
<b>10</b>	<b>Continued Fractions</b>	<b>59</b>
10.1	Finite Continued Fractions . . . . .	59
10.2	Convergents of a Continued Fraction . . . . .	60
10.3	Infinite Continued Fractions . . . . .	60
10.4	Applications of Continued Fractions . . . . .	61

This solution manual accompanies *A Discrete Transition to Advanced Mathematics* by Bettina Richmond and Tom Richmond. The text contains over 650 exercises. This manual includes solutions to parts of 210 of them.

These solutions are presented as an aid to learning the material, and not as a substitute for learning the material. You should attempt to solve each problem on your own and consult the solutions manual only as a last resort.

It is important to note that there are many different ways to solve most of the exercises. Looking up a solution before following through with your own approach to a problem may stifle your creativity. Consulting the solution manual after finding your own solution might reveal a different approach. There is no claim that the solutions presented here are the “best” solutions. These solutions use only techniques which should be familiar to you.

# Chapter 1

## Sets and Logic

### 1.1 Sets

- (a) True (b) The elements of a set are not ordered, so there is no “first” element of a set.
- $|\{M, I, S, S, I, S, S, I, P, P, I\}| = |\{M, I, S, P\}| = 4 < 7 = |\{F, L, O, R, I, D, A\}|$ .
- $\{1, 2, 3\} \subseteq \{1, 2, 3, 4\}$
  - $3 \in \{1, 2, 3, 4\}$
  - $\{3\} \subseteq \{1, 2, 3, 4\}$
  - $\{a\} \in \{\{a\}, \{b\}, \{a, b\}\}$
  - $\emptyset \subseteq \{\{a\}, \{b\}, \{a, b\}\}$
  - $\{\{a\}, \{b\}\} \subseteq \{\{a\}, \{b\}, \{a, b\}\}$
- A 0-element set  $\emptyset$  has  $2^0 = 1$  subset, namely  $\emptyset$ .
  - A 1-element set  $\{1\}$  has  $2^1 = 2$  subsets, namely  $\emptyset$  and  $\{1\}$ .
  - A 2-element set has  $2^2 = 4$  subsets.  
A 3-element set has  $2^3 = 8$  subsets.  
A 4-element set  $\{1, 2, 3, 4\}$  should have  $2^4 = 16$  subsets
  - The 16 subsets of  $\{1, 2, 3, 4\}$  are:  
$$\emptyset, \{1\}, \{2\}, \{3\}, \{4\}, \{1, 2\}, \{1, 3\}, \{1, 4\}, \{2, 3\}, \{2, 4\}, \{3, 4\},$$
$$\{1, 2, 3\}, \{1, 2, 4\}, \{1, 3, 4\}, \{2, 3, 4\}, \{1, 2, 3, 4\}$$
  - A 5-element set has  $2^5 = 32$  subsets.  
A 6-element set has  $2^6 = 64$  subsets.  
An  $n$ -element set has  $2^n$  subsets.

8. (a) 3, 4, 5, and 7:  $|S_3| = |\{t, h, r, e\}| = 4 = |S_4| = |\{f, o, u, r\}| = |S_5| = |\{f, i, v, e\}| = |S_7| = |\{s, e, v, n\}|$ .
- (b)  $S_{21} = S_{22}$  or  $S_{2002} = S_{2000}$ , for example.
- (c)  $a \in S_{1000}$  and  $a \notin S_k$  for  $k = 1, 2, \dots, 999$ .
- (d) (i) True (ii) True (iii) True (iv) False (v) False (vi) True (vii) True  
 (viii) False (ix) True (x) True (xi) True:  $\{n, i, e\} = S_9 \in \mathcal{S}$ . (xii) True  
 (xiii) False (xiv) True (xv) True (xvi) False
9. (a)  $D_1 = \emptyset, D_2 = \{2\}, D_{10} = \{2, 5\}, D_{20} = \{2, 5\}$
- (b) (i) True (ii) False (iii) False (iv) True (v) True (vi) False (vii) True  
 (viii) False (ix) True (x) True (xi) False (xii) True
- (c)  $|D_{10}| = |\{2, 5\}| = 2; |D_{19}| = |\{19\}| = 1$ .
- (d) Observe that  $D_2 = D_4 = D_8 = D_{16}, D_6 = D_{12} = D_{18}, D_3 = D_9, D_{10} = D_{20}$ .  
 Thus  $|\mathcal{D}| = |\{D_1, D_2, \dots, D_{20}\}| = |\{D_1, D_2, D_3, D_5, D_6, D_7, D_{10}, D_{11}, D_{13}, D_{14}, D_{15}, D_{17}, D_{19}\}| = 13$ .
10. For example, let  $S_1 = S_2 = S_3 = \{1, 2, 3\}, S_4 = \{4\}$ , and  $S_5 = \{5\}$ . Now  $\mathcal{S} = \{S_k\}_{k=1}^5 = \{\{1, 2, 3\}, \{4\}, \{5\}\}$ , so  $|\mathcal{S}| = 3$ .


## 1.2 Set Operations

1. (a)  $S \cap T = \{1, 3, 5\}$
- (b)  $S \cup T = \{1, 2, 3, 4, 5, 7, 9\}$
- (c)  $S \cap V = \{3, 9\}$
- (d)  $S \cup V = \{1, 3, 5, 6, 7, 9\}$
- (e)  $(T \cap V) \cup S = \{3\} \cup S = S = \{1, 3, 5, 7, 9\}$
- (f)  $T \cap (V \cup S) = T \cap \{1, 3, 5, 6, 7, 9\} = \{1, 3, 5\}$ .
- (g)  $V \times T = \{(3, 1), (3, 2), (3, 3), (3, 4), (3, 5), (6, 1), (6, 2), (6, 3), (6, 4), (6, 5), (9, 1), (9, 2), (9, 3), (9, 4), (9, 5)\}$
- (h)  $U \times (T \cap S) = \{(3, 1), (3, 3), (3, 5), (6, 1), (6, 3), (6, 5), (9, 1), (9, 3), (9, 5)\}$ .
2. (a)  $A \cap D = \{A \diamond\}$ ; cardinality 1
- (c)  $A \cap (S \cup D) = \{A \spadesuit, A \diamond\}$ ; cardinality 2
- (e)  $(A \cap S) \cup (K \cap D) = \{A \spadesuit, K \diamond\}$ ; cardinality 2
- (g)  $K \cap S^c = \{K \clubsuit, K \diamond, K \heartsuit\}$ ; cardinality 3
- (i)  $(A \cup K)^c \cap S = \{2 \spadesuit, 3 \spadesuit, 4 \spadesuit, 5 \spadesuit, 6 \spadesuit, 7 \spadesuit, 8 \spadesuit, 9 \spadesuit, 10 \spadesuit, J \spadesuit, Q \spadesuit\}$ ; cardinality 11
- (n)  $K \setminus S = \{K \heartsuit, K \clubsuit, K \diamond\}$ ; cardinality 3

7.  $(x, y) \in A \times (B \cap C) \iff x \in A, y \in B \cap C$   
 $\iff x \in A, y \in B \text{ and } y \in C$   
 $\iff x \in A \text{ and } y \in B \text{ and } x \in A \text{ and } y \in C$   
 $\iff (x, y) \in (A \times B) \cap (A \times C).$

This shows that the elements of  $A \times (B \cap C)$  are precisely those of  $(A \times B) \cap (A \times C)$ , and thus the two sets are equal.

9. The conditions are not equivalent. For example, the collection  $\{S_1, S_2\}$  where  $S_1 = S_2 \neq \emptyset$  satisfies  $(S_i \cap S_j \neq \emptyset \Rightarrow S_i = S_j)$ , but not  $(S_i \cap S_j \neq \emptyset \Rightarrow i = j)$ . However, if the sets of the collection  $\{S_i | i \in I\}$  are distinct, the statements will be equivalent.
10. Let  $A$  be the set of students taking Algebra and let  $S$  be the set of students taking Spanish. Now  $|A \cup S| = |A| + |S| - |A \cap S| = 43 + 32 - 7 = 68$ . Thus, there are 68 students taking Algebra or Spanish.
12. A tree diagram for the outcomes will have 2 branches for the choice of meat, each stem of which has 7 branches for the possible choices for vegetables, and each of these stems has 5 branches for the choice of dessert. Thus, 2 choices for meat, 7 choices for vegetable, and 5 choices for dessert give  $2 \cdot 7 \cdot 5 = 70$  choices for the special.
15. Observe that there are not  $4 \cdot 3$  options, for Luis can not take both physics and chemistry at 2:00. There are only 11 scheduling options, as shown in the tree diagram below.



### 1.3 Partitions

3. (a) Not necessarily. Some  $B_i$  may be empty.  
 (b) Yes ( $S \neq \emptyset$  and  $L \neq \emptyset$ ),  $S \cup L = B$ , and  $S \cap L = \emptyset$ .  
 (c) No.  $S$  and  $P$  partition  $A$ , but  $D$  has nonempty intersection with  $S$  or  $P$  yet  $D \neq S$  and  $D \neq P$ .  
 (d) No.  $X = \emptyset$ .  
 (e) No.  $R \cap S = S \neq \emptyset$ , but  $R \neq S$ .

5. (a) Yes.  
 (b) No.  $L_3 \neq L_4$  even though  $L_3 \cap L_4 = \{(0, 0)\} \neq \emptyset$ . Also,  $(0, 1) \notin \bigcup \mathcal{D}$ .  
 (c) Yes.  
 (d) Yes.  
 (e) No.  $(0, 1) \notin \bigcup \mathcal{G}$ . Also,  $P_3 \neq P_4$  yet  $P_3 \cap P_4 = \{(0, 0)\} \neq \emptyset$ .  
 (f) No.  $(\pi, \pi) \notin \bigcup \mathcal{H}$ .

8. Each  $C_i$  is nonempty: Given  $i \in I$ ,  $B_i \neq \emptyset$ , so  $\exists b \in B_i$ , and  $\sqrt{b} \in C_i$ .

$\mathcal{C}$  is a mutually disjoint collection: If  $C_i \cap C_j \neq \emptyset$ , then  $\exists z \in C_i \cap C_j$ , and from the definition of  $C_i$  and  $C_j$ , we have  $z^2 \in B_i \cap B_j$ . Since  $B_i \cap B_j \neq \emptyset$  and  $\{B_i | i \in I\}$  is a partition, it follows that  $B_i = B_j$ , so  $\{x \in \mathbb{R} | x^2 \in B_i\} = \{x \in \mathbb{R} | x^2 \in B_j\}$ , that is,  $C_i = C_j$ .

$\bigcup \mathcal{C} = \mathbb{R}$ : Clearly  $\bigcup \mathcal{C} \subseteq \mathbb{R}$ , so it suffices to show  $\mathbb{R} \subseteq \bigcup \mathcal{C}$ . Given  $x \in \mathbb{R}$ ,  $x^2 \in [0, \infty) = \bigcup \mathcal{B}$ , so  $x^2 \in B_i$  for some  $i \in I$ , which shows  $x \in C_i$ . Thus,  $x \in \mathbb{R} \Rightarrow x \in \bigcup \mathcal{C}$ , as needed.

11. (a) Given any partition  $\mathcal{P}$  of  $S$ , each block of  $\mathcal{P}$  may be partitioned into singleton sets (that is, into blocks of  $\mathcal{D}$ ), so  $\mathcal{D}$  is finer than any partition  $\mathcal{P}$  of  $S$ .  
 (b) The coarsest partition of a set  $S$  is the one-block partition  $\mathcal{I} = \{S\}$ . Given any partition  $\mathcal{P}$  of  $S$ , the block  $S$  of  $\mathcal{I}$  is further partitioned by the blocks of  $\mathcal{P}$ , so ever partition  $\mathcal{P}$  of  $S$  is finer than  $\mathcal{I}$ .  
 (c) Since each block of the coarser partition  $\mathcal{Q}$  is the union of one or more blocks of the finer partition  $\mathcal{P}$ , we have  $|\mathcal{P}| \geq |\mathcal{Q}|$ .  
 (d) No.  $\mathcal{P} = \{(-\infty, 0], (0, \infty)\}$  and  $\mathcal{Q} = \{(-\infty, 5], (5, 6), [6, \infty)\}$  are partitions of  $\mathbb{R}$  with  $|\mathcal{P}| \leq |\mathcal{Q}|$ , but neither partition is a refinement of the other.

## 1.4 Logic and Truth Tables

1. (a)  $S \wedge \sim G$       (b)  $H \vee \sim S$       (c)  $\sim (S \wedge G)$       (d)  $(S \wedge G) \vee (\sim H)$   
 (e)  $(S \vee \sim S) \wedge G$       (f)  $S \wedge H \wedge G$       (g)  $(S \wedge H) \vee (\sim G)$

7. (a)

$P$	$Q$	$P \wedge Q$	$\sim (P \wedge Q)$	$\sim Q$	$\sim (P \wedge Q) \wedge \sim Q$
T	T	T	F	F	F
T	F	F	T	T	T
F	T	F	T	F	F
F	F	F	T	T	T

$\sim (P \wedge Q) \wedge \sim Q = \sim Q$  since the columns for these two statements are identical.

- (b) Note that if  $Q$  fails, then  $(P \wedge Q)$  fails, so that  $Q$  fails and  $(P \wedge Q)$  fails. On the other hand, if  $Q$  fails and some other conditions occur (namely,  $(P \wedge Q)$  fails), then  $Q$  fails.



10. Answers may vary.

$P$	$Q$	(a)	(b)	(c)	(d)	(e)	(f)	(g)	(h)	(i)
T	T	T	F	F	F	T	T	T	F	F
T	F	T	T	T	F	T	F	T	F	T
F	T	T	F	F	F	T	T	F	T	T
F	F	F	T	F	T	T	T	T	F	T

(a)  $P \vee Q$  (b)  $\sim Q$  (c)  $P \wedge \sim Q$  (d)  $\sim P \wedge \sim Q$  (e)  $P \vee \sim P$  (f)  $\sim P \vee Q$   
 (g)  $P \vee \sim Q$  (h)  $\sim P \wedge Q$  (i)  $\sim P \vee \sim Q$

12. The placement of the parentheses in  $P \vee Q \wedge R$  is important:

$(P \vee Q) \wedge R \neq P \vee (Q \wedge R)$ , as the truth table below indicates.

$P$	$Q$	$R$	$(P \vee Q) \wedge R$	$P \vee (Q \wedge R)$
T	T	T	T	T
T	T	F	F	T
T	F	T	T	T
T	F	F	F	T
F	T	T	T	T
F	T	F	F	F
F	F	T	F	F
F	F	F	F	F

14.

$P$	$Q$	$R$	(a)	(b)	(c)	(d)	(e)
T	T	T	T	F	F	T	T
T	T	F	F	F	F	T	T
T	F	T	F	F	F	T	T
T	F	F	F	F	T	F	T
F	T	T	F	F	F	T	T
F	T	F	F	F	F	T	T
F	F	T	F	F	F	T	F
F	F	F	F	T	F	T	T

(a)  $P \wedge Q \wedge R$  (b)  $\sim P \wedge \sim Q \wedge \sim R$  (c)  $P \wedge \sim Q \wedge \sim R$   
 (d)  $\sim (P \wedge \sim Q \wedge \sim R)$  (e)  $\sim (\sim P \wedge \sim Q \wedge \sim R)$

## 1.5 Quantifiers

- $\forall \epsilon \in (0, \infty) \exists n \in \mathbb{N}$  such that  $\frac{1}{n} < \epsilon$ .
  - $\forall e \in \{2k \mid k \in \mathbb{N} \setminus \{1\}\} \exists a \in \{2n \mid n \in \mathbb{Z}\}$  and  $\exists p \in \{\text{prime numbers}\}$  such that  $e = ap$ .
  - $\forall \epsilon \in (0, \infty) \exists \delta \in (0, \infty)$  such that  $x^2 < \epsilon$  whenever  $|x| < \delta$ .
  - $\exists m \in \mathbb{Z}$  such that  $\forall x \in \mathbb{Z} \exists y \in \mathbb{Z}$  with  $xy = m$ .
  - $\forall n \in \mathbb{N} \setminus \{1\} \exists p \in \{\text{prime numbers}\}$  such that  $n < p < n^2$ .
- True. Take  $x = \pm 1$ .  
 Negation:  $\forall x \in \mathbb{Z}, \exists y \in \mathbb{Z}$  such that  $\frac{y}{x} \notin \mathbb{Z}$ .

- (b) False. For  $a = 0$ ,  $\frac{b}{a}$  is not even defined.  
Negation:  $\exists a \in \mathbb{Z}$  such that  $\forall b \in \mathbb{Z}$ ,  $\frac{b}{a} \notin \mathbb{Z}$ .
- (c) True.  $\forall u \in \mathbb{N}$ , take  $v = 2u$ .  
Negation:  $\exists u \in \mathbb{N}$  such that  $\forall v \in \mathbb{N} \setminus \{u\}$ ,  $\frac{v}{u} \notin \mathbb{N}$ .
- (d) False. For  $u = 1$ ,  $\frac{1}{v} \notin \mathbb{N} \forall v \in \mathbb{N}$ .  
Negation:  $\exists u \in \mathbb{N}$  such that  $\forall v \in \mathbb{N} \setminus \{u\}$ ,  $\frac{u}{v} \notin \mathbb{N}$ .
- (e) True.  $\forall a \in \mathbb{N}$ , take  $b = a^2$  and  $c = a$ .  
Negation:  $\exists a \in \mathbb{N}$  such that  $\forall b, c \in \mathbb{N}$ ,  $ab \neq c^3$ .
6. (a)  $\exists a, b \in S$  such that  $\forall n \in \mathbb{N}$ ,  $na \leq b$ .  
(b) (i) No. (ii) Yes. (iii) No. (iv) Yes.

## 1.6 Implications

4. (a)  $S \Rightarrow U$  is false if and only if the stock market goes up but unemployment does not go up.  
(b) The converse of  $S \Rightarrow U$  is false if and only if unemployment goes up but the stock market does not go up.  
(c) The contrapositive of  $\sim I \Rightarrow U$  is false if and only if unemployment does not go up and interest rates do not go down.
6. (a)  $x^2 = 4$  only if  $x = 2$ . False.  
Converse:  $x^2 = 4$  if  $x = 2$ . True.
- (b) If  $2x \leq x$ , then  $x^2 > 0$ . False (consider  $x = 0$ ).  
Converse: If  $x^2 > 0$ , then  $2x \leq x$ . False.
- (c) If 2 is a prime number, then  $2^2$  is a prime number. False.  
Converse: If  $2^2$  is a prime number, then 2 is a prime number. True.
- (d) If  $x$  is an integer then  $\sqrt{x}$  is an integer. False.  
Converse: If  $\sqrt{x}$  is an integer, then  $x$  is an integer. True.
- (e) If every line has a  $y$ -intercept, then every line contains infinitely many points. True.  
Converse: If every line contains infinitely many points then every line has a  $y$ -intercept. False.
- (f) A line has undefined slope only if it is vertical. True.  
Converse: A line has undefined slope if it is vertical. True.
- (g)  $x = -5$  only if  $x^2 - 25 = 0$ . True.  
Converse:  $x = -5$  if  $x^2 - 25 = 0$ . False.
- (h)  $x^2$  is positive only if  $x$  is positive. (Assume  $x \in \mathbb{R}$ .) False.  
Converse:  $x^2$  is positive if  $x$  is positive. True.
7. (a) “ $m$  is a multiple of 8” is sufficient but not necessary for  $\frac{m}{2} \in \mathbb{Z}$ .  
(b) “ $m \in \mathbb{Z}$ ” is necessary but not sufficient for  $\frac{m}{2} \in \mathbb{Z}$ .

(c) “ $m$  is a multiple of 2” is a necessary and sufficient condition on  $m$  for  $\frac{m}{2} \in \mathbb{Z}$ .

10. (a)

$P$	$Q$	$P \Rightarrow Q$	$\sim P \Rightarrow Q$	$\sim P \vee Q$	$P \vee Q$	$\sim(P \Rightarrow \sim Q)$	$P \wedge Q$
T	T	T	T	T	T	T	T
T	F	F	T	F	T	F	F
F	T	T	T	T	T	F	F
F	F	T	F	T	F	F	F

(b) (iii) and (v):  $(P \Rightarrow Q) = (\sim P \vee Q)$ ; (iv) and (vi):  $(\sim P \Rightarrow Q) = (P \vee Q)$ ; (vii) and (viii):  $\sim(P \Rightarrow \sim Q) = (P \wedge Q)$ .

11. (c)

$P$	$Q$	$S$	$P \Rightarrow S$	$Q \Rightarrow S$	$(P \Rightarrow S) \vee (Q \Rightarrow S)$	$P \vee Q$	$(P \vee Q) \Rightarrow S$
T	T	T	T	T	T	T	T
T	T	F	F	F	F	T	F
T	F	T	T	T	T	T	T
T	F	F	F	T	T	T	F
F	T	T	T	T	T	T	T
F	T	F	T	F	T	T	F
F	F	T	T	T	T	F	T
F	F	F	T	T	T	F	T

Because the columns corresponding to  $(P \Rightarrow S) \vee (Q \Rightarrow S)$  and  $(P \vee Q) \Rightarrow S$  are not identical, the statements are not equivalent.



# Chapter 2

## Proofs

### 2.1 Proof Techniques

2. Partition the set  $L$  of lattice points inside a given circle  $C$  into blocks  $B(x, y)$  where, for  $(x, y) \in L$ ,  $B(x, y)$  contains  $(x, y)$  and  $(x, y)$  rotated around the origin by  $90^\circ$ ,  $180^\circ$ , and  $270^\circ$ . Now for each  $(x, y) \in L \setminus \{(0, 0)\}$ , the block  $B(x, y)$  contains 4 elements, and  $B(0, 0)$  contains one element. Since  $|L|$  is the sum of  $|B(x, y)|$  taken over all distinct blocks, we have  $|L| = 4k + 1$  where  $k + 1$  is the number of blocks in this partition.

5. (b) Suppose  $x, y \geq 0$  are given. Then

$$\begin{aligned} 0 \leq \lfloor x \rfloor &\leq x \\ \text{and } 0 \leq \lfloor y \rfloor &\leq y. \end{aligned}$$

Multiplying these equations gives

$$\lfloor x \rfloor \lfloor y \rfloor \leq xy,$$

so  $\lfloor x \rfloor \lfloor y \rfloor$  is an integer which is  $\leq xy$ . By definition,  $\lfloor xy \rfloor$  is the largest integer which is  $\leq xy$ , so  $\lfloor x \rfloor \lfloor y \rfloor \leq \lfloor xy \rfloor$ . This argument holds for any  $x, y \in [0, \infty)$ .

6. Suppose  $p(x) = ax^2 + bx + c$  and  $p(1) = p(-1)$ . The equation  $p(1) = p(-1)$  becomes  $a + b + c = a - b + c$ , and subtracting  $(a + c)$  from both sides gives  $b = -b$ , so  $b = 0$ . Thus,  $p(x) = ax^2 + c$ , so  $p(2) = 2^2a + c = (-2)^2a + c = p(-2)$ .

Conversely, Suppose  $p(x) = ax^2 + bx + c$  and  $p(2) = p(-2)$ . The equation  $p(2) = p(-2)$  becomes  $4a + 2b + c = 4a - 2b + c$ , and again we find that  $b = 0$ . Thus,  $p(x) = ax^2 + c$ , so  $p(1) = 1^2a + c = (-1)^2a + c = p(-1)$ .

8. Note that  $n^3 + n = n(n^2 + 1)$ . Since  $n$  and  $n^2$  have the same parity,  $n$  and  $n^2 + 1$  have opposite parities (that is, one is even and the other is odd). Since any multiple of an even number is even, it follows that  $n(n^2 + 1) = n^3 + n$  is even.
9. (a) Suppose  $a$  is a multiple of 3, say  $a = 3n$  where  $n \in \mathbb{Z}$ . Then  $a = (n-1) + n + (n+1)$ , the sum of three consecutive integers. Conversely, suppose  $a = k + (k+1) + (k+2)$

is the sum of three consecutive integers. Then  $a = 3(k + 1)$ , so  $a$  is a multiple of 3.

- (b) No. The sum  $1 + 2 + 3 + 4 = 10$  of four consecutive integers is not a multiple of 4, and 8, a multiple of 4, cannot be written as a sum of four consecutive integers:  $0 + 1 + 2 + 3 = 6 < 8 < 10 = 1 + 2 + 3 + 4$ .
- (c) The sum of  $k$  consecutive integers has form  $(n + 1) + (n + 2) + \cdots + (n + k) = kn + (1 + 2 + \cdots + k)$ . Since  $kn$  is a multiple of  $k$ , the sum will be a multiple of  $k$  if and only if  $1 + 2 + \cdots + k$  is a multiple of  $k$ . Thus,

$a$  is a multiple of  $k$  if and only if  $a$  may be written as a sum of  $k$  consecutive integers

is true if and only if  $1 + 2 + \cdots + k$  is a multiple of  $k$ .

We will see later that  $1 + 2 + \cdots + k$  is the  $k^{\text{th}}$  triangular number and is given by the formula  $\frac{k(k+1)}{2}$ . Thus,  $1 + 2 + \cdots + k$  is a multiple of  $k$  if and only if  $\frac{k+1}{2} \in \mathbb{Z}$ , that is, if and only if  $k$  is odd.

12. Direct proof: For any  $k \in \mathbb{Z}$ ,  $x_k = \frac{\pi}{2} + 2\pi k$  is a solution to  $\sin x = 1$ , so  $\sin x = 1$  has infinitely many solutions.

Indirect proof: Suppose to the contrary that  $\sin x = 1$  has only finitely many solutions. The solution set is nonempty since  $\sin(\frac{\pi}{2}) = 1$ . Let  $x_m$  be the largest member of the solution set. Now  $\sin(x_m + 2\pi) = \sin x_m = 1$ , so  $x_m + 2\pi$  is an element of the solution set which is larger than  $x_m$ , contrary to the choice of  $x_m$  as the largest solution. Assuming that there were only finitely many solutions gave a contradiction, so there must be infinitely many solutions.

25. Suppose  $k$  and  $l$  are distinct lines that intersect. Suppose  $A$  and  $B$  are points of intersection of  $k$  and  $l$ . If  $A \neq B$ , then the two distinct points  $A$  and  $B$  determine a unique line, contrary to  $k$  and  $l$  being distinct lines through  $A$  and  $B$ . Thus,  $A = B$ . That is,  $k$  and  $l$  intersect in a unique point.
27. Moving a knight out and back to his original position on the first move effectively gave the other player the first move in the double move chess game. In initial double move chess, moving a knight out and back does not exchange the roles of first player and second player, since the first player was playing initial double move chess and the second player is left with a different game—one in which only one player gets an initial double move.

## 2.2 Mathematical Induction

2. (b) For  $n = 1$ , the statement is  $1^3 = \frac{1^2(1+1)^2}{4}$ , which is true. Suppose the statement holds for  $n = k$ , that is, suppose  $1^3 + 2^3 + \cdots + k^3 = \frac{k^2(k+1)^2}{4}$ . We wish to show that the statement holds for  $n = k + 1$ , that is, we wish to show that  $1^3 + 2^3 + \cdots + k^3 + (k + 1)^3 = \frac{(k+1)^2(k+2)^2}{4}$ . Adding  $(k + 1)^3$  to both sides of the

induction hypothesis gives

$$\begin{aligned} 1^3 + 2^3 + \cdots + k^3 + (k+1)^3 &= \frac{k^2(k+1)^2}{4} + (k+1)^3 \\ &= \left(\frac{(k+1)^2}{4}\right)(k^2 + 4(k+1)) \\ &= \frac{(k+1)^2(k+2)^2}{4}, \end{aligned}$$

as needed. Now the statement holds for  $n = 1$  and for  $n = k + 1$  whenever it holds for  $n = k$ , so by mathematical induction,  $1^3 + 2^3 + 3^3 + \cdots + n^3 = \frac{n^2(n+1)^2}{4}$  for every natural number  $n$ .

- (e) For  $n = 1$ , the statement is  $1^2 = \frac{1(2-1)(2+1)}{3}$ , which is true. Suppose the statement holds for  $n = k$ , that is, suppose  $1^2 + 3^2 + 5^2 + \cdots + (2k-1)^2 = \frac{k(2k-1)(2k+1)}{3}$ . Adding  $(2k+1)^2$  to both sides of this equation gives

$$\begin{aligned} 1^2 + 3^2 + \cdots + (2k-1)^2 + (2k+1)^2 &= \frac{k(2k-1)(2k+1)}{3} + (2k+1)^2 \\ &= \frac{2k+1}{3}(k(2k-1) + 3(2k+1)) \\ &= \frac{2k+1}{3}(2k^2 + 5k + 3) \\ &= \frac{2k+1}{3}(2k+3)(k+1) \\ &= \frac{(k+1)(2(k+1)-1)(2(k+1)+1)}{3}, \end{aligned}$$

so the statement holds for  $n = k + 1$ . Now the statement holds for  $n = 1$  and for  $n = k + 1$  whenever it holds for  $n = k$ , so by mathematical induction,  $1^2 + 3^2 + 5^2 + \cdots + (2n-1)^2 = \frac{n(2n-1)(2n+1)}{3}$  for every natural number  $n$ .

4. We wish to show that for any  $n \in \mathbb{N}$ , there exists  $m \in \mathbb{N}$  such that  $n^3 + (n+1)^3 + (n+2)^3 = 9m$ . Taking  $n = 1$ , we find that  $1^3 + 2^3 + 3^3 = 36 = 9m$  where  $m = 4 \in \mathbb{N}$ . Now suppose that the statement holds for  $n = k$ . Then  $k^3 + (k+1)^3 + (k+2)^3 = 9m$  for some  $m \in \mathbb{N}$ . We wish to show that  $(k+1)^3 + (k+2)^3 + (k+3)^3 = 9j$  for some  $j \in \mathbb{N}$ . But

$$\begin{aligned} (k+1)^3 + (k+2)^3 + (k+3)^3 &= k^3 + (k+1)^3 + (k+2)^3 + (k+3)^3 - k^3 \\ &= 9m + (k+3)^3 - k^3 \\ &= 9m + k^3 + 9k^2 + 27k + 27 - k^3 \\ &= 9m + 9(k^2 + 3k + 1) \\ &= 9j \quad \text{where } j = m + k^2 + 3k + 1 \in \mathbb{N}. \end{aligned}$$

Now the statement holds for  $n = 1$  and for  $n = k + 1$  whenever it holds for  $n = k$ , so by mathematical induction, for any  $n \in \mathbb{N}$ , there exists  $m \in \mathbb{N}$  such that  $n^3 + (n+1)^3 + (n+2)^3 = 9m$ .

9. Suppose  $\alpha > -1, \alpha \neq 0$ . We wish to show  $(1 + \alpha)^n > 1 + n\alpha$  for  $n \geq 2$ . Observe that  $\alpha > -1$  guarantees that the powers  $(1 + \alpha)^n$  are all positive. For  $n = 2$ , the statement is  $(1 + \alpha)^2 > 1 + 2\alpha$ , which is true since  $(1 + \alpha)^2 = 1 + 2\alpha + \alpha^2$ , and  $\alpha^2 > 0$  for  $\alpha \neq 0$ . Now suppose  $(1 + \alpha)^k > 1 + k\alpha$  for  $n = k \geq 2$ . We wish to show  $(1 + \alpha)^{k+1} > 1 + (k + 1)\alpha$ . But

$$\begin{aligned} (1 + \alpha)^{k+1} &= (1 + \alpha)^k(1 + \alpha) \\ &> (1 + k\alpha)(1 + \alpha) \text{ (Induction hypothesis)} \\ &= 1 + k\alpha + \alpha + k\alpha^2 \\ &= 1 + (k + 1)\alpha + k\alpha^2 \\ &> 1 + (k + 1)\alpha \text{ since } k\alpha^2 > 0 \text{ for } \alpha \neq 0. \end{aligned}$$

Now the statement holds for  $n = 2$  and for  $n = k + 1$  whenever it holds for  $n = k$ , so by mathematical induction,  $(1 + \alpha)^n > 1 + n\alpha$  for every natural number  $n \geq 2$ .

14. (b) Any combination of  $m$  4-cent stamps and  $n$  10-cent stamps gives  $(4m + 10n)$ -cents postage. Since  $4m + 10n$  is always even, 4-cent and 10-cent stamps can never be combined to give any odd amount.
15. Assuming that all horses of any  $n$ -element set have the same color, the induction step argues that all horses of an  $n + 1$ -element set  $H = \{h_1, \dots, h_{n+1}\}$  have the same color since all horses of the  $n$ -element set  $H \setminus \{h_1\}$  have the same color C, all horses of the  $n$ -element set  $H \setminus \{h_{n+1}\}$  have the same color D, and  $C = D$  since  $H \setminus \{h_1\} \cap H \setminus \{h_{n+1}\} \neq \emptyset$ . However,  $H \setminus \{h_1\} \cap H \setminus \{h_{n+1}\} = \emptyset$  if  $n = 1$ . Thus, the first induction step (if true for  $n = 1$ , then true for  $n = 2$ ) fails.

## 2.3 The Pigeonhole Principle

3. (a) 24. Worst case: First 8 nickels, 10 dimes, 3 quarters, then 3 pennies.  
 (b) 9. The pigeonhole principle applies. Worst case: 2 of each of the 4 types, then one more.  
 (c) All 33. Worst case: the last coin drawn is a quarter.  
 (d) 25. Worst case: First 12 pennies, 8 nickels, 3 quarters, then 2 dimes.  
 (e) 16. Worst case: First 12 pennies, then 4 more coins to get a second pair.
5. Given 5 lattice points  $(a_i, b_i)$   $i = 1, 2, 3, 4, 5$ , the pigeonhole principle implies that at least three of the integers  $a_1, \dots, a_5$  have the same parity. Without loss of generality, assume  $a_1, a_2$ , and  $a_3$  have the same parity. Now by the pigeonhole principle, at least two of the points  $b_1, b_2, b_3$  must have the same parity. Without loss of generality, assume  $b_1$  and  $b_2$  have the same parity. Now the midpoint of the segment from  $(a_1, b_1)$  to  $(a_2, b_2)$  is  $(\frac{a_1+a_2}{2}, \frac{b_1+b_2}{2})$ , and this is a lattice point since  $a_1$  and  $a_2$  have the same parity and  $b_1$  and  $b_2$  have the same parity.
7. In the worst case, each of the six cameras would receive 23 exposures before the next exposure would give one camera 24 exposures. Thus,  $23 \times 6 + 1 = 139$  exposures are needed to guarantee that one camera has 24 exposures.



9. (a) Partition the balls into “50-sum” sets  $\{1, 49\}, \{2, 48\}, \dots, \{24, 26\}$  and two unpaired singleton  $\{25\}$  and  $\{50\}$ . This gives 26 sets. If balls are drawn and assigned to the appropriate set, to insure that one set receives two balls, we must draw 27 balls.



# Chapter 3

## Number Theory

### 3.1 Divisibility

4. If  $d \mid n^2$ , then it need not be true that  $d \mid n$ . For example,  $4 \mid 6^2$  but  $4 \nmid 6$ .
6. (a) Suppose  $a \mid b$ . Then  $b = na$  for some  $n \in \mathbb{Z}$ . If  $c \in D_a$ , then  $c \mid a$ , so  $a = mc$  for some  $m \in \mathbb{Z}$ , so  $b = na = n(mc) = (nm)c$  where  $nm \in \mathbb{Z}$ , so  $c \mid b$ , that is,  $c \in D_b$ . Thus,  $D_a \subseteq D_b$ . Conversely, suppose  $D_a \subseteq D_b$ . Now  $a \in D_a$  so  $a \in D_b$ , and thus  $a \mid b$ .
- (b) Suppose  $a \mid b$ . Then  $b = na$  for some  $n \in \mathbb{Z}$ . If  $c \in M_b$ , then  $c = mb$  for some  $m \in \mathbb{Z}$ , so  $c = mb = m(na) = (mn)a$  where  $mn \in \mathbb{Z}$ , so  $a \mid c$ . Thus,  $c \in M_a$ . This shows that  $M_b \subseteq M_a$ . Conversely, suppose  $M_b \subseteq M_a$ . Since  $b \in M_b$ , we have  $b \in M_a$ , so  $a \mid b$ .
- (c)  $D_a = D_b \iff D_a \subseteq D_b \text{ and } D_b \subseteq D_a$   
 $\iff a \mid b \text{ and } b \mid a \quad \text{by part (a)}$   
 $\iff a = \pm b \quad \text{(Theorem 3.1.7)}$   
 $\iff |a| = |b|$
- (d)  $M_a = M_b \iff M_a \subseteq M_b \text{ and } M_b \subseteq M_a$   
 $\iff b \mid a \text{ and } a \mid b \quad \text{by part (b)}$   
 $\iff a = \pm b \quad \text{(Theorem 3.1.7)}$   
 $\iff |a| = |b|$
9. (a)  $a = 73, b = 25: q = 0, r = 25$ .  
(b)  $a = 25, b = 73: q = 2, r = 23$ .  
(c)  $a = -73, b = -25: q = 1, r = 48$ .  
(d)  $a = -25, b = -73: q = 3, r = 2$ .  
(e)  $a = 79, b = -17: q = -1, r = 62$ .

- (f)  $a = -17, b = 79: q = -4, r = 11$ .  
 (g)  $a = -37, b = 13: q = 0, r = 13$ .  
 (h)  $a = 13, b = -37: q = -3, r = 2$ .
17. (a) If  $a$  and  $b$  leave a remainder of 2 when divided by 7, then  $a = 7q + 2$  and  $b = 7s + 2$  for some integers  $q$  and  $s$ , and thus  $a - b = 7q + 2 - (7s + 2) = 7(q - s)$  where  $q - s \in \mathbb{Z}$ , so  $7|(a - b)$ .  
 (b) If  $a = 7q + 2$ , then  $10a = 70q + 20 = 70q + 14 + 6 = 7(10q + 2) + 6$ . Thus, by uniqueness of the quotient and remainder when  $10a$  is divided by 7, we have a quotient of  $10q + 2$  and a remainder of 6.
20. We will show  $4|(13^n - 1) \forall n \in \mathbb{N}$  by mathematical induction. If  $n = 1$ , then  $4|(13^1 - 1)$  since  $4|12$ . Suppose  $4|(13^k - 1)$ . We wish to show that  $4|(13^{k+1} - 1)$ . Now

$$\begin{aligned} 13^{k+1} - 1 &= 13(13^k - 1 + 1) - 1 \\ &= 13(13^k - 1) + 13 - 1 \\ &= 13(13^k - 1) + 12 \end{aligned}$$

Now  $4|(13^k - 1)$  by the induction hypothesis and  $4|12$ , so  $4|(13^{k+1} - 1)$ . Now by mathematical induction,  $4|(13^n - 1) \forall n \in \mathbb{N}$ .

Alternatively, the result of Exercise 23 shows that  $12|(13^n - 1) \forall n \in \mathbb{N}$ , and since  $4|12$ , we have  $4|(13^n - 1) \forall n \in \mathbb{N}$ .

## 3.2 The Euclidean Algorithm

5. If  $a, b \in \mathbb{Z}$  and  $z$  and  $w$  are linear combinations of  $a$  and  $b$  using integer coefficients, say  $z = ja + kb$  and  $w = la + mb$  ( $j, k, l, m \in \mathbb{Z}$ ) then a linear combination of  $z$  and  $w$  with integer coefficients has form  $sz + tw$  ( $s, t \in \mathbb{Z}$ ). Now

$$\begin{aligned} sz + tw &= s(ja + kb) + t(la + mb) \\ &= (sj + tl)a + (ks + tm)b \end{aligned}$$

is a linear combination of  $a$  and  $b$  with integer coefficients  $sj + tl$  and  $ks + tm$ .

7.  $\gcd(15, 39) = 3$ , so 3 should divide  $15s + 39t$  for any  $s, t \in \mathbb{Z}$ . The bill should be of form  $15s + 39t$  ( $s, t \in \mathbb{N} \cup \{0\}$ ), so the bill should be a multiple of 3 cents. It is not.
11. Suppose  $a, b, q, r \in \mathbb{Z} \setminus \{0\}$  and  $a = bq + r$ .
- (a)  $\gcd(a, b) = \gcd(b, r)$  is true.

*Proof:* If  $d$  is any common divisor of  $a$  and  $b$ , then  $d$  is a divisor of  $a - bq = r$ . Thus, any common divisor of  $a$  and  $b$  is a common divisor of  $b$  and  $r$ . Conversely, any common divisor of  $b$  and  $r$  must also divide  $b$  and  $bq + r = a$ , and therefore must be a common divisor of  $a$  and  $b$ . This shows that the common divisors of  $a$  and  $b$  are precisely the common divisors of  $b$  and  $r$ , so  $\gcd(a, b) = \gcd(b, r)$ .

- (c) In general,  $\gcd(q, r)$  does not divide  $b$ . For example, with  $a = 45, b = 7, q = 6$ , and  $r = 3$ , we have  $\gcd(q, r) = 3$  but 3 does not divide 7.

### 3.3 The Fundamental Theorem of Arithmetic

3. Substituting the expressions for  $\text{lcm}(m, n)$  and  $\text{gcd}(m, n)$  given in Corollary 3.3.4 into  $mn = \text{gcd}(m, n)\text{lcm}(m, n)$  and observing that  $\min\{m_i, n_i\} + \max\{m_i, n_i\} = m_i + n_i$  proves Corollary 3.3.5.
8. (a) If  $n = p_1^{n_1} p_2^{n_2} \cdots p_j^{n_j}$ , then  $n^k = p_1^{kn_1} p_2^{kn_2} \cdots p_j^{kn_j}$ , and it follows that  $m = n^k$  is a perfect  $k^{\text{th}}$  power if and only if the multiplicity of each prime factor of  $m$  is a multiple of  $k$ .
- (b) Suppose  $m = p_1^{n_1} p_2^{n_2} \cdots p_j^{n_j}$ . If  $m$  is a perfect square, then  $2|n_i$  for each  $i = 1, \dots, j$ . If  $m$  is a perfect cube, then  $3|n_i$  for each  $i = 1, \dots, j$ . If  $m$  is simultaneously a perfect square and a perfect cube, then  $2|n_i$  and  $3|n_i$  for each  $i$ , so the prime factorization of each  $n_i$  contains a 2 and a 3. Thus,  $6|n_i$  for each  $i = 1, \dots, j$ , and it follows that  $m$  is a perfect  $6^{\text{th}}$  power.
10.  $d|a \Rightarrow a = dq$  for some  $q \in \mathbb{Z}$   
 $\Rightarrow a^2 = d^2 q^2$  for  $q^2 \in \mathbb{Z}$   
 $\Rightarrow d^2|a^2$ .

Conversely, suppose  $d^2|a^2$ . Then  $a^2 = d^2 s$  for some  $s \in \mathbb{Z}$ . Consider the prime factorization of  $s = \frac{a^2}{d^2}$ . If the prime factorizations of  $a^2$  and  $d^2$  are  $p_1^{2n_1} \cdots p_j^{2n_j}$  and  $p_1^{2m_1} \cdots p_j^{2m_j}$  respectively, then by dividing we find that the prime factorization of  $s = \frac{a^2}{d^2}$  must be  $p_1^{2(n_1-m_1)} \cdots p_j^{2(n_j-m_j)} = t^2$  where  $t = p_1^{n_1-m_1} \cdots p_j^{n_j-m_j} = \frac{a}{d}$ . Now since  $s = t^2$  is a perfect square,  $a^2 = d^2 s \Rightarrow a^2 = d^2 t^2 \Rightarrow a = \pm dt \Rightarrow d|a$ .

18. Suppose  $p(x) = c_n x^n + c_{n-1} x^{n-1} + \cdots + c_2 x^2 + c_1 x + c_0$  is a polynomial with integer coefficients  $c_0, \dots, c_n$ , and  $r = \frac{a}{b}$  ( $a, b \in \mathbb{Z}, b \neq 0, \text{gcd}(a, b) = 1$ ) is a rational number with  $p(r) = 0$ . Since  $p(r) = 0$ , we have

$$\frac{c_n a^n}{b^n} + \frac{c_{n-1} a^{n-1}}{b^{n-1}} + \cdots + \frac{c_2 a^2}{b^2} + \frac{c_1 a}{b} + c_0 = 0. \quad (3.1)$$

Multiplying both sides of this equation by  $b^n$  and rearranging the terms gives

$$c_n a^n = -c_{n-1} a^{n-1} b - \cdots - c_2 a^2 b^{n-2} - c_1 a b^{n-1} - c_0 b^n.$$

Since  $b$  divides the right hand side of this equation, it must divide the left hand side, so  $b|c_n a^n$ . Since  $\text{gcd}(a, b) = 1$ , we have  $\text{gcd}(b, a^n) = 1$  and thus  $b|c_n$ .

Again multiplying Equation (3.1) by  $b^n$  and rearranging the terms, we find that

$$c_n a^n + c_{n-1} a^{n-1} b + \cdots + c_2 a^2 b^{n-2} + c_1 a b^{n-1} = -c_0 b^n.$$

Since  $a$  divides the left hand side of this equation,  $a$  must divide the right hand side as well, so  $a|c_0 b^n$ . Since  $\text{gcd}(a, b) = 1 = \text{gcd}(a, b^n)$ , it follows that  $a|c_0$ . Together with the result of the previous paragraph, this proves the Rational Root Theorem.

19. (a) To count the number of factors of form  $(a_i - a_j)$  where  $1 \leq i < j \leq m + 1$ , observe that once  $i$  is selected, the inequality  $i < j \leq m + 1$  implies that there are  $m + 1 - i$  possibilities for  $j$ . As  $i$  may assume any value from 1 to  $m$ , the number of factors is  $\sum_{i=1}^m (m + 1 - i) = m + (m - 1) + \cdots + 2 + 1 = T_m$ .
- (b) The pigeonhole principle implies that  $\lceil \frac{m+1}{2} \rceil$  of the integers  $a_1, \dots, a_{m+1}$  must have the same parity. The argument of (a) shows that from any  $s$  integers  $b_1, \dots, b_s$ , we may form  $T_{s-1}$  distinct factors  $(b_i - b_j)$ . Thus, the  $\lceil \frac{m+1}{2} \rceil$  integers from  $a_1, \dots, a_{m+1}$  of the same parity give  $k = T_{\lceil \frac{m+1}{2} \rceil - 1}$  distinct even factors  $(a_i - a_j)$  in  $P$ , and it follows that  $2^k | P$ .

### 3.4 Divisibility Tests

1. (a)  $10a = 110q + 10r = 110q + 11r - r = 11(10q + r) - r = 11q' - r$  where  $q' = 10q + r \in \mathbb{Z}$ .
- (b) The case  $m = 0$  is clear:  $10^0 a = a = 11q + (-1)^0 r$  as given. If  $10^k a = 11q' + (-1)^k r$ , then applying (a) gives  $10(10^k a) = 11q'' - (-1)^k r$  or  $10^{k+1} a = 11q'' + (-1)^{k+1} r$  for some  $q'' \in \mathbb{Z}$ . By mathematical induction,  $10^m a = 11q'' + (-1)^m r$  for any integer  $m \geq 0$ .
- (c) In  $a = 11q + r$ , take  $a = 1, q = 0$ , and  $r = 1$ , so that for any integer  $m \geq 0$ ,  $10^m = 11q'' + (-1)^m$  for some integer  $q''$ .

4. If  $s = \langle d_{2j-1} \cdots d_2 d_1 d_0 \rangle$  has  $2j$  digits then  $t = \langle d_0 d_1 d_2 \cdots d_{2j-1} \rangle$ , so

$$\begin{aligned}
 s + t &= (d_{2j-1}10^{2j-1} + d_{2j-2}10^{2j-2} + \cdots + d_210^2 + d_110 + d_0) \\
 &\quad + (d_010^{2j-1} + d_110^{2j-2} + \cdots + d_{2j-3}10^2 + d_{2j-2}10^1 + d_{2j-1}) \\
 &= d_{2j-1}(10^{2j-1} + 1) + d_{2j-2}(10^{2j-2} + 10^1) \\
 &\quad + d_{2j-3}(10^{2j-3} + 10^2) + \cdots + d_j(10^j + 10^{j-1}) \\
 &\quad + d_{j-1}(10^{j-1} + 10^j) + \cdots + d_2(10^2 + 10^{2j-3}) \\
 &\quad + d_1(10^1 + 10^{2j-2}) + d_0(1 + 10^{2j-1}) \\
 &= d_{2j-1}(10^{2j-1} + 1) + 10d_{2j-2}(10^{2j-3} + 1) \\
 &\quad + 10^2d_{2j-3}(10^{2j-5} + 1) + \cdots + 10^{j-1}d_j(10^1 + 1) \\
 &\quad + 10^{j-1}d_{j-1}(1 + 10^1) + \cdots + 10^2d_2(1 + 10^{2j-5}) \\
 &\quad + 10d_1(1 + 10^{2j-3}) + d_0(1 + 10^{2j-1}).
 \end{aligned}$$

Recalling that  $11|(10^m + 1)$  for any odd number  $m$ , and observing that each term in the last expression above contains a factor of form  $(10^m + 1)$  ( $m$  odd), we have  $11|(s + t)$ .

8. For  $n = \langle d_k d_{k-1} \cdots d_2 d_1 d_0 \rangle$ , we have

$$\begin{aligned}
 n &= \langle d_k d_{k-1} \cdots d_3 000 \rangle + 100d_2 + 10d_1 + d_0 \\
 &= \langle d_k d_{k-1} \cdots d_3 000 \rangle + 96d_2 + 4d_2 + 8d_1 + 2d_1 + d_0 \\
 &= [\langle d_k d_{k-1} \cdots d_3 000 \rangle + 96d_2 + 8d_1] + [4d_2 + 2d_1 + d_0].
 \end{aligned}$$

Since 8 divides each term in the first bracketed expression above, 8 divides that bracketed expression, and it follows from Corollary 3.4.2 that  $8|n$  if and only if 8 divides the second bracketed expression. That is,  $8|n$  if and only if  $8|(4d_2 + 2d_1 + d_0)$ .

11. All of the tests below are direct consequences of Theorem 3.4.3.
- (a) For  $n = \langle d_k \cdots d_2 d_1 d_0 \rangle$ ,  $12|n$  if and only if  $[3|(d_k + \cdots + d_1 + d_0)$  and  $4|\langle d_1 d_0 \rangle]$ .  
Proof:  $12|n$  if and only if  $3|n$  and  $4|n$ , since 3 and 4 are relatively prime.
  - (b) For  $n = \langle d_k \cdots d_2 d_1 d_0 \rangle$ ,  $14|n$  if and only if  $[2|d_0$  and  $7|n]$ . Proof:  $14|n$  if and only if  $2|n$  and  $7|n$ , since 2 and 7 are relatively prime.
  - (c) For  $n = \langle d_k \cdots d_2 d_1 d_0 \rangle$ ,  $15|n$  if and only if  $[5|d_0$  and  $3|(d_k + \cdots + d_1 + d_0)]$ .  
Proof:  $15|n$  if and only if  $5|n$  and  $3|n$ , since 5 and 3 are relatively prime.
  - (d) For  $n = \langle d_k \cdots d_2 d_1 d_0 \rangle$ ,  $18|n$  if and only if  $[9|(d_k + \cdots + d_1 + d_0)$  and  $2|d_0]$ .  
Proof:  $18|n$  if and only if  $9|n$  and  $2|n$ , since 9 and 2 are relatively prime.
  - (e) For  $n = \langle d_k \cdots d_2 d_1 d_0 \rangle$ ,  $75|n$  if and only if  $[25|\langle d_1 d_0 \rangle$  and  $3|(d_k + \cdots + d_1 + d_0)]$ .  
Proof:  $75|n$  if and only if  $25|n$  and  $3|n$ , since 25 and 3 are relatively prime.
17. Suppose that the sum of the digits of  $a$  and the sum of the digits of  $5a$  both equal  $k$ . Then  $a = 9q + k$  and  $5a = 9n + k$  for some integers  $q, n$ . Now  $4a = 5a - a = 9(n - q)$ , so  $9|4a$ . Since 9 and 4 are relatively prime, we have  $9|a$ .

### 3.5 Number Patterns

1. 
$$\begin{aligned} \frac{1 + 3 + \cdots + (2n - 1)}{(2n + 1) + \cdots + (4n - 1)} &= \frac{1 + 3 + \cdots + (2n - 1)}{(1 + 3 + \cdots + (4n - 1)) - (1 + 3 + \cdots + (2n - 1))} \\ &= \frac{n^2}{(2n)^2 - n^2} \\ &= \frac{n^2}{3n^2} \\ &= \frac{1}{3} \end{aligned}$$
3. (c)  $1 \cdot 2 \cdots j + 2 \cdot 3 \cdots (j + 1) + \cdots (n)(n + 1) \cdots (n + j - 1) = \frac{n(n+1)\cdots(n+j)}{j+1}$  for all  $j, n \in \mathbb{N}$ . Suppose  $j \in \mathbb{N}$  is given. The case  $n = 1$  is clearly true. If  $1 \cdot 2 \cdots j + 2 \cdot 3 \cdots (j + 1) + \cdots (k)(k + 1) \cdots (k + j - 1) = \frac{k(k+1)\cdots(k+j)}{j+1}$ , then
- $$\begin{aligned} &1 \cdot 2 \cdots j + 2 \cdot 3 \cdots (j + 1) + \cdots (k)(k + 1) \cdots (k + j - 1) + (k + 1)(k + 2) \cdots (k + j) \\ &= \frac{k(k + 1) \cdots (k + j)}{j + 1} + (k + 1)(k + 2) \cdots (k + j) \\ &= (k + 1) \cdots (k + j) \left( \frac{k}{j + 1} + 1 \right) \\ &= (k + 1) \cdots (k + j) \left( \frac{k + j + 1}{j + 1} \right), \end{aligned}$$

as needed. Now by mathematical induction, the result holds for all  $n \in \mathbb{N}$ . Since  $j \in \mathbb{N}$  was arbitrary, this completes the proof.





$$\begin{aligned} &= \sum_{j=1}^n (16T_n^2 + j^2) + 8T_n \sum_{j=1}^n j \\ &= \sum_{j=1}^n (16T_n^2 + 8jT_n + j^2) \\ &= \sum_{j=1}^n (4T_n + j)^2. \end{aligned}$$

12.  $(1, 3, \dots, (2n-1)) \cdot (n, n, \dots, n)$

$$\begin{aligned} &= 1n + 3n + \dots + (2n-1)n \\ &= n(1 + 3 + \dots + (2n-1)) \\ &= n(n^2) \\ &= n^3 \end{aligned}$$



## Chapter 4

# Combinatorics

### 4.1 Getting from Point A to Point B

2.  $\binom{7}{4} = 35$ .
  
4. Since  $2310 = 2 \cdot 3 \cdot 5 \cdot 7 \cdot 11$ , a divisor of 2310 having exactly 3 prime factors will be of form  $p_1 p_2 p_3$  where  $p_1, p_2, p_3 \in \{2, 3, 5, 7, 11\}$ . There are  $\binom{5}{3} = 10$  ways to pick three primes  $p_1, p_2, p_3$  from the set  $\{2, 3, 5, 7, 11\}$ , and thus there are 10 divisors of 2310 having exactly three prime factors.
  
8. One edge, say the top edge, of the  $n \times n$  grid requires  $n$  matchsticks. There are  $n + 1$  parallel copies of  $n$  matchsticks in the grid, the last one being the bottom edge. Thus,  $n(n + 1)$  matchsticks are required to draw in the horizontal part of the grid. Similarly,  $n(n + 1)$  matchsticks are needed for the vertical part, for a total of  $2n(n + 1) = 4T_n$  matchsticks.
  
9. For  $n = 1, \dots, 6$ , the table below shows the routes from  $A$  to  $B$  which make exactly  $n$  turns. Each route is seven blocks, three of which are to the west (denoted by  $w$ ) and four of which are to the east (denoted by  $e$ ). The numbers in the bottom row of the table show the number of routes from  $A$  to  $B$  which make exactly  $n$  turns.

1	2	3	4	5	6
wwweeee	wweeeew	wweweee	weweeew	wewewee	ewewewe
eeeeeww	ewwweee	wweewee	ewwewee	weweeew	
	weeeeww	wweeewe	ewweewe	weewewe	
	eeewwee	wewweee	weeweew	eweweeew	
	eeewwwe	ewweeew	ewewwee	eweewew	
		weewwee	eweewwe	eewewew	
		weewwee	weewwee		
		weeeeww	eewwewe		
		eewweew	eewewwe		
		eewweew			
		eewweew			
		eewweew			
		eewweew			
		eewweew			
2	5	12	9	6	1

## 4.2 The Fundamental Principle of Counting

1. A positive divisor of  $n = p_1^{n_1} p_2^{n_2} p_3^{n_3} \cdots p_k^{n_k}$  has form  $p_1^{m_1} p_2^{m_2} p_3^{m_3} \cdots p_k^{m_k}$  where  $0 \leq m_j \leq n_j$  for each  $j = 1, \dots, k$ . Thus, the number of positive divisors of  $n$  is the number of ways to choose a sequence  $(m_1, \dots, m_k)$  of whole numbers satisfying  $0 \leq m_j \leq n_j$  for each  $j = 1, \dots, k$ . As there are  $n_j + 1$  choices for  $m_j$  ( $j = 1, \dots, k$ ), the Fundamental Principle of Counting tells us that there are  $(n_1 + 1)(n_2 + 1) \cdots (n_k + 1)$  positive divisors of  $n$ . Applying this to  $2^3 3^2 7^1 11^1$ , we see that there are  $(4)(3)(2)(2) = 48$  positive divisors of  $2^3 3^2 7^1 11^1$ .

3. There are two choices—depressed or not—for each of the four valves, so there are  $2^4 = 16$  fingering positions for a four-valve instrument. Equivalently, each fingering position corresponds to a subset of valves to be depressed. There are 4 valves and  $2^4 = 16$  possible subsets.

If the third and fourth valves are not to be depressed simultaneously, then there are three choices for positions of the third and fourth valves: only the third valve depressed, only the fourth valve depressed, or neither valve depressed. These 3 options follow the 2 options (depressed or not) for the first valve and the 2 options (depressed or not) for the second valve. This gives a total of  $2 \cdot 2 \cdot 3 = 12$  fingering positions in which the third and fourth valve are not depressed simultaneously.

5. There are  $\binom{7}{4}$  ways to select the Democrats and  $\binom{9}{4}$  ways to select the Republicans, so there are  $\binom{7}{4} \binom{9}{4} = 35 \cdot 126 = 4410$  ways to make the appointments.
8. (a) There are 26 choices for the first letter, 26 choices for the second letter, 26 choices for the third letter, 10 choices for the first digit, 10 choices for the second digit, and 10 choices for the third digit, for a total of  $26^3 10^3 = 17,576,000$  possible license plates.
  - (b)  $26^3 10^3 \binom{6}{3} = 351,520,000$ ; There are 26 choices for each of the three letters,

10 choices for each of the three digits, and  $\binom{6}{3}$  ways to choose three of the six positions for the letters.

### 4.3 A Formula for the Binomial Coefficients

5. (a)  $P(11, 3) = 11 \cdot 10 \cdot 9 = 990$   
 (b)  $P(11, 3)P(11, 3)P(10, 3)P(8, 3) = (11 \cdot 10 \cdot 9)(11 \cdot 10 \cdot 9)(10 \cdot 9 \cdot 8)(8 \cdot 7 \cdot 6) = 237,105,792,000$
8. (a)  $\binom{52}{5} = 2,598,960$   
 (b)  $\binom{4}{3} \binom{4}{2} = 24$

### 4.4 Combinatorics with Indistinguishable Objects

1. The frequency of letters in each anagram are given, followed by an application of Theorem 4.4.2.
- (a) c, 1; o, 1; m, 2; i, 1; t, 2; e, 2; s, 1;  $\frac{10!}{2!2!2!} = 453,600$   
 (b) m, 2; e, 3; a, 1; s, 2; u, 1; r, 1; n, 1; t, 1;  $\frac{12!}{2!2!3!} = 19,958,400$   
 (c) t, 1; h, 1; e, 3; p, 1; r, 3; o, 2; f, 1, a, 1; d, 1, s, 1;  $\frac{15!}{2!3!3!} = 18,162,144,000$   
 (d) r, 1; e, 3; v, 1; i, 3; s, 1; d, 2; t, 1; o, 1; n, 1;  $\frac{14!}{2!3!3!} = 1,210,809,600$   
 (e) t, 2; h, 1; e, 4; o, 4; d, 1; r, 2; s, 1, v, 1; l, 1;  $\frac{17!}{2!2!4!4!} = 154,378,224,000$   
 (f) t, 3; r, 2; u, 1; s, 3; w, 1; o, 1; h, 1; i, 1; n, 1; e, 1;  $\frac{15!}{2!3!3!} = 18,162,144,000$   
 (g) w, 1; i, 2; l, 2; a, 3; m, 1; s, 2; h, 1; k, 1; e, 3; p, 1; r, 1;  $\frac{18!}{2!2!2!3!3!} = 22,230,464,256,000$   
 (h) t, 4; h, 2; e, 6; u, 3; n, 1; i, 3; d, 1; s, 4; a, 2; b, 1; r, 2; o, 1; f, 2;  $\frac{32!}{2!2!2!2!3!3!4!4!6!} = 1,101,524,811,141,375,548,928,000,000$
3. (a) There are  $\frac{15!}{6!6!3!} = 420,420$  distinguishable permutations of the six indistinguishable nut crunch bars, six indistinguishable chocolate bars, and three indistinguishable toffee bars, and thus there are 420,420 distinguishable ways to distribute the bars to a row of 15 students.
- (b) We have three tasks: distributing the nut crunch bars, the chocolate bars, and the toffee bars. Distributing six nut crunch bars to 15 students can be done in  $\binom{6+15-1}{6} = \binom{20}{6} = 38,760$  ways. Distributing six chocolate bars to 15 students can also be done in 38,760 ways. Finally, distributing three toffee bars to 15 students can be done in  $\binom{3+15-1}{3} = \binom{17}{3} = 680$  ways. By the Fundamental Principle of Counting, the 15 bars can be distributed to 15 students in  $(38,760)(38,760)(680) = 1,021,589,568,000$  ways.
7. (a) Each child has 7 choices.  $7^5 = 16,807$ .

(b) Five drinks can be placed in 11 drink-or-divider slots in  $\binom{11}{5} = 462$  ways.

8. Let F represent a football toss ticket, C a cakewalk ticket, and G a miniature golf ticket. The number of indistinguishable arrangements of the tickets

F F F C C C C G G G G G

is  $\frac{13!}{3!4!6!} = 60,060$ .

11. (a) The budget increase will be divided into 100 equal one-percent increments which will be distributed among three areas. This may be done in  $\binom{100+3-1}{100} = \binom{102}{2} = 5151$  ways.
- (b) After 15% increases are distributed to each of the three areas, there remain 55 one-percent increments to be divided among the three areas. This can be done in  $\binom{55+3-1}{2} = \binom{57}{2} = 1596$  ways.
- (c) After a 50% increase is allotted for salaries, the remaining 50 one-percent increments can be distributed to the three areas in  $\binom{50+3-1}{2} = \binom{52}{2} = 1326$  ways.

## 4.5 Probability

2. (a)  $\frac{1}{435}$ . There are  $\binom{30}{2} = 435$  possible pairs, and {Sarah, Becky} constitute only one such pair.
- (b)  $\frac{28}{435}$ . Of the  $\binom{30}{2} = 435$  possible pairs, there are 28 of form {Sarah,  $x$ } where  $x$  is a member other than Sarah or Becky.
- (c)  $\frac{57}{435}$ . Of the  $\binom{30}{2} = 435$  possible pairs, there are 28 in which Sarah is selected but not Becky (see (b)) and likewise, 28 in which Becky is selected but not Sarah. Together with one pair in which both are selected, this gives  $28 + 28 + 1 = 57$  pairs including Sarah or Becky.
- (d)  $\frac{378}{435}$ . From (c), 57 of the 435 pairs include Sarah or Becky, so the remaining  $435 - 57 = 378$  pairs include neither Sarah nor Becky.
4. Note that the sample space  $S$  consists of  $\binom{52}{5} = 2,598,960$  possible 5-card hands.

- (a)  $\frac{4 \binom{13}{5}}{\binom{52}{5}} = \frac{5148}{2,598,960} \approx 0.00198079$ . There are 4 possible suits, and once the suit is selected,  $\binom{13}{5}$  possible hands within that suit.

(b)  $\frac{\binom{13}{1} \binom{4}{4} \binom{48}{1}}{\binom{52}{5}} = \frac{624}{2,598,960} \approx 0.000240096$ . From 13 kinds, we choose 1. From

the 4 of this kind, we choose all 4, and from the 48 cards not of this kind, we choose 1.

(c)  $\frac{13 \cdot 12 \binom{4}{3} \binom{4}{2}}{\binom{52}{5}} = \frac{3744}{2,598,960} \approx 0.00144058$ . There are 13 ways to choose the

kind to get 3 of, and  $\binom{4}{3}$  ways to select the three from 4 cards of this kind, and

There are 12 ways to choose the kind to get 2 of, and  $\binom{4}{2}$  ways to select the two from 4 cards of this kind.

(d)  $\frac{\binom{10}{1} \binom{4}{1} \binom{4}{1} \binom{4}{1} \binom{4}{1} \binom{4}{1}}{\binom{52}{5}} = \frac{10240}{2,598,960} \approx 0.00394004$ . There are 10 choices

(A, 2, 3, ..., 10) for the lowest card in the straight. This determines which 5 values will be in the straight. There are 4 cards of each of these values and we wish to choose 1 of each.

We have found that there are

624 ways to get a four of a kind

3744 ways to get a full house

5148 ways to get a flush, and

10240 ways to get a straight.

Thus, these hands are here listed in order from rarest to most common, so four of a kind beats a full house, a full house beat a flush, and a flush beats a straight.

8.  $\frac{30}{1200} = \frac{1}{40}$ . Since  $1200 = 2^4 3^1 5^2$ , any positive divisor of 1200 has form  $2^r 3^s 5^t$  where  $0 \leq r \leq 4$ ,  $0 \leq s \leq 1$ , and  $0 \leq t \leq 2$ . As there are 5 choices for  $r$ , 2 choices for  $s$ , and 3 choices for  $t$ , there are  $(5)(2)(3) = 30$  divisors of 1200 in the set  $\{1, 2, \dots, 1200\}$ .

10. As seen in Example 4.5.4, the sample space contains  $\binom{19}{7} = 50,388$  elements.

(b)  $\frac{19,305}{50,388}$ . There are  $\binom{13}{5}$  ways to select the five colors. Since there must be one gumball of each color, this accounts for 5 gumballs. The remaining 2 may be distributed among the 5 colors (requiring 4 dividers) in  $\binom{2+4}{2} = \binom{6}{2} = 15$  ways. Thus, there are  $1287 \cdot 15 = 19,305$  assortments with exactly 5 colors.

(c)  $\frac{19,071}{50,388}$ . No more than 4 colors means 1 color, 2 colors, 3 colors, or 4 colors. In (a) we found that there are 13 assortments with one color, and in Example 4.5.4 we found that there are 14300 assortments with exactly 4 colors.

*Assortments with exactly 2 colors:* There are  $\binom{13}{2} = 78$  ways to choose the 2 colors. Since there must be one gumball of each color, this accounts for 2 gumballs. The remaining 5 may be distributed among the 2 colors (requiring 1 divider) in  $\binom{5+1}{5} = \binom{6}{5} = 6$  ways. Thus, there are  $78 \cdot 6 = 468$  assortments

with exactly 2 colors.

*Assortments with exactly 3 colors:* There are  $\binom{13}{3} = 286$  ways to choose the 3 colors. Since there must be one gumball of each color, this accounts for 3 gumballs. The remaining 4 may be distributed among the 3 colors (requiring 2 dividers) in  $\binom{4+2}{4} = \binom{6}{4} = 15$  ways. Thus, there are  $286 \cdot 15 = 4290$  assortments with exactly 3 colors.

Combining our results, there are  $13 + 468 + 4290 + 14300 = 19,071$  assortments with no more than 4 colors.



# Chapter 5

## Relations

### 5.1 Relations

3. If  $A = \{1, 2, 3\}$  and  $R = \{(1, 2), (2, 1), (1, 3)\}$ , then  $R$  is not symmetric (for  $(1, 3) \in R$  but  $(3, 1) \notin R$ ) and is not antisymmetric (for  $(1, 2) \in R$  and  $(2, 1) \in R$ , but  $1 \neq 2$ ). This shows that neither implication holds.
5. The ordered pairs given with some negative answers suggest points at which the property in question fails.

Relation	Domain	Range	Refl.	Sym.	Antisym.	Trans.
(a) S	$\{1, 3, 5\}$	$\{3, 5\}$	No	No	No	No: $(3, 5), (5, 3)$
(b) R	$\mathbb{N}$	$\mathbb{N} \setminus \{1\}$	No $(1, 1)$	No $(1, 3)$	No $(7, 8)$	No $(100, 15), (15, 5)$
(c) T	$\{0, 4, 7\}$	$\{0, 4, 7\}$	Yes	Yes	No $(0, 7)$	Yes
(d) U	$\mathbb{Z} \setminus \{0\}$	$\mathbb{Z} \setminus \{0\}$	No $(0, 0)$	Yes	No $(1, 2)$	Yes
(e) P	$\mathbb{Z} \setminus \{0\}$	$\mathbb{Z} \setminus \{0\}$	No $(5, 5)$	Yes	No $(3, 7)$	No $(6, 5), (5, 2)$

7. (a)  $S \times S$  has 9 elements, and thus has  $2^9 = 512$  subsets. Relations on  $S$  are subsets of  $S \times S$ , so there are 512 relations on  $S$ .
- (b) Every reflexive relation on  $S$  has form  $\{(1, 1), (2, 2), (3, 3)\} \cup C$  where  $C$  is a subset of the remaining six elements of  $S \times S$ . There are  $2^6$  such subsets  $C$ , and thus  $2^6 = 64$  reflexive relations on  $S$ .
- (c) The relations described are of form

$$\{(1, 1), (2, 2), (3, 3), (1, 2), (1, 3), (2, 3)\} \cup C$$

where  $C$  is a subset of the remaining three elements of  $S \times S$ , that is, where  $C \subseteq \{(2, 1), (3, 1), (3, 2)\}$ . There are  $2^3 = 8$  such subsets. They are  $C_1 = \emptyset$ ,  $C_2 = \{(2, 1)\}$ ,  $C_3 = \{(3, 1)\}$ ,  $C_4 = \{(3, 2)\}$ ,  $C_5 = \{(2, 1), (3, 1)\}$ ,  $C_6 = \{(2, 1), (3, 2)\}$ ,

$C_7 = \{(3, 1), (3, 2)\}$ , and  $C_8 = \{(2, 1), (3, 1), (3, 2)\}$ . Now the 8 such relations are given by  $R_i = \{(1, 1), (2, 2), (3, 3), (1, 2), (1, 3), (2, 3)\} \cup C_i$  for  $i = 1, \dots, 8$ .

(d) Each relation  $R_i$  ( $i = 1, \dots, 8$ ) has  $\{1, 2, 3\}$  as domain and range.

(e)

	Refl.	Sym.	Antisym.	Trans.
$R_1$	Yes	No	Yes	Yes
$R_2$	Yes	No	No	Yes
$R_3$	Yes	No	No	No
$R_4$	Yes	No	No	Yes
$R_5$	Yes	No	No	No
$R_6$	Yes	No	No	No
$R_7$	Yes	No	No	No
$R_8$	Yes	Yes	No	Yes

10. (a) i.  $\{(2,3), (2,1), (3,5), (4,4)\}$   
 ii.  $\{(1,3), (3,5), (5,4), (5,2)\}$   
 iii.  $\{(5,4), (5,2)\}$
- (b) The graph of  $S$  is a parabola in  $\mathbb{R}^2$  with vertex at the origin and having the  $y$ -axis as axis of symmetry. The graph of  $S|_{[0, \infty)}$  is the right half of that parabola, including the vertex  $(0, 0)$ .
- (c) i.  $\{(1,1), (1,2), (1,3), (1,4), (1,5), (1,6), (2,2), (2,3), (2,4), (2,5), (2,6), (3,3), (3,4), (3,5), (3,6)\}$   
 ii.  $\{(2,2), (2,3), (2,4), (2,5), (2,6), (4,4), (4,5), (4,6)\}$   
 iii.  $\{(6,6)\}$
13. (a)  $R_1 \circ R_n = R_1$  and  $R_n \circ R_1 = R_1$  for all  $n \in \{1, 2, \dots, 16\}$ .  
 (b)  $R_8 \circ R_n = R_n$  and  $R_n \circ R_8 = R_n$  for all  $n \in \{1, 2, \dots, 16\}$ .  
 (c) For all  $n \in \{1, 2, \dots, 16\}$ ,  $R_{16} \circ R_n$  is the largest relation on  $\{1, 2\}$  having the same domain as  $R_n$ , and  $R_n \circ R_{16}$  is the largest relation on  $\{1, 2\}$  having the same range as  $R_n$ .

## 5.2 Equivalence Relations

5. In measuring angles in radian measure, angles  $x$  and  $y$  are coterminal if and only if  $x = y + 2\pi n$  for some  $n \in \mathbb{Z}$ , that is, if and only if  $x \equiv y \pmod{2\pi}$ .
8. (a)  $\#$  is not reflexive and not transitive, and thus is not an equivalence relation.  
 (b)  $\Delta$  is an equivalence relation. The equivalence classes are  $\{\emptyset\}$ ,  $\{\{1\}, \{2\}, \{3\}\}$ ,  $\{\{1, 2\}, \{1, 3\}, \{2, 3\}\}$ , and  $\{\{1, 2, 3\}\}$ .  
 (c)  $*$  is an equivalence relation. The equivalence classes are  $\{\emptyset, \{2\}\}$ ,  $\{\{1\}, \{1, 2\}\}$ ,  $\{\{3\}, \{2, 3\}\}$ , and  $\{\{1, 3\}, \{1, 2, 3\}\}$ .  
 (d)  $\approx$  is not symmetric and is thus not an equivalence relation.

### 5.3 Partial Orders

3. (a) The relation is a partial order.
- (b)  $\sqsubseteq$  is reflexive since  $x \leq x^2 \forall x \in \mathbb{N}$ .  $\sqsubseteq$  is not antisymmetric. For example,  $7 \sqsubseteq 8$  and  $8 \sqsubseteq 7$ , yet  $7 \neq 8$ .  $\sqsubseteq$  is not transitive. For example,  $15 \sqsubseteq 4$  and  $4 \sqsubseteq 2$ , but  $15 \not\sqsubseteq 2$ . Thus  $\sqsubseteq$  is not a partial order.
- (c) The relation  $\ll$  is reflexive, but is neither antisymmetric ( $3 \ll 4$  and  $4 \ll 3$  but  $3 \neq 4$ , for example) nor transitive ( $6 \ll 4$  and  $4 \ll 2$ , but  $6 \not\ll 2$ , for example). Thus,  $\ll$  is not a partial order.
5.  $(\mathcal{P}(S), \subseteq)$  is totally ordered if and only if  $|S| \leq 1$ .

If  $|S| = 0$ , then  $\mathcal{P}(S) = \{\emptyset\}$ , a one-element collection totally ordered by inclusion. If  $|S| = 1$ , then  $\mathcal{P}(S) = \{\emptyset, S\}$ , a two-element collection totally ordered by inclusion. If  $|S| \geq 2$ , then there exist distinct elements  $a, b \in S$ , and  $\{a\}, \{b\} \in \mathcal{P}(S)$  but  $\{a\} \not\subseteq \{b\}$  and  $\{b\} \not\subseteq \{a\}$ . Thus, if  $|S| \geq 2$ , then  $(\mathcal{P}(S), \subseteq)$  is not totally ordered.

10. (a) Yes. The maximum element of  $S$  is an upper bound of  $C$ .
- (b) No. Let  $P = [0, 1) \cup (2, 3]$  in  $\mathbb{R}$  with the usual order. The upper bounds of  $C = [0, 1)$  are precisely the points of  $(2, 3]$ . Thus,  $C = [0, 1)$  has upper bounds, but no least upper bound.
- (c) No. Let  $P = \{\{a\}, \{b\}, \{a, b, c\}, \{a, b, d\}, \{a, b, c, d\}\}$  ordered by inclusion, and let  $C = \{\{a\}, \{b\}\}$ . Now the set of upper bounds of  $C$  is  $UB = \{\{a, b, c\}, \{a, b, d\}, \{a, b, c, d\}\}$ . Now since  $UB$  has no minimum element,  $C$  has no least upper bound.
- (d) Yes. If  $C$  has a least upper bound, then the set  $UB$  of upper bounds of  $C$  is nonempty, and if  $UB$  has a minimum element, it must be unique. (See Theorem 5.3.6.)
11.  $a \prec b$  if and only if there is a line from  $a$  upward to  $b$  in the Hasse diagram for the poset.
- (a)  $\{2\} \prec \{2, 3\}$ ;  $\{3\} \prec \{2, 3\}$ ;  $\{2, 3\} \prec \{2, 3, 4\}$ ;  $\{4, 5\} \prec \{4, 5, 6\}$ .

17. (a) Yes. If each  $P_i$  has a maximum element  $m_i$ , then  $(m_i)_{i \in I}$  is the maximum element in  $P$ , for given any  $(x_i)_{i \in I} \in P$ , we have  $x_i \leq_i m_i \forall i \in I$ , so by the definition of the product order,  $(x_i)_{i \in I} \leq (m_i)_{i \in I}$ .
- (b) Yes. Suppose  $(m_i)_{i \in I}$  is the maximum element in  $P$ . Then  $\forall (x_i)_{i \in I} \in P$ , we have  $(x_i)_{i \in I} \leq (m_i)_{i \in I}$  and hence  $x_i \leq_i m_i \forall i \in I$ . We claim  $m_{i_0}$  is the maximum element of  $P_{i_0}$ . Suppose  $x \in P_{i_0}$ . Define  $(a_i)_{i \in I} \in P$  by  $a_i = m_i$  for  $i \in I \setminus \{i_0\}$  and  $a_{i_0} = x$ . Now  $(a_i)_{i \in I} \leq (m_i)_{i \in I}$  implies  $x = a_{i_0} \leq m_{i_0}$ . Since  $x \in P_{i_0}$  was arbitrary, this shows that  $m_{i_0}$  is maximum in  $P_{i_0}$ .
- (c) If, for all  $i \in I$ ,  $m_i$  is a maximal element in  $P_i$ , then  $(m_i)_{i \in I}$  is maximal in  $P$ , for if  $(x_i)_{i \in I} \in P$  with  $(x_i)_{i \in I} \geq (m_i)_{i \in I}$ , then  $m_i \leq_i x_i \forall i \in I$ . Since  $m_i$  is maximal in  $P_i$ , this implies  $m_i = x_i \forall i \in I$ , so  $(x_i)_{i \in I} = (m_i)_{i \in I}$ , and  $(m_i)_{i \in I}$  is maximal in  $P$ .

If  $(m_i)_{i \in I}$  is maximal in  $P$ , then  $m_i$  is maximal in  $P_i \forall i \in I$ , for if not, there exists  $i_0 \in I$  and  $x \in P_{i_0}$  with  $m_{i_0} <_{i_0} x$ . Now  $(a_i)_{i \in I} \in P$  defined by  $a_i = m_i \forall i \in I \setminus \{i_0\}$  and  $a_{i_0} = x$  is a point of  $P$  strictly larger than  $(m_i)_{i \in I}$ , contrary to the maximality of  $(m_i)_{i \in I}$ .

## 5.4 Quotient Spaces

3. (a) For  $0 < \epsilon < 2$ , consider the line  $l_\epsilon$  through the point  $(2 - \epsilon, 200) \in P(0, 100)$  and the point  $(2, 202 - \epsilon) \in P(1, 100)$ . The slope of  $l_\epsilon$  is  $\frac{2-\epsilon}{\epsilon}$ , and these slopes range from 0 to  $\infty$  as  $\epsilon$  ranges from 2 to 0. With arbitrary positive slopes allowed, we may find a line of this form passing through the point  $(8, 2n + 1) \in P(4, n)$  for every integer  $n \geq 100$ . Thus,  $P(0, 100)$ ,  $P(1, 100)$  and  $P(4, n)$  are collinear for all integers  $n \geq 100$ . To show that  $P(0, 100)$ ,  $P(1, 100)$  and  $P(4, n)$  are collinear for all integers  $n < 100$ , for  $0 < \epsilon < 2$ , consider the lines through  $(2 + \epsilon, 200) \in P(1, 100)$  and  $(2 - \epsilon, 202 - \epsilon) \in P(0, 100)$ . Such a line has slope  $\frac{2-\epsilon}{-2\epsilon}$ , and as  $\epsilon$  ranges from 0 to 2, these slopes range from  $-\infty$  to 0. Thus, there exists such a line through  $P(0, 100)$ ,  $P(1, 100)$  and  $P(4, n)$  for all integers  $n < 100$ .
  - (c) No. Let  $\mathcal{L}_1$  be the collinear set  $\{P(0, 100), P(2, 102)\}$ , let  $\mathcal{L}_2$  be the collinear set  $\{P(0, 100), P(2, 100)\}$ , and let  $\mathcal{L}_3$  be the collinear set  $\{P(0, 100), P(2, 98)\}$ . It is easy to see that (i) there is a line  $l_1$  of slope  $m_1$  which illuminates all the pixels of  $\mathcal{L}_1$  if and only if  $m_1 \in (\frac{1}{3}, 3)$ , (ii) there is a line  $l_2$  of slope  $m_2$  which illuminates all the pixels of  $\mathcal{L}_2$  if and only if  $m_2 \in (-1, 1)$ , and (iii) there is a line  $l_3$  of slope  $m_3$  which illuminates all the pixels of  $\mathcal{L}_3$  if and only if  $m_3 \in (-3, \frac{1}{3})$ .  
Now  $\mathcal{L}_1$  is parallel to  $\mathcal{L}_2$  since there exist parallel lines  $l_i$  of slope  $m_i = \frac{2}{3}$  illuminating all the pixels of  $\mathcal{L}_i$  ( $i = 1, 2$ ), and  $\mathcal{L}_2$  is parallel to  $\mathcal{L}_3$  since there exist parallel lines  $l_i$  of slope  $m_i = \frac{-2}{3}$  illuminating all the pixels of  $\mathcal{L}_i$  ( $i = 2, 3$ ). However, no line  $l_1$  illuminating all the pixels of  $\mathcal{L}_1$  can be parallel to any line  $l_3$  illuminating all the pixels of  $\mathcal{L}_3$ , since  $m_1 \in (\frac{1}{3}, 3)$  and  $m_3 \in (-3, \frac{1}{3})$  imply  $m_1 \neq m_3$ .
6. Suppose  $[a] = [a_1]$  and  $[b] = [b_1]$  in  $\mathbb{Z}/n$ . Then  $a = a_1 + kn$  and  $b = b_1 + jn$  for some  $j, k \in \mathbb{Z}$ . Thus,  $ab = (a_1 + kn)(b_1 + jn) = a_1b_1 + n(a_1j + kb_1 + knj)$  where  $a_1j + kb_1 + knj \in \mathbb{Z}$ , so  $[ab] = [a_1b_1]$  in  $\mathbb{Z}/n$ . Thus,  $[a] \times [b] = [a_1] \times [b_1]$  whenever  $[a] = [a_1]$  and  $[b] = [b_1]$ , so the operation  $\times$  is well defined.
  7. (a) Since  $[3] \times [5] = [15] = [1]$  in  $\mathbb{Z}/7$ ,  $[5]$  is the multiplicative inverse of  $[3]$  in  $\mathbb{Z}/7$ .  
(b) Since  $[3] \times [2] = [6] = [1]$  in  $\mathbb{Z}/5$ ,  $[2]$  is the multiplicative inverse of  $[3]$  in  $\mathbb{Z}/5$ .  
(c) Since  $[3] \times [3] = [9] = [1]$  in  $\mathbb{Z}/4$ ,  $[3]$  is the multiplicative inverse of  $[3]$  in  $\mathbb{Z}/4$ .  
(d) In  $\mathbb{Z}/6$ , We have  $[3] \times [0] = [0]$ ,  $[3] \times [1] = [3]$ ,  $[3] \times [2] = [0]$ ,  $[3] \times [3] = [3]$ ,  $[3] \times [4] = [0]$ , and  $[3] \times [5] = [3]$ . Thus, there is no  $[n] \in \mathbb{Z}/6$  with  $[3] \times [n] = [1]$ , so  $[3]$  has no multiplicative inverse in  $\mathbb{Z}/6$ .
11. (b) Reflexive: For any triangle  $t_1$ ,  $t_1$  has an angle whose measure is greater than or equal to that of every angle in  $t_1$ . Transitive: If  $t_1$  has an angle whose measure is greater than or equal to that of every angle in  $t_2$  and  $t_2$  has an angle whose

measure is greater than or equal to that of every angle in  $t_3$ , then  $t_1$  has an angle whose measure is greater than or equal to that of every angle in  $t_3$ .

Now  $t_1 \preceq t_2$  if and only if the largest angle in  $t_2$  is greater than or equal to the largest angle in  $t_1$ . Thus,  $t_1 \sim t_2$  if and only if the largest angle in  $t_1$  has the same measure as the largest angle in  $t_2$ . The resulting partial order on equivalence classes is a total order.

14. (a) For any  $a \in \emptyset$ , we have  $b \leq a \Rightarrow b \in \emptyset$  vacuously, so  $\emptyset \in \mathcal{T}$ . For  $a, b \in S$ , clearly  $a \in S$  and  $b \leq a$  implies  $b \in S$ , so  $S \in \mathcal{T}$ .
- (b) Suppose  $D_1, D_2, \dots, D_n \in \mathcal{T}$ ,  $a \in D_1 \cap D_2 \cap \dots \cap D_n$ , and  $b \leq a$ . Since  $D_1, D_2, \dots, D_n$  are each decreasing,  $b \in D_i$  ( $i = 1, 2, \dots, n$ ), so  $b \in D_1 \cap \dots \cap D_n$ . Thus,  $D_1 \cap \dots \cap D_n \in \mathcal{T}$ .
- (c) Suppose  $J$  is an arbitrary index set (finite or infinite),  $D_j \in \mathcal{T} \forall j \in J$ ,  $a \in \cup_{j \in J} D_j$ , and  $b \leq a$ . Now  $a \in \cup_{j \in J} D_j \Rightarrow \exists j_0 \in J$  such that  $a \in D_{j_0}$ . Now  $D_{j_0}$  decreasing implies  $b \in D_{j_0}$ , and thus  $b \in \cup_{j \in J} D_j$ . Thus  $\cup_{j \in J} D_j \in \mathcal{T}$ .



## Chapter 6

# Functions and Cardinality

### 6.1 Functions

1. (a) The graph represents a function from  $\mathbb{R}$  to  $\mathbb{R}$  if and only if every vertical line intersects the graph in exactly one point.  
If the graph represents a function  $f$ , the line  $x = a$  intersects the graph in exactly one point  $(a, f(a))$ . If every vertical line  $x = a$  intersects the graph in exactly one point, let this point be  $(a, f(a))$  and this defines a function from  $\mathbb{R}$  to  $\mathbb{R}$ .
- (b) The graph represents a function from a *subset* of  $\mathbb{R}$  to  $\mathbb{R}$  if and only if every vertical line intersects the graph in no more than one point.  
If the graph represents a function  $f : S \rightarrow \mathbb{R}$  where  $S \subseteq \mathbb{R}$ , then  $x = a$  intersects the graph in exactly one point  $(a, f(a))$  for each  $a \in S$ , and in no points for  $a \notin S$ . Conversely, if every vertical line intersects the graph  $G$  in no more than one point, the graph represents  $f : S \rightarrow \mathbb{R}$  where  $S = \{a \in \mathbb{R} \mid x = a \text{ intersects } G\}$  and  $(a, f(a))$  is the intersection of  $x = a$  with  $G$ .
8. We will show  $f$  is not one-to-one if and only if there is a horizontal line intersecting the graph more than once. If  $f$  is not one-to-one, then there exist  $a \neq b$  such that  $f(a) = f(b)$ , so the horizontal line  $y = f(a)$  intersects the graph of  $f$  in (at least) two points, namely  $(a, f(a))$  and  $(b, f(b))$ . Conversely, if a horizontal line  $y = c$  intersects the graph of  $f$  in two points  $(a, c)$  and  $(b, c)$  with  $a \neq b$ , then  $f(a) = f(b) = c$ , so  $f$  is not one-to-one.
10. We will describe  $f : \{1, 2, 3\} \rightarrow \{1, 2, 3\}$  by listing the ordered triple  $(f(1), f(2), f(3))$ . Now the functions  $(1, 2, 3)$ ,  $(1, 3, 2)$ ,  $(2, 1, 3)$ , and  $(3, 2, 1)$  are the only ones with  $f \circ f = id$ . If  $g \circ f$  is one-to-one, then  $f$  must be, so all functions with  $f \circ f = id$  are one-to-one. (In fact, if  $f \circ f = id$ , then  $f = f^{-1}$ , so  $f$  is invertible and is thus one-to-one and onto.)
13. (a)  $\sim_f$  is reflexive since  $f(a) = f(a) \forall a \in A$ .  $\sim_f$  is symmetric since  $f(a) = f(b) \Rightarrow f(b) = f(a)$ .  $\sim_f$  is transitive since  $f(a) = f(b)$  and  $f(b) = f(c)$  imply  $f(a) = f(c)$ .

- (b) If  $f$  is injective, then  $a \sim_f b \iff f(a) = f(b) \iff a = b$ , so  $\sim_f$  is  $\Delta_A$ . If  $\sim_f$  is  $\Delta_A$  and  $f(a) = f(b)$ , then  $a \sim_f b$ , so  $(a, b) \in \Delta_A$ , so  $a = b$ , and thus  $f$  is injective.
17. (a) Let  $S = [-1, 0)$ ,  $T = (0, 1]$ , and  $A = S \cup T$  have the usual order from  $\mathbb{R}$ . Consider  $f : A \rightarrow \mathbb{R}$  (where  $\mathbb{R}$  has the usual order) defined by  $f(x) = x + 1$  if  $x < 0$  and  $f(x) = x$  if  $x > 0$ . Now  $f$  is increasing on  $S$  and on  $T$ , but not on  $A = S \cup T$  since, for example,  $\frac{-1}{4} \leq \frac{1}{4}$  but  $f(\frac{-1}{4}) = \frac{3}{4} \not\leq \frac{1}{4} = f(\frac{1}{4})$ .
- (b) Suppose  $f$  is increasing on  $S$  and on  $T$ .  
 $S \cap T \neq \emptyset$  is not necessary for  $f$  to be increasing on  $A = S \cup T$ : consider  $f : S \cup T \rightarrow \mathbb{R}$  given by  $f(x) = x$ , where  $S = [-1, 0)$  and  $T = (0, 1]$  are subsets of  $\mathbb{R}$  with the usual order.  
 $S \cap T \neq \emptyset$  is not sufficient for  $f$  to be increasing on  $A = S \cup T$ : consider  $A = S \cup T$  where  $S = \{\{1\}, \{1, 2\}\}$  and  $T = \{\{1\}, \{1, 2, 3\}\}$  with set inclusion as the order. Define  $f : A \rightarrow \mathbb{N}$  (where  $\mathbb{N}$  has the usual order) by  $f(\{1\}) = 1$ ,  $f(\{1, 2\}) = 5$ , and  $f(\{1, 2, 3\}) = 2$ . Now  $f$  is increasing on  $S$  and on  $T$  but not on  $A = S \cup T$  since  $\{1, 2\} \subseteq \{1, 2, 3\}$  but  $f(\{1, 2\}) = 5 \not\leq 2 = f(\{1, 2, 3\})$ .
20. (a) Observe that  $f \preceq g$  if and only if  $\frac{f(1)-f(0)}{1-0} \leq \frac{g(1)-g(0)}{1-0}$ , that is, if and only if the slope of  $f$  is less than or equal to the slope of  $g$ . Clearly  $f \preceq f$  for any  $f \in \mathcal{F}$  since the slope of  $f$  is less than or equal to the slope of  $f$ . If  $f \preceq g$  and  $g \preceq h$ , then the slope of  $f$  is less than or equal to that of  $g$ , and the slope of  $g$  is less than or equal to that of  $h$ , so the slope of  $f$  is less than or equal to that of  $h$ , so  $f \preceq h$ , and thus  $\preceq$  is transitive.

## 6.2 Inverse Relations and Inverse Functions

2. (a) The inverse relation  $\{(2,1), (1,2), (4,3), (3,4)\}$  is a function.  
 (b) The inverse relation  $\{(1,1), (1,3), (1,2), (1,4)\}$  is not a function.  
 (c) The inverse relation  $\{(1,1), (2,1), (3,1), (4,1)\}$  is a function.  
 (d) The inverse relation  $\{(3,1), (4,2), (3,3), (3,4)\}$  is not a function.  
 (e) The inverse relation  $\{(3,1), (1,2), (2,3)\}$  is not a function on  $\{1,2,3,4\}$ .
3. (a)  $f^{-1}(x) = \frac{3x+1}{5}$   
 (f)  $f^{-1}(x) = \frac{1+2x}{x-2}$
4. (b)  $T = [3, \infty)$ ,  $f|_{T^{-1}}(x) = \sqrt{x+4} + 3$ , or  $T' = (-\infty, 3]$ ,  $f|_{T'^{-1}}(x) = -\sqrt{x+4} + 3$   
 (d)  $T = \mathcal{P}(\{2n-1|n \in \mathbb{N}\} \cup \{4n|n \in \mathbb{N}\})$  and  $f|_{T^{-1}}(S) = \{s \in S | s \text{ is odd}\} \cup \{2s | s \in S \text{ and } s \text{ is even}\}$ .  
 (e)  $T = \mathbb{N} \cup \{0\}$  and  $f|_{T^{-1}}(x) = x - 1$ , or  $T' = \{2n-1|n \in \mathbb{N}\} \cup \{2-2n|n \in \mathbb{N}\}$  and  $f|_{T'^{-1}}(x) = x - 1$  if  $x$  is even and  $f|_{T'^{-1}}(x) = 1 - x$  if  $x$  is odd.  
 (g)  $T = \{\emptyset, \{1\}, \{1, 2\}, \{1, 2, 3\}, \dots\}$  and  $f|_{T^{-1}}(x) = \{1, 2, \dots, x\}$  if  $x \neq 0$ ;  $f|_{T^{-1}}(0) = \emptyset$ .



(h)  $T = A$ ,  $f(T) = \{[x, y] \subseteq \mathbb{R} \mid x \leq y + 4\}$ , and  $f|_T^{-1}([x, y]) = [x + 1, y - 3]$ .

12. Suppose  $f : A \rightarrow B$  and  $C \subseteq D \subseteq B$ . Now

$$\begin{aligned} x \in f^{-1}(C) &\Rightarrow f(x) \in C \subseteq D \\ &\Rightarrow f(x) \in D \\ &\Rightarrow x \in f^{-1}(D). \end{aligned}$$

Thus,  $f^{-1}(C) \subseteq f^{-1}(D)$ .

The converse fails. If  $f(x) = x^2$ , then  $f^{-1}([-10, 1]) \subseteq f^{-1}([-3, 1])$  but  $[-10, 1] \not\subseteq [-3, 1]$ .

19. (d)  $f^{-1}(5)$  contains numbers of the following types:

TYPE	form/choices for each digit	total number
5 odd digits:		
$\underbrace{5}$	$\underbrace{5}$	$5^5 = 3125$
odd	odd	
3 odd digits, 1 even:		
$\underbrace{4}$	$\underbrace{5}$	$4 \cdot 5^3 = 500$
even $\neq 0$	odd	
$\underbrace{5}$	$\underbrace{5}$	$5^4 = 625$
odd	even	
$\underbrace{5}$	$\underbrace{5}$	$5^4 = 625$
odd	odd	
$\underbrace{5}$	$\underbrace{5}$	$5^4 = 625$
odd	odd	
1 odd digit, 2 evens:		
$\underbrace{4}$	$\underbrace{5}$	$4 \cdot 5^2 = 100$
even $\neq 0$	odd	
$\underbrace{4}$	$\underbrace{5}$	$4 \cdot 5^2 = 100$
even $\neq 0$	even	
$\underbrace{5}$	$\underbrace{5}$	$5^3 = 125$
odd	even	

Adding the numbers in the right column above gives  $|f^{-1}(5)| = 5825$ .

### 6.3 Cardinality of Infinite Sets

6. A function  $f : \{1, 2\} \rightarrow \mathbb{N}$  is completely characterized by the ordered pair  $(f(1), f(2)) \in \mathbb{N} \times \mathbb{N}$ . This gives a bijection between the set of all functions  $f : \{1, 2\} \rightarrow \mathbb{N}$  and  $\mathbb{N} \times \mathbb{N}$ .

Since  $\mathbb{N} \times \mathbb{N}$  is countable, so is the set of all functions  $f : \{1, 2\} \rightarrow \mathbb{N}$ .

7. (a) Suppose  $A_i$  is countable for  $i = 1, 2, \dots, n$ . Consider the products  $A_1, A_1 \times A_2, A_1 \times A_2 \times A_3, \dots, \prod_{i=1}^n A_i$ . Clearly  $A_1$  is countable. If  $\prod_{i=1}^k A_i$  is countable, then  $\prod_{i=1}^{k+1} A_i = (\prod_{i=1}^k A_i) \times A_{k+1}$  is a product of two countable sets and is thus countable. By mathematical induction, it follows that  $\prod_{i=1}^n A_i$  is countable for any  $n$  if each  $A_i$  is countable.
- (b) Corresponding  $f : \{1, 2, \dots, n\} \rightarrow \mathbb{N}$  to  $(f(1), f(2), \dots, f(n)) \in \mathbb{N}^n$  gives a bijection from the set  $B$  to  $\mathbb{N}^n$ . Since the latter set is countable by (a), the former set is also countable.
13. “The smallest natural number that cannot be defined using less than twenty words” is a thirteen-word description of that natural number.

## 6.4 An Order Relation on Cardinal Numbers

3. Each nondegenerate interval in  $\mathbb{R}$  contains a rational number. Since there are only countably many rational numbers, there can be only countably many pairwise disjoint intervals.
4. Let  $C'_r = C_r \cap \{(x, y) \in \mathbb{R}^2 \mid x \geq 0, y \geq 0\}$  be the first quadrant part of the circle  $C_r$ . The projection function  $f : C'_r \rightarrow [0, r]$  defined by  $f((x, y)) = x$  is a bijection. Since  $[0, r]$  is uncountable (see Exercise 3 of Section 6.3), it follows that  $C'_r$  is uncountable. Now  $C'_r = A \cup B \cup C$  where  $A = \{(x, y) \in C'_r \mid x \in \mathbb{Q}\}$ ,  $B = \{(x, y) \in C'_r \mid y \in \mathbb{Q}\}$ , and  $C = \{(x, y) \in C'_r \mid x \notin \mathbb{Q}, y \notin \mathbb{Q}\}$ . Now  $A$  is indexed by a subset of  $\mathbb{Q}$ , so  $A$  is countable. ( $h : A \rightarrow [0, r] \cap \mathbb{Q}$  defined by  $h((x, y)) = x$  is a bijection.) Similarly,  $B$  is countable. Now  $A \cup B \cup C = C'_r$  is uncountable, so  $C$  must be uncountable. Thus,  $C'_r \subseteq C_r$  contains uncountably many points  $(x, y)$  with  $x \notin \mathbb{Q}$  and  $y \notin \mathbb{Q}$ .
9. (a) No.  $\frac{3}{2} = \frac{6}{4}$  but  $g(\frac{3}{2}) = 2 \neq 4 = g(\frac{6}{4})$ .
- (b)  $h$  is well defined since every element of  $\mathbb{Q}^+$  has a unique representation as  $\frac{m}{n}$  where  $m$  and  $n$  are relatively prime natural numbers.  $h$  is not one-to-one since  $h(\frac{1}{4}) = 4 = h(\frac{3}{4})$ .  $h$  is onto, since for any  $n \in \mathbb{N}$ ,  $n = h(\frac{1}{n})$ .
14. Given an algebraic number  $\alpha$ , pick a polynomial  $p(x) = c_0 + c_1x + \dots + c_nx^n$  with integer coefficients  $c_0, \dots, c_n$  such that  $p(\alpha) = 0$ . (In fact, there exists a unique such polynomial of minimal degree such that  $c_0, \dots, c_n$  are relatively prime and  $c_n > 0$ .) Suppose  $\alpha$  is the  $m^{\text{th}}$  zero of  $p(x)$  when the distinct real zeros of  $p(x)$  are listed in increasing order. Map  $\alpha$  to the natural number whose base 12 representation is the sequence of digits

$$m^1 m^2 \dots m^j \pm c_0^1 c_0^2 \dots c_0^{k_0} \pm c_1^1 c_1^2 \dots c_1^{k_1} \dots \pm c_{n-1}^1 c_{n-1}^2 \dots c_{n-1}^{k_{n-1}} \pm c_n^1 c_n^2 \dots c_n^{k_n}$$

where the digits base 12 are  $0, 1, \dots, 9, +$ , and  $-$ ;  $m = m^1 m^2 \dots m^j$  where  $m^1, \dots, m^j$  are the base 10 digits of  $m$ ; and  $c_i = \pm c_i^1 c_i^2 \dots c_i^{k_i}$  where  $c_i^1, \dots, c_i^{k_i}$  are the base 10 digits of  $c_i$  if  $c_i \neq 0$ , and  $c_i = +c_i^1 = +0$  if  $c_i = 0$ . This gives an injection from the set  $A$  of algebraic numbers to  $\mathbb{N}$ , so  $A$  is countable.


## Chapter 7

# Graph Theory

### 7.1 Graphs

2. (a)  $6^5 = 7776$   
(b)  $15^7 = 170859375$   
(c)  $\left(n + \binom{n}{2}\right)^k$ . Any edge has either one endpoint (and there are  $n$  choices for the vertex at which such a loop may be based) or has two endpoints (and there are  $\binom{n}{2}$  choices for the two end points). This gives  $n + \binom{n}{2} = \frac{n^2+n}{2} = T_n$  ways to construct one edge, so there are  $\left(n + \binom{n}{2}\right)^k = T_n^k$  ways to construct  $k$  edges on  $n$  vertices.  
(d) 0  
(e) 120  
(f) There are  $\binom{n}{2}$  possible edges (with distinct endpoints) on  $n$  vertices, and we wish to choose  $k$  of them:  $\binom{\binom{n}{2}}{k}$ .
6. (a)  $e_1, e_2, e_1, e_6, e_{10}$ , for example.  
(c) Impossible. If a walk has distinct vertices, it must have distinct edges, so every path is a trail.  
(e)  $e_7, e_8, e_{11}, e_{10}$ .
10.  $G$  is a connected graph if and only if  $\bigcup_{i \in I} D_i$  is a connected subset of the plane. That is,  $G$  is a connected graph if and only if for every  $a, b \in \bigcup_{i \in I} D_i$ , there is a continuous curve contained in  $\bigcup_{i \in I} D_i \subseteq \mathbb{R}^2$  from  $a$  to  $b$ .
13. By placing a doorway in each edge of the graph, the problem becomes analogous to those of Exercise 12. The associated graph is shown below. Since more than two

vertices have odd degree (namely  $B, D, E,$  and  $F$ ), the graph has no Eulerian trail, so it is impossible to draw a continuous curve bisecting each edge of the original graph.




## 7.2 Matrices, Digraphs, and Relations

2.

$$(a) \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

$$(c) \begin{bmatrix} 0 & 2 & 0 & 1 & 0 \\ 2 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 1 \end{bmatrix}$$




(b) Since the  $(1, 3)$  entry of  $A^3$  is 4, there are four  $(v_1, v_3)$ -walks (i.e.,  $(1, 3)$ -walks) of length three. Referring to the edge labels in (a), they are  $e_1e_1e_2$ ,  $e_1e_2e_3$ ,  $e_2e_3e_3$ , and  $e_2e_4e_2$ .

13. (a)

$$\left[ \begin{array}{cccc|cc} 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 & 1 \\ \hline 1 & 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 1 & 0 & 0 \end{array} \right]$$

(b) The adjacency matrix will be an  $(m+n) \times (m+n)$  matrix containing an  $m \times m$  square of zeros in the upper left corner, an  $n \times n$  square of zeros in the lower right corner, and all other entries are ones.

15. (c)



$$\begin{matrix} & e_1 & e_2 & e_3 & e_4 & e_5 & e_6 \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{bmatrix} 1 & 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 2 \\ 0 & 0 & 0 & 1 & 1 & 0 \end{bmatrix} \end{matrix}$$

17. (c) Let  $P = \prod_{i=1}^n \prod_{j \neq i}^n (1 - a_{ij}a_{ji})$ . Then

$$\begin{aligned}
 P \neq 0 &\iff 1 - a_{ij}a_{ji} \neq 0 \quad \forall i, j \in \{1, 2, \dots, n\}, i \neq j \\
 &\iff a_{ij}a_{ji} = 0 \quad \forall i, j \in \{1, 2, \dots, n\}, i \neq j \\
 &\iff a_{ij} = 0 \text{ or } a_{ji} = 0 \quad \forall i, j \in \{1, 2, \dots, n\}, i \neq j \\
 &\iff [a_{ij} = 1 \text{ and } a_{ji} = 1] \text{ implies } i = j \quad \forall i, j \in \{1, \dots, n\} \\
 &\iff iRj \text{ and } jRi \text{ imply } i = j \quad \forall i, j \in \{1, 2, \dots, n\} \\
 &\iff R \text{ is antisymmetric.}
 \end{aligned}$$

### 7.3 Shortest Paths in Weighted Graphs

6. (b)  $agbhfe$  is a shortest  $(a, e)$ -path, and thus  $bhfe$  must be a shortest  $(b, e)$ -path (for if there were a shorter  $(b, e)$ -path, appending it to  $agb$  would give a shorter  $(a, e)$ -path, contrary to  $agbhfe$  being a shortest  $(a, e)$ -path).
8. (a) The shortest  $(d, v)$ -paths found by the implementation of Dijkstra's algorithm below are  $dcba, deb, dc, d, dgfe, dgf, dg$ .

v	a	b	c	d	e	f	g
	$\infty_d$	$\infty_d$	$1_d$	<b><math>0_d</math></b>	$\infty_d$	$\infty_d$	$2_d$
	$\infty_d$	$5_c$	<b><math>1_d</math></b>	<b><math>0_d</math></b>	$\infty_d$	$4_c$	$2_d$
	$\infty_d$	$5_c$	<b><math>1_d</math></b>	<b><math>0_d</math></b>	$\infty_d$	$3_g$	<b><math>2_d</math></b>
	$12_f$	$5_c$	<b><math>1_d</math></b>	<b><math>0_d</math></b>	$9_f$	<b><math>3_g</math></b>	<b><math>2_d</math></b>
	$8_b$	<b><math>5_c</math></b>	<b><math>1_d</math></b>	<b><math>0_d</math></b>	$9_f$	<b><math>3_g</math></b>	<b><math>2_d</math></b>
	<b><math>8_b</math></b>	<b><math>5_c</math></b>	<b><math>1_d</math></b>	<b><math>0_d</math></b>	$9_f$	<b><math>3_g</math></b>	<b><math>2_d</math></b>
	<b><math>8_b</math></b>	<b><math>5_c</math></b>	<b><math>1_d</math></b>	<b><math>0_d</math></b>	<b><math>9_f</math></b>	<b><math>3_g</math></b>	<b><math>2_d</math></b>

9. (c) The shortest  $(f, v)$ -paths found by the implementation of Dijkstra's algorithm below are  $fa, fb, fgc, fgd, fbe, f, fg, fgch$ .

v	a	b	c	d	e	f	g	h
	$4_f$	$2_f$	$\infty_f$	$\infty_f$	$5_f$	<b><math>0_f</math></b>	$2_f$	$\infty_f$
	$4_f$	<b><math>2_f</math></b>	$10_b$	$\infty_f$	$4_b$	<b><math>0_f</math></b>	$2_f$	$\infty_f$
	$4_f$	<b><math>2_f</math></b>	$7_g$	$8_g$	$4_b$	<b><math>0_f</math></b>	<b><math>2_f</math></b>	$11_g$
	<b><math>4_f</math></b>	<b><math>2_f</math></b>	$7_g$	$8_g$	$4_b$	<b><math>0_f</math></b>	<b><math>2_f</math></b>	$11_g$
	<b><math>4_f</math></b>	<b><math>2_f</math></b>	$7_g$	$8_g$	<b><math>4_b</math></b>	<b><math>0_f</math></b>	<b><math>2_f</math></b>	$11_g$
	<b><math>4_f</math></b>	<b><math>2_f</math></b>	<b><math>7_g</math></b>	$8_g$	<b><math>4_b</math></b>	<b><math>0_f</math></b>	<b><math>2_f</math></b>	$10_c$
	<b><math>4_f</math></b>	<b><math>2_f</math></b>	<b><math>7_g</math></b>	<b><math>8_g</math></b>	<b><math>4_b</math></b>	<b><math>0_f</math></b>	<b><math>2_f</math></b>	$10_c$
	<b><math>4_f</math></b>	<b><math>2_f</math></b>	<b><math>7_g</math></b>	<b><math>8_g</math></b>	<b><math>4_b</math></b>	<b><math>0_f</math></b>	<b><math>2_f</math></b>	<b><math>10_c</math></b>

10. (a) The shortest  $(a, v)$ -paths found by the implementation of Dijkstra's algorithm below are  $a, ab, ac, aefd, ae, aef$ .


v	a	b	c	d	e	f
<b>0a</b>	$4_a$	$8_a$	$\infty_a$	$6_a$	$\infty_d$	
<b>0a</b>	<b>4a</b>	$8_a$	$\infty_a$	$6_a$	$\infty_d$	
<b>0a</b>	<b>4a</b>	$8_a$	$\infty_a$	<b>6a</b>	$12_e$	
<b>0a</b>	<b>4a</b>	<b>8a</b>	$\infty_a$	<b>6a</b>	$12_e$	
<b>0a</b>	<b>4a</b>	<b>8a</b>	$17_f$	<b>6a</b>	<b>12e</b>	
<b>0a</b>	<b>4a</b>	<b>8a</b>	<b>17f</b>	<b>6a</b>	<b>12e</b>	

11. (b) The shortest  $(c, v)$ -paths found by the implementation of Dijkstra's algorithm below are  $ca, cab, c, cd, cgfe, cgf, cg, cdh$ .

v	a	b	c	d	e	f	g	h
$2_c$	$8_c$	<b>0c</b>	$6_c$	$\infty_c$	$\infty_c$	$5_c$	$\infty_c$	
<b>2c</b>	$5_a$	<b>0c</b>	$6_c$	$\infty_c$	$\infty_c$	$5_c$	$\infty_c$	
<b>2c</b>	<b>5a</b>	<b>0c</b>	$6_c$	$19_b$	$15_b$	$5_c$	$\infty_c$	
<b>2c</b>	<b>5a</b>	<b>0c</b>	$6_c$	$19_b$	$9_g$	<b>5c</b>	$9_g$	
<b>2c</b>	<b>5a</b>	<b>0c</b>	<b>6c</b>	$19_b$	$9_g$	<b>5c</b>	$7_d$	
<b>2c</b>	<b>5a</b>	<b>0c</b>	<b>6c</b>	$19_g$	$9_g$	<b>5c</b>	$7_d$	
<b>2c</b>	<b>5a</b>	<b>0c</b>	<b>6c</b>	$12_f$	<b>9g</b>	<b>5c</b>	$7_d$	
<b>2c</b>	<b>5a</b>	<b>0c</b>	<b>6c</b>	<b>12f</b>	<b>9g</b>	<b>5c</b>	$7_d$	

## 7.4 Trees

7. (a) Each of the first five spanning trees shown below can be rotated  $60^\circ$  or  $120^\circ$  to obtain other spanning trees. These  $5 \times 3 = 15$  spanning trees and the sixth spanning tree shown below give 16 spanning trees.



- (b) 100%. Classify the edges as “spokes” (the edges of the sixth spanning tree shown above) and “rim edges”. By the Pigeonhole principle, three edges selected at random must contain at least two spokes or at least two rim edges. In either case, these two edges form a connected subgraph which contains 3 of the 4 vertices of  $G$ . The third edge has two endpoints, and one of them must be already among the three vertices incident on the first two edges. It follows that any three edges selected at random form a connected subgraph.
10. Using the Pythagorean theorem to find the lengths  $BE$  and  $AE$ , we find the lengths of the edges, in increasing order, are as shown:

$$BD\ 3, CE\ 3, DE\ 4, BC\ 4, BE\ 5, AC\ \text{unknown}, AB\ 12, AE\ 13.$$

Applying Kruskal's algorithm, we select edges  $BD, CE$ , either  $DE$  or  $BC$ , and  $AC$ . Since the side  $AC$  of unknown length does appear in the minimal spanning tree, we must compute its length to find the length of the minimal spanning tree. Since angles  $BEC$  and  $ABC$  are both complements of angle  $CBE$ , they have the same measure  $\theta$ . From the 3-4-5 triangle, we see that  $\cos \theta = \frac{3}{5}$ . Applying the law of cosines to triangle  $ABC$ , we have  $AC^2 = 4^2 + 12^2 - 2(4)(12)\frac{3}{5} = 102.4$ , so  $AC = \sqrt{102.4} \approx 10.1193$ . Thus, the length of the minimal spanning tree is approximately  $3 + 3 + 4 + 10.1193 = 20.1193$ .

12. The weights of the edges of graph  $Q$  are given below in increasing order.

FG 5   BD 6   EH 7   FH 7   AB 8   EF 8   DE 9   CF 9  
 CG 9   AD 10   CD 10   GH 10   BE 11   DH 11   BC 12   CH 12

We go through the list and select the following edges to form a minimal spanning tree for  $Q$ : FG, BD, EH, FH, AB, DE, and CF. The only other edges forming a minimal spanning tree for  $G$  are FG, BD, EH, FH, AB, DE, and CG. The weight of the minimal spanning trees is 51.

14. (a) The algorithm for maximal spanning trees is this: Start with any edge of maximal weight. From the remaining edges, add any edge of maximal weight which does not create a cycle. Repeat until all vertices are used. The result is a maximal spanning tree.

We now prove that the algorithm works. Given a connected weighted graph  $G(V, E)$  with weight function  $w : E \rightarrow [0, \infty)$ . Let  $m - 1 = \max\{w(e) | e \in E\}$  be the maximum weight in  $G$ , and define  $G'$  to be the graph  $G'(V, E)$  having the same vertices and edges, but with the new weight function  $w'(e) = m - w(e)$ . Note that a list of the edges of  $G$  in increasing order of weights gives a list of the edges of  $G'$  in decreasing order of weights. Let  $\mathcal{S}$  be the set of all spanning trees for  $G(V, E)$ . Then  $\mathcal{S}$  is also the set of all spanning trees for  $G'(V, E)$ . Each  $T \in \mathcal{S}$  has  $v - 1$  edges where  $v = |V|$ . If  $T = \{e_1, \dots, e_{v-1}\} \in \mathcal{S}$ , then  $w(T) = w(e_1) + \dots + w(e_{v-1})$  and  $w'(T) = (m - w(e_1)) + \dots + (m - w(e_{v-1})) = (v - 1)m - w(T)$ . Thus, for  $T \in \mathcal{S}$ ,  $w(T)$  is maximum when  $w'(T)$  is minimum, and conversely. This shows that a minimal spanning tree for  $G'$  is a maximal spanning tree for  $G$ , and conversely.

- (b) As in the solution to Exercise 11, we list the edges of the graph of Exercise 11 in order.

AF 315   HK 320   CD 330   AB 332   BF 340   GK 345   FJ 350   BC 360  
 CF 360   JK 365   DG 370   FH 375   EI 375   IJ 378   AE 380   EF 380

Since we want a maximal spanning tree, we proceed greedily through the edges from the heaviest backwards through the list to the lightest, including edges as long as they do not create a cycle. The edges required are: EF, AE, IJ, EI, FH, DG, JK, CF, BC, GK. These edges form a maximal spanning tree, and the weight of this tree is  $380 + 380 + 378 + 375 + 375 + 370 + 365 + 360 + 360 + 345 = 3688$ .





# Chapter 8

## Sequences

### 8.1 Sequences

5. (151, 144, 137, 130, 123, 116, 109, 102, 95, 88, 81, 74, 67, 60, 53, 46, 39, 32, 25, 18, 11, 4).  
There are 22 nonnegative terms in this sequence, indicating that 22 is the largest number of sevens which can be subtracted from 158 so that the remaining difference (namely, 4) is nonnegative. This tells us that  $158 \div 7$  gives a quotient of 22 with a remainder of 4.
6. (e)  $(b_n)_{n=1}^{\infty} = (|3n - 4| - 5)_{n=1}^{\infty}$  is a subsequence of  $(a_n)_{n=1}^{\infty}$ . ( $f : \mathbb{R} \rightarrow \mathbb{R}$  is not increasing, but  $f : \mathbb{N} \rightarrow \mathbb{N}$  is.)
11. (b) Suppose  $b_n = b + ns$  and  $a_n = ar^n$  for  $n = 0, 1, 2, \dots$ . Then  $a_{b_n} = ar^{b+ns} = ar^b \cdot r^{ns} = ar^b (r^s)^n$ , so  $(a_{b_n})_{n=0}^{\infty}$  is geometric with first term  $ar^b$  and ratio  $r^s$ .
12. (a) No. For example, if  $(a_n)_{n=1}^{\infty} = (2, 4, 6, 8, 10, \dots)$  and  $(b_n)_{n=1}^{\infty} = (1, 2, 4, 8, \dots)$ , then  $(a_{b_n})_{n=0}^{\infty} = (a_1, a_2, a_4, a_8, \dots) = (2, 4, 8, 16, \dots)$  which is not arithmetic.
16. (a) If  $D$  is a nonempty countable set, then either  $D = \{d_1, d_2, \dots, d_n\}$  is finite or  $D$  is countably infinite. If  $D$  is finite, then

$$(d_1, d_2, \dots, d_{n-1}, d_n, d_n, d_n, d_n, d_n, \dots)$$

is a sequence whose set of terms is  $D$ . If  $D$  is countably infinite, then there exists a bijection  $f$  from  $\mathbb{N}$  to  $D$ , and  $(f(n))_{n=1}^{\infty}$  is a sequence whose set of terms is  $D$ .

- (b) If  $D = \{d_1, d_2, \dots, d_n\}$ , then any sequence in  $D$  is a subsequence of

$$(d_1, d_2, \dots, d_n, d_1, d_2, \dots, d_n, d_1, d_2, \dots, d_n, \dots).$$

If  $D = \{d_1, d_2, d_3, \dots\}$  is countably infinite, then any sequence in  $D$  is a subsequence of

$$(d_1, d_1, d_2, d_1, d_2, d_3, d_1, d_2, d_3, d_4, d_1, d_2, d_3, d_4, d_5, d_1, d_2, \dots).$$

## 8.2 Finite Differences

1. (a) The sequence of second differences is constantly 4. This tells us that the sequence is generated by a second degree polynomial  $p(n) = an^2 + bn + c$ . Since the second differences of the sequence determined by  $p(n)$  are constantly  $2!a = 4$ , we find that  $a = 2$ . Since the first term 4 is  $p(0) = c$ , we have  $p(n) = 2n^2 + bn + 4$ . Now  $p(1) = 3 = 2(1^2) + b(1) + 4$  implies  $b = -3$ , so  $p(n) = 2n^2 - 3n + 4$ .
  - (c) The sequence of third differences is constantly 18. This tells us that the sequence is generated by a third degree polynomial  $p(n) = an^3 + bn^2 + cn + d$ . Since the third differences of the sequence determined by  $p(n)$  are constantly  $3!a = 18$ , we find that  $a = 3$ . Since the first term  $-1$  is  $p(0) = d$ , we have  $p(n) = 3n^3 + bn^2 + cn - 1$ . The equations  $p(1) = 2$  and  $p(2) = 23$  yield, respectively,  $b + c = 0$  and  $2b + c = 0$ , and the only simultaneous solution to these equations is  $b = c = 0$ . Thus,  $p(n) = 3n^3 - 1$ .
  - (e) The sequence of third differences is constantly  $30 = 5 \cdot 3!$  and the initial term is 9, so the sequence is generated by a third degree polynomial of form  $5n^3 + bn^2 + cn + 9$ . The equations  $p(1) = 11$  and  $p(2) = 43$  yield  $b + c = -3$  and  $2b + c = -3$ , giving  $b = 0$  and  $c = -3$ . Thus,  $p(n) = 5n^3 - 3n + 9$ .
  - (g) The sequence of fourth differences is constantly  $48 = 2 \cdot 4!$  and the initial term is 38, so the sequence is generated by a polynomial of form  $p(n) = 2n^4 + bn^3 + cn^2 + dn + 38$ . The equations  $p(1) = 40$ ,  $p(2) = 70$ , and  $p(3) = 200$  yield  $b + c + d = 0 = 4b + 2c + d = 9b + 3c + d$ . Clearly  $b = c = d = 0$  is a solution, so  $p(n) = 2n^4 + 38$ .
3. (c)  $a_n = f(n) = n^2 + 2^n$ . Observe that the  $3^{rd}$  differences (and all  $m^{th}$  differences for  $m \geq 3$ ) are  $1, 2, 4, 8, \dots$ . The sequence  $(2^n)_{n=0}^{\infty}$  has  $1, 2, 4, 8, \dots$  as  $m^{th}$  differences for all natural numbers  $m$ . The fact that the first and second differences of our sequence are not  $1, 2, 4, 8, \dots$  suggests that the terms of our sequence are  $2^n + p(n)$  where  $p(n)$  is a second degree polynomial. (The addition of such a polynomial will alter only the first and second differences, since all subsequent differences of  $p(n)$  would be zero.) Subtracting  $2^n$  from the  $n^{th}$  term of the original sequence leaves the sequence  $n^2$ , so the original sequence is given by  $a_n = n^2 + 2^n$ .
  5. If  $a_n = f(n) = \sum_{i=0}^n i^2$ , then the sequence of first differences is  $(0^2, 1^2, 2^2, 3^2, \dots)$  and thus the sequence of third differences is constantly  $2! = 2$ . Thus,  $a_n = f(n) = an^3 + bn^2 + cn + d$ , and since the third differences of this sequence are  $3!a = 2$ , we have  $a = \frac{2}{3!} = \frac{1}{3}$ . Since  $f(0) = 0 = d$ , we now have  $f(n) = \frac{1}{3}n^3 + bn^2 + cn$ . From  $f(1) = 1^2 = \frac{1}{3} + b + c$  and  $f(2) = 1^2 + 2^2 = 5 = \frac{8}{3} + 4b + 2c$ , we find that  $b = \frac{1}{2}$  and  $c = \frac{1}{6}$ , so

$$f(n) = 1^2 + \dots + n^2 = \frac{n^3}{3} + \frac{n^2}{2} + \frac{n}{6} = \frac{n}{6}(2n^2 + 3n + 1) = \frac{n(n+1)(2n+1)}{6}.$$

7. If the  $k^{th}$  differences of  $(a_i)_{i=1}^{\infty}$  are generated by an  $n^{th}$  degree polynomial, then the  $n^{th}$  differences of the  $k^{th}$  differences of  $(a_i)_{i=1}^{\infty}$  are a nonzero constant. Thus, the  $(n+k)^{th}$  differences of  $(a_i)_{i=1}^{\infty}$  are constant and nonzero, so  $(a_i)_{i=1}^{\infty}$  is generated by an  $(n+k)^{th}$  degree polynomial.

12. (b) The first differences of  $-5, -2, 4, 16, 40, \dots$  agree with those of  $(3 \cdot 2^i)_{i=0}^\infty = (3, 6, 12, 24, 48, \dots)$ , so these two sequences differ by a constant. The original sequence is  $(3 \cdot 2^i - 8)_{i=0}^\infty$ .

### 8.3 Limits of Sequences of Real Numbers

3. Given  $\epsilon > 0$ , we wish to find  $N \in \mathbb{N}$  such that

$$n \geq N \Rightarrow \left| \frac{1-n^2}{3n^2+1} - \frac{-1}{3} \right| < \epsilon.$$

But

$$\left| \frac{1-n^2}{3n^2+1} - \frac{-1}{3} \right| = \left| \frac{3-3n^2}{3(3n^2+1)} + \frac{3n^2+1}{3(3n^2+1)} \right| = \left| \frac{4}{3(3n^2+1)} \right| = \frac{4}{9n^2+3}.$$

Choose  $N \in \mathbb{N}$  such that  $\frac{1}{N} < \epsilon$ .

$$\begin{aligned} n \geq N &\Rightarrow 9n^2 + 3 \geq 4N \\ &\Rightarrow \frac{4}{9n^2 + 3} \leq \frac{4}{4N} = \frac{1}{N} < \epsilon. \end{aligned}$$

Thus,  $n \geq N \Rightarrow \left| \frac{1-n^2}{3n^2+1} - \frac{-1}{3} \right| < \epsilon$ , as needed.

[Or, choose  $N \in \mathbb{N}$  such that  $\frac{4}{9N^2+3} < \epsilon$ , if you believe such an  $N$  exists.]

5. Given  $M < 0$ , we wish to find  $N \in \mathbb{N}$  such that  $n \geq N \Rightarrow \frac{2n^2+1}{3-n} < M$ . Choose  $N = \max\{4, -M\}$  and suppose  $n \geq N$ . Now  $n \geq N \Rightarrow n \geq 4$ , which implies

$$\begin{aligned} \frac{2n^2+1}{3-n} &= \frac{2n^2}{3-n} + \frac{1}{3-n} \\ &< \frac{2n^2}{3-n} = \frac{-n(-2n)}{3-n} = -n \left( \frac{2n}{n-3} \right) = -n \left( \frac{n+n}{n-3} \right) \\ &< -n \quad (\text{since } \frac{n+n}{n-3} > 1). \end{aligned}$$

Now because  $-n \leq -N \leq M$ , we have  $n \geq N$  now implies  $\frac{2n^2+1}{3-n} < M$ , as needed.

8. The functions that preserve all limits are known as *continuous functions*. Our example will necessarily be discontinuous. Let  $f(x) = 1$  if  $x \neq 0$  and  $f(0) = 0$ . Let  $(a_n)_{n=1}^\infty = (\frac{1}{n})_{n=1}^\infty$ . Now  $f(a_n) = f(\frac{1}{n}) = 1$  for any  $n \in \mathbb{N}$ , so  $\lim_{n \rightarrow \infty} f(a_n) = \lim_{n \rightarrow \infty} 1 = 1$ , but  $f(\lim_{n \rightarrow \infty} a_n) = f(\lim_{n \rightarrow \infty} \frac{1}{n}) = f(0) = 0 \neq 1 = \lim_{n \rightarrow \infty} f(a_n)$ .
12. (b) The following statements are equivalent:
- i.  $\lim_{n \rightarrow \infty} a_n = \infty$
  - ii.  $\forall M > 0 \exists N \in \mathbb{N}, N > 100$  such that  $n \geq N \Rightarrow a_n > M$
  - iii.  $\forall M > 0 \exists N \in \mathbb{N}, N > 100$  such that  $n + 100 \geq N \Rightarrow a_{n+100} > M$

- iv.  $\forall M > 0 \exists N' = N - 100 \in \mathbb{N}$  such that  $n \geq N' \Rightarrow b_n > M$   
 v.  $\lim_{n \rightarrow \infty} b_n = \infty$

14. Suppose  $\lim_{n \rightarrow \infty} a_n = A$  and  $\lim_{n \rightarrow \infty} b_n = B$ . Suppose  $\epsilon > 0$  is given. Then there exist  $N_a, N_b \in \mathbb{N}$  such that  $n \geq N_a \Rightarrow |a_n - A| < \frac{\epsilon}{2}$  and  $n \geq N_b \Rightarrow |b_n - B| < \frac{\epsilon}{2}$ . Now for  $n \geq \max\{N_a, N_b\}$ , we have

$$\begin{aligned} |A_n + b_n - (A + B)| &= |a_n - A + b_n - B| \\ &\leq |a_n - A| + |b_n - B| \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon. \end{aligned}$$

Thus,  $\lim_{n \rightarrow \infty} (a_n + b_n) = A + B = \lim_{n \rightarrow \infty} a_n + \lim_{n \rightarrow \infty} b_n$ .

## 8.4 Some Convergence Properties

1. (a) False. Consider  $a_n = \frac{1}{n}, b_n = \frac{1}{2n}$ .  
 (b) True. If  $\lim_{n \rightarrow \infty} a_n = A < B = \lim_{n \rightarrow \infty} b_n$ , take  $\epsilon = \frac{B-A}{2}$ . Now there exists  $M_a, M_b \in \mathbb{N}$  such that

$$A - \epsilon < a_n < A + \epsilon \leq B - \epsilon < b_j < B + \epsilon$$

for any  $n \geq M_a$  and any  $j \geq M_b$ . Now for  $M = \max\{M_a, M_b\}$ , we have  $a_n < b_n \forall n \geq M$ .

6. Any decreasing sequence  $(a_n)_{n=1}^{\infty}$  which is not bounded below by any  $M$  must diverge to  $-\infty$ , for given  $M < 0, \exists N \in \mathbb{N}$  such that  $a_N < M$ , and therefore  $a_n \leq M \forall n \geq N$ . If  $(a_n)_{n=1}^{\infty}$  is a decreasing sequence of real numbers which is bounded below, then  $(-a_n)_{n=1}^{\infty}$  is an increasing sequence of real numbers bounded above, and therefore  $(-a_n)_{n=1}^{\infty}$  converges to a limit  $-L$  by the proof of Theorem 8.4.1. It follows that  $(a_n)_{n=1}^{\infty}$  converges to  $L$ .

Thus, any decreasing sequence of real numbers either converges or diverges to  $-\infty$ .

9. (a) Dividing the numerator and denominator of the expression for  $a_n$  by  $n^2$  gives

$$a_n = \frac{1 - \frac{100}{n}}{1 + \frac{2}{n^2}} = \frac{p(\frac{1}{n})}{q(\frac{1}{n})} \quad \text{where } p(x) = 1 - 100x \quad \text{and } q(x) = 1 + 2x^2.$$

Similarly, we find  $r(x) = 1 + 100x$  and  $s(x) = 1 + 2x^2$ .

$$(b) \quad \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} \frac{p(\frac{1}{n})}{q(\frac{1}{n})} = \frac{\lim_{n \rightarrow \infty} p(\frac{1}{n})}{\lim_{n \rightarrow \infty} q(\frac{1}{n})} = \frac{p(0)}{q(0)} = \frac{1}{1} = 1,$$

and similarly,  $\lim_{n \rightarrow \infty} b_n = 1$ .

- (c) Since  $a_n \leq c_n \leq b_n \forall n \in \mathbb{N}$  and the outer two sequences converge to 1 as  $n \rightarrow \infty$ , it follows that  $\lim_{n \rightarrow \infty} c_n = 1$ .

## 8.5 Infinite Arithmetic

2.  $1 - 1 + 1 - 1 + 1 - \dots$  diverges. The odd partial sums are all 1 and the even partial sums are all 0, so the sequence  $(1, 0, 1, 0, 1, 0, \dots)$  of partial sums does not converge.
4.  $d_n = 0.\underbrace{111\dots1}_{n \text{ digits}}$  and  $\lim_{n \rightarrow \infty} d_n = 0.\overline{111} = 1$ . (Just as  $0.999 = 1.0$  in base 10, in base 2 we have  $0.\overline{111} = 1.0$ .)
9. (c) Let  $P_n = \prod_{j=0}^n \left(1 + \frac{1}{r^{2^j}}\right)$  and  $s_n = \sum_{j=0}^n r^j$ . Note that

$$\begin{aligned} P_0 &= 1 + \frac{1}{r} = s_1 \\ P_1 &= \left(1 + \frac{1}{r}\right) \left(1 + \frac{1}{r^2}\right) = 1 + \frac{1}{r} + \frac{1}{r^2} + \frac{1}{r^3} = s_3 \\ P_2 &= P_1 \left(1 + \frac{1}{r^4}\right) = s_7 \\ P_3 &= s_{15} \end{aligned}$$

and in general,  $P_n = s_{2^{n+1}-1}$ .

Now if  $r \in (0, 1)$ , then  $\sum_{j=0}^{\infty} r^j$  converges, and this implies the convergence of  $(s_n)_{n=0}^{\infty}$  and thus  $(s_{2^{n+1}-1})_{n=0}^{\infty} = (P_n)_{n=0}^{\infty}$ . If  $r \geq 1$ , then  $\sum_{j=0}^{\infty} r^{(2^j)}$  diverges since  $\lim_{j \rightarrow \infty} r^{(2^j)} \neq 0$ , and by Theorem 8.5.4,  $\lim_{n \rightarrow \infty} P_n$  also diverges. Thus, for  $r > 0$ ,  $\sum_{j=0}^{\infty} r^j = \prod_{j=0}^{\infty} \left(1 + \frac{1}{r^{(2^j)}}\right)$ .

10. Let  $p_k$  be the  $k^{\text{th}}$  partial product.
- (b)  $(p_{100k})_{k=1}^{10} = (0.6667326, 0.6666832, 0.6666740, 0.6666708, 0.6666693, 0.6666685, 0.6666680, 0.6666677, 0.6666674, 0.6666673)$ . This suggests that the partial products decrease to  $\frac{2}{3}$ .
11. (b) As  $n \rightarrow \infty$ , the graphs of  $f_n(x)$  converge to the graph of  $y = \cos(x)$ .
13. (a) Let  $p_k = \sqrt{3 + \sqrt{2 + \dots + \sqrt{a_k}}}$  where  $(a_i)_{i=1}^{\infty} = (3, 2, 3, 2, 3, 2, \dots)$ . Now  $p_{k+2} = \sqrt{3 + \sqrt{2 + p_k}}$ . Observe that  $p_1 = \sqrt{3} < 3$  and  $p_2 = \sqrt{3 + \sqrt{2}} < \sqrt{3 + 2} < \sqrt{9} = 3$ . Now suppose  $p_1, \dots, p_{k+1} < 3$ . Then  $p_{k+2} = \sqrt{3 + \sqrt{2 + p_k}} < \sqrt{3 + \sqrt{2 + 3}} = \sqrt{3 + \sqrt{5}} < \sqrt{3 + 5} < \sqrt{9} = 3$ , since  $g(x) = \sqrt{3 + \sqrt{2 + x}}$  is an increasing function. By mathematical induction,  $(p_k)_{k=1}^{\infty}$  is bounded above by 3.
17. Any periodic sequence of nonnegative real numbers is bounded above, and thus the sequence of partial expressions for the associated infinite additive nested radical is increasing and bounded above, and hence is convergent.
18. (a) If  $\sqrt{n + \sqrt{n + \sqrt{n + \dots}}} = 3$ , then  $n + \sqrt{n + \sqrt{n + \sqrt{n + \dots}}} = 9$ , or  $n + 3 = 9$ , so  $n = 6$ .

## 8.6 Recurrence Relations

2. Given a  $k^{\text{th}}$ -order recurrence relation and  $k$  initial conditions  $a_1, \dots, a_k$ , this uniquely determines  $a_{k+1}$ . Now suppose  $a_{j-k+1}, a_{j-k+2}, \dots, a_j$  have been uniquely determined. The recurrence relation then gives  $a_{j+1}$ . By mathematical induction, we see that  $a_n$  is uniquely determined for any  $n \in \mathbb{N}$ , and thus  $f(n) = a_n$  is the unique solution to the recurrence relation.
5. The Fibonacci sequence is given by  $F_0 = 0, F_1 = 1$ , and  $F_{n+2} = F_{n+1} + F_n \forall n \in \mathbb{N} \cup \{0\}$ . The characteristic equation  $r^2 = r + 1$  has roots  $r = \frac{1 \pm \sqrt{5}}{2}$ , which provide the basic solutions  $((\frac{1+\sqrt{5}}{2})^n)_{n=0}^\infty$  and  $((\frac{1-\sqrt{5}}{2})^n)_{n=0}^\infty$  to the recurrence relation. We wish to find a linear combination  $c(\frac{1+\sqrt{5}}{2})^n + d(\frac{1-\sqrt{5}}{2})^n$  which satisfies the initial conditions:

$$\begin{aligned} 0 = F_0 &= c + d && \text{(so } d = -c) \\ 1 = F_1 &= c \left( \frac{1 + \sqrt{5}}{2} \right) + d \left( \frac{1 - \sqrt{5}}{2} \right) \\ &= c \left( \frac{1 + \sqrt{5}}{2} \right) - c \left( \frac{1 - \sqrt{5}}{2} \right) \\ &= \sqrt{5}c \end{aligned}$$

It follows that  $c = \frac{1}{\sqrt{5}}$  and  $d = \frac{-1}{\sqrt{5}}$ , so

$$F_n = \frac{(\frac{1+\sqrt{5}}{2})^n - (\frac{1-\sqrt{5}}{2})^n}{\sqrt{5}} \quad \forall n \in \mathbb{N} \cup \{0\}.$$

6. (a) The recurrence relation  $a_{n+2} = 3a_{n+1} + 10a_n$  has characteristic equation  $r^2 - 3r - 10 = 0 = (r - 5)(r + 2)$ , so the general solution to the recurrence relation is  $a_n = b5^n + c(-2)^n$ . The initial condition  $a_0 = -2$  gives  $-2 = b + c$  and the initial condition  $a_1 = 11$  gives  $11 = 5b - 2c$ . These two linear equations in  $b$  and  $c$  have a unique solution  $b = 1, c = -3$ . Thus,  $a_n = 1 \cdot 5^n - 3(-2)^n = 5^n - 3(-2)^n$ .
- (c) The recurrence relation  $a_{n+4} = 13a_{n+2} - 36a_n$  has characteristic equation  $r^4 - 13r^2 + 36 = 0 = (r^2 - 9)(r^2 - 4)$ , which has roots  $\pm 3, \pm 2$ , so the general solution to the recurrence relation is  $a_n = b3^n + c(-3)^n + d2^n + e(-2)^n$ . The initial conditions  $a_0 = 14, a_1 = -5, a_2 = 101$  and  $a_3 = -35$  give

$$\begin{aligned} b + c + d + e &= 14 \\ 3b - 3c + 2d - 2e &= -5 \\ 9b + 9c + 4d + 4e &= 101 \\ 27b - 27c + 8d - 8e &= -35 \end{aligned}$$

This system may be solved using standard linear algebra techniques, or we may reduce this system of 4 equations in 4 unknowns to two systems of 2 equations

in 2 unknowns: The first and third equations form a system in unknowns  $(b + c)$  and  $(d + e)$

$$\begin{aligned}(b + c) + (d + e) &= 14 \\ 9(b + c) + 4(d + e) &= 101\end{aligned}$$

with solutions  $b + c = 9, d + e = 5$ . The second and fourth equations form a system in unknowns  $(b - c)$  and  $(d - e)$

$$\begin{aligned}3(b - c) + 2(d - e) &= -5 \\ 27(b - c) + 8(d - e) &= -35\end{aligned}$$

with solutions  $b - c = -1, d - e = -1$ . Now combining  $b + c = 9$  and  $b - c = -1$ , we find  $b = 4, c = 5$ , and combining  $d + e = 5$  and  $d - e = -1$ , we find  $d = 2, e = 3$ . Thus, the solution to the recurrence relation with the given initial conditions is  $a_n = 4 \cdot 3^n + 5(-3)^n + 2 \cdot 2^n + 3(-2)^n$ .

8. (a) The characteristic equation is  $r^2 = 4r - 4$  or  $(r - 2)^2 = 0$ , so  $r = 2$  is a repeated root of multiplicity 2.
- (b) Substituting  $a_n = 2^n$  into the recurrence relation, we get  $2^{n+2} = 4 \cdot 2^{n+1} - 4 \cdot 2^n$ , or upon dividing by  $2^n$ ,  $2^2 = 4 \cdot 2 - 4$ , which is true. Substituting  $a_n = n2^n$  into the recurrence relation, we get  $(n + 2)2^{n+2} = 4(n + 1)2^{n+1} - 4n2^n$ , or upon dividing by  $4 \cdot 2^n$ ,  $(n + 2) = (n + 1)2 - n$ , which is true. Now by Theorem 8.6.2,  $a_n = c2^n + dn2^n$  is a solution to the recurrence relation.
- (c) The initial conditions give  $5 = a_0 = c + 0d$  and  $-4 = a_1 = 2c + 2d$ , so  $c = 5$  and  $d = -7$ , and thus  $a_n = 5(2^n) - 7n(2^n) = 2^n(5 - 7n) \forall n \geq 0$ .





## Chapter 9

# Fibonacci Numbers and Pascal's Triangle

### 9.1 Pascal's Triangle

3. (a) 

Ways to write 4 as an ordered sum of natural numbers		
using one term	4	1 way
using two terms	1+3 = 3+1 = 2+2	3 ways
using three terms	1+1+2 = 1+2+1 = 2+1+1	3 ways
using four terms	1+1+1+1	1 way

There are  $8 = \sum_{j=0}^3 \binom{3}{j}$  solutions. By the results of Section 4.4, the number of natural number solutions to  $x_1 + \cdots + x_k = 4$  is the same as the number of whole number solutions to  $x'_1 + \cdots + x'_k = 4 - k$ , which will be  $\binom{4-k+k-1}{4-k} = \binom{3}{4-k}$ . Summing from  $k = 1$  to 4 gives the number we wish, namely  $\sum_{k=1}^4 \binom{3}{4-k} = \sum_{j=0}^3 \binom{3}{j} = 2^3 = 8$ .

- (b) The number of natural number solutions to  $x_1 + \cdots + x_k = m$  is the same as the number of whole number solutions to  $x'_1 + \cdots + x'_k = m - k$ , and this number is  $\binom{m-k+k-1}{m-k} = \binom{m-1}{m-k}$ . Summing from  $k = 1$  to  $m$  gives the number we wish, namely

$$\sum_{k=1}^m \binom{m-1}{m-k} = \sum_{j=0}^{m-1} \binom{m-1}{j} = 2^{m-1}.$$

The last equality holds from the result of Example 9.1.1.

5. Of the  $2^4 = 1 + 4 + 6 + 4 + 1 = 16$  subsets of  $\{a, b, c, d\}$ , half of them ( $2^3 = 1 + 6 + 1 =$

$\binom{4}{0} + \binom{4}{2} + \binom{4}{4} = 8$  of them) have an even number of elements. These subsets are  $\emptyset, \{a, b\}, \{a, c\}, \{a, d\}, \{b, c\}, \{b, d\}, \{c, d\}$ , and  $\{a, b, c, d\}$ .

10. The seven entries around  $\binom{n}{k}$  are

$$\begin{array}{ccccc} & & \binom{n-1}{k-1} & & \binom{n-1}{k} & & \\ & & & & & & \\ \binom{n}{k-1} & & \binom{n}{k} & & \binom{n}{k+1} & & \\ & & & & & & \\ & & \binom{n+1}{k} & & \binom{n+1}{k+1} & & \end{array} \quad \text{which we label as } \begin{array}{ccccc} & & a & & b & & \\ & & c & d & e & & \\ & & & f & g & & \end{array} .$$

Let  $h = \binom{n+2}{k+1}$ . We wish to show that  $a + b + c + d + e + f + g = 2h$ . Now

$$\begin{aligned} a + b + c + d + e + f + g &= [(a + b) + c] + (d + e) + (f + g) \\ &= [d + c] + (g) + (h) \\ &= f + g + h \\ &= 2h, \quad \text{as needed.} \end{aligned}$$

13. (a)  $\left( \binom{n}{0}, \binom{n}{1}, \binom{n}{2}, \dots, \binom{n}{n} \right) \cdot (0, 1, 2, \dots, n) = 2^{n-1} \cdot n$ .

14. (a) 
$$\begin{aligned} &\left( \binom{n}{0}, \binom{n}{1}, \binom{n}{2}, \dots, \binom{n}{n} \right) \cdot (0, 1, 2, \dots, n) \\ &= \binom{n}{1} + 2 \binom{n}{2} + 3 \binom{n}{3} + \dots + (n-1) \binom{n}{n-1} + n \binom{n}{n} \\ &= n + \frac{n!}{(n-2)!1!} + \frac{n!}{(n-3)!2!} + \dots + \frac{n!}{1!(n-2)!} + n \\ &= n \left[ 1 + \binom{n-1}{1} + \binom{n-1}{2} + \dots + \binom{n-1}{n-2} + 1 \right] \\ &= n \cdot 2^{n-1} \text{ (by the result of Example 9.1.1).} \end{aligned}$$

17. (a) Observe that row  $k$  contains only odd entries if and only if row  $k + 1$  contains only even entries except for the initial and final 1's. Thus, by Theorem 9.1.8, the rows which contain only odd entries are rows  $2^m - 1$  for  $m \in \mathbb{N} \cup \{0\}$ .


(b) The entries of row  $m$  alternate odd, even, odd, even,  $\dots$  if and only if the entries of row  $m + 1$  are all odd, and by part (a), this occurs if and only if  $m + 1 = 2^n - 1$  ( $n \in \mathbb{N}$ ), if and only if  $m = 2^n - 2$  for some  $n \in \mathbb{N}$ .

## 9.2 The Fibonacci Numbers


1. (a)  $F_7 = 13$ . See Example 9.2.2, and interpret  $1''$  high blocks as \$5 payments and  $2''$  high blocks as \$10 payments.
- (b)  $F_{12} = 144$ .
- (c)  $F_{n+1}$ .

5. (d)

$$F_{n+1} + F_{n-2} = 2F_n$$



Or, consider the following rectangle with area  $2F_n$ . The shaded region has area  $F_{n+1}$  (since  $F_{n-1} + F_n = F_{n+1}$ ) but also has area  $2F_n - F_{n-2}$ .



6.  $F_0 = 0 = (-1)^1 F_0$  and  $F_{-1} = 1 = (-1)^2 F_1$ . Now suppose  $F_{-k} = (-1)^{k+1} F_k$  for  $k = 0, 1, \dots, j$ . Now

$$\begin{aligned} F_{-(j+1)} &= F_{-(j-1)} - F_{-j} \\ &= (-1)^j F_{j-1} - (-1)^{-j+1} F_j \\ &= (-1)^{j+2} (F_{j-1} + F_j) \\ &= (-1)^{j+2} F_{j+1} \end{aligned}$$

By mathematical induction,  $F_{-n} = (-1)^{n+1} F_n$  for any integer  $n \geq 0$ , and dividing by  $(-1)^{n+1}$  shows that the result holds for all negative integers, as well.

8.  $F_m^2 + F_{m-1}^2 = F_{2m-1}$ . Apply Theorem 9.2.4 with  $n = 2m - 1$  and  $j = m - 1$ .

10. After trying a few values of  $n$ , the formula is easily recognized to be

$$F_n F_{n+1} - F_{n-1} F_{n+2} = (-1)^{n+1}.$$

Dividing by  $(-1)^{n+1}$  gives

$$(-1)^{n+1} F_n F_{n+1} + (-1)^n F_{n-1} F_{n+2} = 1.$$

Observing that  $(-1)^{j+1} F_j = F_{-j}$ , we have

$$F_{-n} F_{n+1} + F_{1-n} F_{n+2} = 1 = F_1 = F_2 = F_{-1}.$$

This formula looks very similar to one proved in Theorem 9.2.4:

$$F_{j+1} F_{m-j} + F_j F_{m-j-1} = F_m.$$

We would hope to find appropriate values of  $m$  and  $j$  which transform the result of Theorem 9.2.4 into the formula we wish to prove. Taking  $m = 2$  and  $j = -n$  gives the result.

12. (b)  $(F_{n+1}^2 - F_{n-1}^2)_{n=2}^7 = (3, 8, 21, 55, 144, 377)$ . The formula is  $F_{n+1}^2 - F_{n-1}^2 = F_{2n}$ . Applying Theorem 9.2.4 with  $m = j$  gives

$$\begin{aligned} F_{2j} &= F_{j+1}F_j + F_jF_{j-1} \\ &= F_j(F_{j+1} + F_{j-1}) \\ &= (F_{j+1} - F_{j-1})(F_{j+1} + F_{j-1}) \\ &= F_{j+1}^2 - F_{j-1}^2. \end{aligned}$$

### 9.3 The Golden Ratio

4. The sequence of partial expressions is

$$(\sqrt{1}, \sqrt{1 - \sqrt{1}}, \sqrt{1 - \sqrt{1 - \sqrt{1}}}, \dots) = (1, 0, 1, 0, 1, 0, \dots),$$

which diverges. [Were one not to notice this divergence, one would be tempted to say the value of the nested radical is  $x$  where  $x = \sqrt{1 - x}$ , so that  $x = \frac{-1 \pm \sqrt{5}}{2}$ . All this shows, however, is that *if* the nested radical converged, its value would be one of those given.]

6. Let  $ABCD$ ,  $M$ ,  $E$ , and  $F$  be as described and take  $AB = 1$ . Then  $BC = 1$  and  $MB = \frac{1}{2}$ , so  $CM = \sqrt{1^2 + (\frac{1}{2})^2} = \frac{\sqrt{5}}{2}$ . Now  $AE = AM + ME = AM + MC = \frac{1}{2} + \frac{\sqrt{5}}{2} = \varphi$ , so  $\frac{AE}{AD} = \frac{\varphi}{1} = \varphi$ , and  $AEFD$  is a golden rectangle.
7. The restrictions  $lwh = 1$ ,  $\sqrt{l^2 + w^2 + h^2} = 2$ , and  $h = 1$  give  $l^2 + w^2 = 3$  and  $l = \frac{1}{w}$ . Substituting the latter equation into the former and multiplying through by  $w^2$  gives  $1 + w^4 = 3w^2$ , a quadratic in  $w^2$  with solutions

$$w^2 = \frac{3 + \sqrt{5}}{2} = 1 + \varphi = \varphi^2$$

and

$$w^2 = \frac{3 - \sqrt{5}}{2} = \frac{2}{3 + \sqrt{5}} = \frac{1}{\varphi^2}.$$

Since  $w$  must be positive, we have  $w = \varphi$  and  $l = \frac{1}{w} = \frac{1}{\varphi}$ , or  $w = \frac{1}{\varphi}$  and  $l = \frac{1}{w} = \varphi$ .

### 9.4 Fibonacci Numbers and the Golden Ratio

1. These problems use the fact that  $\varphi^2 = \varphi + 1$ , and (multiplying by  $\varphi^n$ )  $\varphi^{n+2} = \varphi^{n+1} + \varphi^n$ .

$$\begin{aligned}
 \text{(b)} \quad 2\varphi^4 - 3\varphi^2 - 8 &= 2(\varphi + 1)^2 - 3(\varphi + 1) - 8 \\
 &= 2(\varphi^2 + 2\varphi + 1) - 3\varphi - 3 - 8 \\
 &= 2(\varphi + 1) + 4\varphi + 2 - 3\varphi - 11 \\
 &= 3\varphi - 7
 \end{aligned}$$

$$\begin{aligned}
 \text{(d)} \quad 2\varphi^5 - 3\varphi^4 + 1 &= 2(\varphi^4 + \varphi^3) - 3\varphi^4 + 1 \\
 &= -1\varphi^4 + 2\varphi^3 + 1 \\
 &= -(\varphi^3 + \varphi^2) + 2\varphi^3 + 1 \\
 &= \varphi^3 - \varphi^2 + 1 \\
 &= (\varphi^2 + \varphi) - \varphi^2 + 1 \\
 &= \varphi + 1
 \end{aligned}$$

6. (a) Yes. If  $(a_n)_{n=1}^{\infty}$  and  $(b_n)_{n=1}^{\infty}$  are additive sequences and  $c_n = a_n + b_n$ , then

$$\begin{aligned}
 c_{n+2} &= a_{n+2} + b_{n+2} \\
 &= (a_{n+1} + a_n) + (b_{n+1} + b_n) \\
 &= (a_{n+1} + b_{n+1}) + (a_n + b_n) \\
 &= c_{n+1} + c_n,
 \end{aligned}$$

so  $(c_n)_{n=1}^{\infty}$  is additive as well.

- (b) i.  $(F_{n-1} + F_{n+1})_{n=1}^{\infty} = (1, 3, 4, 7, 11, \dots) = (L_n)_{n=1}^{\infty}$   
 ii.  $(L_{n-1} + L_{n+1})_{n=2}^{\infty} = (5, 10, 15, 25, 40, \dots) = (5F_n)_{n=2}^{\infty}$   
 iii.  $(\varphi^n + (\varphi')^n)_{n=1}^{\infty} = (1, 3, 4, 7, 11, \dots) = (L_n)_{n=1}^{\infty}$   
 iv.  $(\frac{F_{2n}}{F_n})_{n=1}^{\infty} = (1, 3, 4, 7, 11, \dots) = (L_n)_{n=1}^{\infty}$

7. (a) From Exercise 4 (a), we have

$$\begin{aligned}
 \lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} &= \lim_{n \rightarrow \infty} \frac{F_{n-1}a_1 + F_n a_2}{F_{n-2}a_1 + F_{n-1}a_2} \\
 &= \lim_{n \rightarrow \infty} \frac{F_{n-1}(a_1 + \frac{F_n}{F_{n-1}}a_2)}{F_{n-2}(a_1 + \frac{F_{n-1}}{F_{n-2}}a_2)} \\
 &= \varphi \left( \frac{a_1 + \varphi a_2}{a_1 + \varphi a_2} \right) \\
 &= \varphi.
 \end{aligned}$$

11. 
$$\frac{x}{1+x-x^2} = 1x - 1x^2 + 2x^3 - 3x^4 + 5x^5 - \dots + (-1)^{n+1}F_n x^n + \dots$$

## 9.5 Pascal's Triangle and the Fibonacci Numbers

2. (c)	Number of \$50 calculators	Number of \$100 calculators	Number of ways to distribute
	8	0	$\binom{15}{8} = 6435$
	6	1	$\binom{15}{6} \binom{9}{1} = 45045$
	4	2	$\binom{15}{4} \binom{11}{2} = 75075$
	2	3	$\binom{15}{2} \binom{13}{3} = 30030$
	0	4	$\binom{15}{4} = 1365$
			157,950

The total number of outcomes is 157,950, which is not a Fibonacci number. It falls between  $F_{26} = 121,393$  and  $F_{27} = 196,418$ .

4. The formula follows immediately from Theorem 9.5.2 and the fact that  $\binom{m}{j} = \binom{m}{m-j}$ . This new formula would have been suggested by Example 9.5.1 if the right column of the tables there had listed the number of ways to distribute \$50 calculators (rather than the \$100 ones) among those receiving new calculators.
7. (a)  $5^n F_{2n}$   
 (b)  $5^n F_{2n+1}$   
 (c)  $5^{n-1} L_{2n}$   
 (d)  $5^n F_{2n+3}$ .

## Chapter 10

# Continued Fractions

### 10.1 Finite Continued Fractions

2. (a)  $[4; ]$

(c)  $[4; 10]$

(f) 
$$\frac{-23}{7} = -3\frac{2}{7} = -4 + \frac{5}{7} = -4 + \frac{1}{\left(\frac{7}{5}\right)} = -4 + \frac{1}{1 + \frac{2}{5}} = -4 + \frac{1}{1 + \frac{1}{\left(\frac{5}{2}\right)}}$$

$$= -4 + \frac{1}{1 + \frac{1}{2 + \frac{1}{2}}} = [-4; 1, 2, 2]$$

(i) 
$$\frac{-7}{5} = -2 + \frac{1}{\left(\frac{5}{3}\right)} = -2 + \frac{1}{1 + \frac{1}{\left(\frac{3}{2}\right)}} = -2 + \frac{1}{1 + \frac{1}{1 + \frac{1}{2}}} = [-2; 1, 1, 2]$$

3. (b)  $[0; ] = 0$ ,  $[0; 2] = \frac{1}{2}$ ,  $[0; 2, 5] = \frac{5}{11}$ ,  $[0; 2, 5, 4] = \frac{21}{46}$ .

(d)  $[-4; ] = -4$ ,  $[-4; 1] = -3$ ,  $[-4; 1, 1] = \frac{-7}{2}$ ,  $[-4; 1, 1, 1] = \frac{-10}{3}$ ,  $[-4; 1, 1, 1, 2] = \frac{-27}{8}$ .

6. The expressions in (a) both equal  $2\frac{3}{7}$ ; those in (b) both equal  $2\frac{1}{4}$ . The expressions on the left are not regular continued fractions (see the  $-2$  in (a) and the second  $2$  in (b)), so Theorem 10.1.6 does not apply.

10.	$k$	$\frac{k}{14}$	$1 - \frac{k}{14} = \frac{14-k}{14}$
	1	[0;14]	[0;1,13]
	2	[0;7]	[0;1,6]
	3	[0;4,1,2]	[0;1,3,1,2]
	4	[0;3,2]	[0;1,2,2]
	5	[0;2,1,4]	[0;1,1,1,4]
	6	[0;2,3]	[0;1,1,3]
	7	[0;2]	[0;2] = [0;1,1]

For  $0 < k < 7$ , the continued fraction for  $1 - \frac{k}{14}$  is of form  $[0; a_1, a_2, \dots, a_n]$  where  $n \geq 1$ , and the continued fraction for  $\frac{k}{14}$  is  $[0; a_1 + a_2, \dots, a_n]$ . Furthermore, in this notation,  $a_1 = 1$ . For  $n = 7$ , the two representations  $[0; 2] = [0; 1, 1] = [0; 1 + 1]$  for  $\frac{7}{14}$  allow us to apply the pattern in this case as well. The pattern is described in general in Exercise 11.

## 10.2 Convergents of a Continued Fraction

1. No.  $a_0$  is the integral part of  $[a_0; a_1, \dots, a_n]$ . If  $x \notin \mathbb{Z}$ , the integral part of  $-x$  is not the negative of the integral part of  $x$ . For example,  $[2; 3] = \frac{7}{3}$ , so  $-[2; 3] = -\frac{7}{3}$ . The integral part of  $-\frac{7}{3}$  is  $-3$ , so the continued fraction for  $-[2; 3]$  is not of form  $[-2; a_1, \dots, a_n]$ . In fact,  $-[2; 3] = [-3; 1, 2]$  and  $[-2; 3] = -\frac{5}{3}$ .
2. (a)  $C_0 = 0, C_1 = 1, C_2 = \frac{1}{2}, C_3 = \frac{3}{5}, C_4 = \frac{7}{12}, C_5 = \frac{10}{17}, C_6 = \frac{17}{29}, C_7 = \frac{44}{75}, C_8 = \frac{105}{179}$   
 (b)  $C_0 = 2, C_1 = 3, C_2 = \frac{11}{4}, C_3 = \frac{14}{5}, C_4 = \frac{67}{24}, C_5 = \frac{81}{29}, C_6 = \frac{472}{169}, C_7 = \frac{553}{198}$
3. Since  $q_k = a_k q_{k-1} + q_{k-2}$ , the  $q_k$ 's will increase most slowly if every  $a_k = 1$ .

$$([1; 1], [1; 1, 1], \dots, [1; 1, 1, 1, 1, 1, 1]) = \left( \frac{2}{1}, \frac{3}{2}, \frac{5}{3}, \frac{8}{5}, \frac{13}{8}, \frac{21}{13} \right).$$

Each expression is of form  $\frac{F_{n+2}}{F_{n+1}}$  where  $F_n$  is the  $n^{\text{th}}$  Fibonacci number.

7. (a)  $\frac{225}{157} = \frac{p_4}{q_4} = [1; 2, 3, 4, 5]; [1; 2, 3, 4] = \frac{43}{30} = \frac{p_3}{q_3}; [1; 2, 3] = \frac{10}{7} = \frac{p_2}{q_2}; [1; 2] = \frac{3}{2} = \frac{p_1}{q_1}; [1; ] = 1$ .  
 (b)  $\frac{157}{30} = [5; 4, 3, 2] = \frac{q_4}{q_3} \cdot \frac{30}{7} = [4; 3, 2] = \frac{q_3}{q_2} \cdot \frac{7}{2} = [3; 2] = \frac{q_2}{q_1} \cdot \frac{2}{1} = [2; ] = \frac{q_1}{q_0}$ .  
 (c) Each is of form  $\frac{q_k}{q_{k-1}}$ .

## 10.3 Infinite Continued Fractions

$$3. \quad 1 + \frac{1}{1 + \sqrt{2}} = \frac{(1 + \sqrt{2}) + 1}{1 + \sqrt{2}} = \frac{2 + \sqrt{2}}{1 + \sqrt{2}} \left( \frac{1 - \sqrt{2}}{1 - \sqrt{2}} \right) = \frac{-\sqrt{2}}{-1} = \sqrt{2}.$$

Putting this expression for  $\sqrt{2}$  in place of the  $\sqrt{2}$  appearing on the left gives

$$\sqrt{2} = 1 + \frac{1}{1 + 1 + \frac{1}{1 + \sqrt{2}}} = 1 + \frac{1}{2 + \frac{1}{1 + \sqrt{2}}}$$



Repeating this gives  $\sqrt{2} = [1; 2, 2, 2, \dots] = [1; \overline{2}]$ .

5. (a) If  $r \neq 0$  is a root of  $p(x) = ax^2 + bx + c$  ( $a, b, c \in \mathbb{Z}$ ) and  $k \in \mathbb{Z}$ , then translating  $p(x)$  by  $k$  units to the right gives a parabola with zero at  $r + k$ . That is,  $r + k$  is a root of the polynomial  $p(x - k)$ . Furthermore, if  $ar^2 + br + c = 0$  and  $r \neq 0$ , dividing by  $r^2$  gives  $a + b(\frac{1}{r}) + c(\frac{1}{r})^2 = 0$ , so  $\frac{1}{r}$  is a root of  $q(x) = cx^2 + bx + a$ .
- (b) Given a periodic continued fraction  $[a_0; a_1, \dots, a_{k-1}, \overline{a_k, \dots, a_{k+j}}]$ , Exercise 4 shows that  $r_1 = [0; \overline{a_k, \dots, a_{k+j}}]$  is a root of a quadratic equation with integer coefficients. Now  $r_2 = [a_{k-1}; \overline{a_k, \dots, a_{k+j}}] = a_{k-1} + \frac{1}{r_1}$  is also a root of a quadratic equation with integer coefficients by an application of part (a). Similarly,

$$r_3 = [a_{k-2}; a_{k-1}, \overline{a_k, \dots, a_{k+j}}] = a_{k-2} + \frac{1}{r_2}$$

is also a root of a quadratic equation with integer coefficients. Continuing this iterative process, we find that the original continued fraction  $[a_0; a_1, \dots, a_{k-1}, \overline{a_k, \dots, a_{k+j}}] = r_{k+1}$  is a root of a quadratic equation with integer coefficients.

## 10.4 Applications of Continued Fractions

1. (a) Since 17 and 13 are relatively prime, Theorem 10.4.1 tells us that all solutions of the Diophantine equation  $17x + 13y = 981$  have form  $(21 + 13j, 48 - 17j)$  for  $j \in \mathbb{Z}$ . Four other solutions may be found by taking  $j = 1, -1, 2$ , and  $10$ , giving solutions  $(34, 31), (8, 65), (47, 14)$ , and  $(151, -122)$ .
2. (a)  $26x + 53y = 3938$ : Since 26 and 53 are relatively prime, we find  $\frac{26}{53} = [0; 2, 26]$ , a continued fraction of order  $n = 2$  with convergent  $C_{n-1} = C_1 = \frac{1}{2}$ . By Theorem 10.4.2, the solutions  $(x, y)$  have form

$$((-1)^1 3938(2) + 53k, (-1)^2 3938(1) - 26k) = (53k - 7876, 3938 - 26k).$$

As we want positive solutions,  $x = 53k - 7876 > 0$  implies  $k > \frac{7876}{53} \approx 148.6$ , and  $y = 3938 - 26k > 0$  implies  $k < \frac{3938}{26} \approx 151.5$ . Thus,  $k = 149, 150$ , or  $151$ , yielding solutions  $(x, y) = (21, 64), (74, 38)$ , and  $(127, 12)$ .

- (d)  $213x + 121y = 6714$ : Since 213 and 121 are relatively prime, we find  $\frac{213}{121} = [1; 1, 3, 5, 1, 4]$ , a continued fraction of order  $n = 5$  with convergent  $C_{n-1} = C_4 = \frac{44}{25}$ . By Theorem 10.4.2, the solutions have form

$$\begin{aligned} (x, y) &= ((-1)^4 6714(25) + 121k, (-1)^5 6714(44) - 213k) \\ &= (167850 + 121k, -295416 - 213k). \end{aligned}$$

As we want positive solutions,  $x = 167850 + 121k > 0$  implies  $k > \frac{-167850}{121} \approx -1387.2$ , and  $y = -295416 - 213k > 0$  implies  $k < \frac{-295416}{213} \approx -1386.9$ . Thus,  $k = -1387$  and the only solution is  $(x, y) = (23, 15)$ .

5. Each congruence given has form  $ax \equiv c \pmod{b}$  where  $a$  and  $b$  are relatively prime, so by Theorem 10.4.4, the solution is of form  $x = (-1)^{n-1} c q_{n-1} + bk$  where  $k \in \mathbb{Z}$ ,  $\frac{a}{b} = [a_0; a_1, \dots, a_n]$ , and  $[a_0; a_1, \dots, a_{n-1}] = \frac{p_{n-1}}{q_{n-1}}$ .

- (a) For  $25x \equiv 18 \pmod{26}$ , we find  $\frac{25}{26} = [0; 1, 25]$ , a continued fraction of order  $n = 2$  and with  $C_{n-1} = C_1 = 1$ , so  $q_1 = 1$  and all solutions are of form  $(-1)^1 18(1) + 26k = -18 + 26k = 8 + 26k'$  ( $k, k' \in \mathbb{Z}$ ). Thus, all solutions are congruent to 8 modulo 26.
7. Let  $x$  be the number of minutes the press ran. Then  $28x$  newspapers were printed. Because enough newsprint for 65 newspapers remained on the last spool, the press had printed  $235 - 65 = 170$  newspapers from its last spool. Thus, the number of newspapers printed is congruent to 170 modulo 235. That is,  $28x \equiv 170 \pmod{235}$ . Since  $28 = 2^2 \cdot 7$  and  $235 = 5 \cdot 47$  are relatively prime, Theorem 10.4.4 applies. We find that  $\frac{28}{235} = [0; 8, 2, 1, 1, 5]$ , a continued fraction of order  $n = 5$ . The first two convergents of this continued fraction are  $C_0 = \frac{0}{1}$  and  $C_1 = \frac{1}{8}$ , so  $q_0 = 1$  and  $q_1 = 8$ . From the recurrence relation  $q_k = a_k q_{k-1} + q_{k-2}$ , we find that  $q_2 = 2(8) + 1 = 17$ ,  $q_3 = 1(17) + 8 = 25$ , and  $q_4 = q_{n-1} = 1(25) + 17 = 42$ . Since the solutions are congruent modulo 235 to  $(-1)^{n-1} c q_{n-1} = (-1)^4 (170)(42) = 7140 = 90 + 235(30)$  and only positive answers are possible, the solution set is  $\{90 + 235j | j \in \mathbb{N}\}$ . The operator should not have spent  $90 + 235$  minutes at lunch, so the only possible answer in the appropriate range is 90 minutes.
9. We find that  $C_4 = \frac{134}{35}$  and  $C_5 = \frac{1229}{321}$ . Since all even convergents of  $\alpha$  are below  $\alpha$ ,  $C_4$  is an underestimate and  $|\alpha - C_4| = \alpha - C_4$ . Now Lemma 10.4.6 gives

$$\frac{1}{12,460} = \frac{1}{35(35 + 321)} < \alpha - \frac{134}{35} < \frac{1}{35(321)} = \frac{1}{11,235} \approx 0.000089.$$

It follows that

$$\frac{47,705}{12,460} = \frac{1}{35(35 + 321)} + \frac{134}{35} < \alpha < \frac{134}{35} + \frac{1}{35(321)} = \frac{1229}{321},$$

so  $\alpha \in (\frac{47,705}{12,460}, \frac{1229}{321}) \subset (\frac{134}{35}, \frac{1229}{321}) = (C_4, C_5)$ .

11. (b)  $3\pi = [9; 2, 2, 1, 4, 1, 1, 1, 97, 4, \dots]$  has convergent  $C_5 = \frac{377}{40}$ , and this must be the best approximation to  $3\pi$  by a rational number with denominator  $\leq 40$ .